

Biomembranes

Cumulative Contents

Rapid report

Raspberry vesicles

A.-L. Bernard, M.-A. Guedeau-Boudeville,
 L. Jullien and J.-M. di Meglio (France)

Regular papers

The interaction of α -tocopherol with bilayers of
 1-palmitoyl-2-oleoyl-phosphatidylcholine

X. Wang and P.J. Quinn (United Kingdom)

Mammalian Pex14p: membrane topology and char-
 acterisation of the Pex14p–Pex14p interaction

M.E.M. Oliveira, C. Reguenga, A.M.M. Gouveia,
 C.P. Guimarães, W. Schliebs, W.-H. Kunau,
 M.T. Silva, C. Sá-Miranda and J.E. Azevedo
 (Portugal, Germany)

Evaluation of lipid-based reagents to mediate intra-
 cellular gene delivery

H. Faneca, S. Simões and M.C. Pedrosa de Lima
 (Portugal)

Indanedione spin labelling of Na,K-ATPase

M. Esmann, C.P. Sár, K. Hideg and D. Marsh
 (Denmark, Hungary, Germany)

Viability of *Escherichia coli* after combined osmotic
 and thermal treatment: a plasma membrane impli-
 cation

Y. Mille, L. Beney and P. Gervais (France)

A molecular basis explanation of the dynamic and
 thermal effects of vinblastine sulfate upon dipal-
 mitoylphosphatidylcholine bilayer membranes

H. Maswadeh, C. Demetzos, I. Daliani, I. Kyrikou,
 T. Mavromoustakos, A. Tsortos and G. Nounesis
 (Greece)

Identification of lysine-411 in the human reduced
 folate carrier as an important determinant of sub-
 strate selectivity and carrier function by system-
 atic site-directed mutagenesis

T.L. Witt and L.H. Matherly (USA)

Comparative molecular dynamics simulations of am-
 photericin B–cholesterol/ergosterol membrane
 channels

M. Baginski, H. Resat and E. Borowski (Poland,
 Turkey)

Interfacial tension of bilayer lipid membrane formed
 from phosphatidylethanolamine

A.D. Petelska and Z.A. Figaszewski (Poland)

Transepithelial ultrafiltration and fractal power diffu-
 sion of D-glucose in the perfused rat intestine

G.M. Kochak and S. Mangat (USA)

Functional characterization of the human intestinal
 NaPi-IIb cotransporter in hamster fibroblasts and
Xenopus oocytes

H. Xu, M. Inouye, T. Missey, J.F. Collins and
 F.K. Ghishan (USA)

Cellular and molecular aspects of thiamin uptake by
 human liver cells: studies with cultured HepG2
 cells

H.M. Said, J.C. Reidling and A. Ortiz (USA)

Estimation of the radius of the pores formed by the
Bacillus thuringiensis Cry1C δ -endotoxin in pla-
 nar lipid bilayers

O. Peyronnet, B. Nieman, F. G  n  reux,
 V. Vachon, R. Laprade and J.-L. Schwartz
 (Canada)

A cholesterol-binding and transporting protein from
 rat liver mitochondria

A.M. Campbell, A. Capuano and S.H.P. Chan
 (USA)

Functional analysis of candidate ABC transporter
 proteins for sitosterol transport

C. Albrecht, J.I. Elliott, A. Sardini, T. Litman,
 B. Stieger, P.J. Meier and C.F. Higgins (UK,
 Denmark, Switzerland)

Syngotoxin pore formation and inactivation in hu-
 man red blood cell and model bilayer lipid mem-
 branes

Z. Szab  , P. Gr  f, L.V. Schagina, P.A. Gurnev,
 J.Y. Takemoto, E. M  tyus and K. Blask  
 (Hungary, Russia, USA)

Doxorubicin induces early lipid peroxidation associated with changes in glucose transport in cultured cardiomyocytes <i>S. Hrelia, D. Fiorentini, T. Maraldi, C. Angeloni, A. Bordoni, P.L. Biagi and G. Hakim (Italy)</i>	150	Ion transport across a phospholipid membrane mediated by the peptide trichogin GA IV <i>T.N. Kropacheva and J. Raap (Russia, The Netherlands)</i>	193
Identification of a conserved domain of the HIV-1 transmembrane protein gp41 which interacts with cholesteryl groups <i>N. Vincent, C. Genin and E. Malvoisin (France)</i>	157	Attaching histidine-tagged peptides and proteins to lipid-based carriers through use of metal-ion-chelating lipids <i>G.G. Chikh, W.M. Li, M.-P. Schutze-Redelmeier, J.-C. Meunier and M.B. Bally (Canada, France)</i>	204
Differential localization of glucose transporter isoforms in non-polarized mammalian cells: distribution of GLUT1 but not GLUT3 to detergent-resistant membrane domains <i>T. Sakyo and T. Kitagawa (Japan)</i>	165	Characterization of the macrocyclic carbon suboxide factors as potent Na,K-ATPase and SR Ca-ATPase inhibitors <i>F. Kerek, R. Stimac, H.-J. Apell, F. Freudenmann and L. Moroder (Germany)</i>	213
Phosphatidylserine content is a more important contributor than transmembrane potential to interactions of merocyanine 540 with lipid bilayers <i>I. Waczulikova, M. Rozalski, J. Rievaj, K. Nagyova, M. Bryszewska and C. Watala (Slovak Republic, Poland)</i>	176	Structural changes are induced in human neutrophil cytochrome <i>b</i> by NADPH oxidase activators, LDS, SDS, and arachidonate: intermolecular resonance energy transfer between trisulfopyrenyl-wheat germ agglutinin and cytochrome <i>b</i> ₅₅₈ <i>T.R. Foubert, J.B. Burritt, R.M. Taylor and A.J. Jesaitis (USA)</i>	221
Erythrocyte ghost cell-alkaline phosphatase: construction and characterization of a vesicular system for use in biomineralization studies <i>D.F. Ierardi, J.M. Pizauro and P. Ciancaglini (Brazil)</i>	183	<i>Author Index</i>	232
		<i>Cumulative Contents</i>	234