

Biomembranes Cumulative Contents

Regular papers

- | | | | |
|---|----|---|-----|
| Computational analysis of mutations in the trans-membrane region of Vpu from HIV-1
<i>A. Candler, M. Featherstone, R. Ali, L. Maloney, A. Watts and W.B. Fischer</i> | 1 | Lysolipid incorporation in dipalmitoylphosphatidylcholine bilayer membranes enhances the ion permeability and drug release rates at the membrane phase transition
<i>J.K. Mills and D. Needham</i> | 77 |
| Membrane-disruptive abilities of β -hairpin antimicrobial peptides correlate with conformation and activity: A ^{31}P and ^1H NMR study
<i>R. Mani, A.J. Waring, R.I. Lehrer and M. Hong</i> | 11 | Organization of protochlorophyllide oxidoreductase in prolamellar bodies isolated from etiolated carotenoid-deficient wheat leaves as revealed by fluorescence probes
<i>I.D. Denev, G.T. Yahubyan, I.N. Minkov and C. Sundqvist</i> | 97 |
| Effect of ischemia reperfusion on sodium-dependent phosphate transport in renal brush border membranes
<i>S.J. Khundmiri, M. Asghar, A.A. Banday, F. Khan, S. Salim, M. Levi and A.N.K. Yusufi</i> | 19 | The novel chelator lipid 3(nitrilotriacetic acid)-ditetradecylamine (NTA ₃ -DTDA) promotes stable binding of His-tagged proteins to liposomal membranes: Potent anti-tumor responses induced by simultaneously targeting antigen, cytokine and costimulatory signals to T cells
<i>C.L. van Broekhoven and J.G. Altin</i> | 104 |
| Investigation of a novel artificial antimicrobial peptide by fluorescence correlation spectroscopy: An amphipathic cationic pattern is sufficient for selective binding to bacterial type membranes and antimicrobial activity
<i>L. Yu, J.L. Ding, B. Ho and T. Wohland</i> | 29 | Porous membranes for reconstitution of ion channels
<i>M.A. Dhoke, P.J. Ladha, F.J. Boerio, L.B. Lessard, D.H. Malinowska, J. Cuppoletti and D.S. Wieczorek</i> | 117 |
| Composition dependence of vesicle morphology and mixing properties in a bacterial model membrane system
<i>B. Pozo Navas, K. Lohner, G. Deutsch, E. Sevcsik, K.A. Riske, R. Dimova, P. Garidel and G. Pabst</i> | 40 | Membrane perturbation effects of peptides derived from the N-termini of unprocessed prion proteins
<i>M. Magzoub, K. Ogłęcka, A. Pramanik, L.E. Göran Eriksson and A. Gräslund</i> | 126 |
| Coverage-dependent changes of cytochrome <i>c</i> transverse location in phospholipid membranes revealed by FRET
<i>Y.A. Domanov, J.G. Molotkovsky and G.P. Gorbenko</i> | 49 | Membrane-bound peptides mimicking transmembrane Vph1p helix 7 of yeast V-ATPase: A spectroscopic and polarity mismatch study
<i>R.W. Hesselink, R.B.M. Koehorst, P.V. Nazarov and M.A. Hemminga</i> | 137 |
| Using micellar mole fractions to assess membrane protein stability in mixed micelles
<i>P. Sehgal, J.E. Mogensen and D.E. Otzen</i> | 59 | <i>Cumulative Contents</i> | 146 |
| Large-scale purification of the proton pumping pyrophosphatase from <i>Thermotoga maritima</i> : A "Hot-Solve" method for isolation of recombinant thermophilic membrane proteins
<i>R.L. López-Marqués, J.R. Pérez-Castiñeira, M.J. Buch-Pedersen, S. Marco, J.-L. Rigaud, M.G. Palmgren and A. Serrano</i> | 69 | | |