



Expression of purinergic receptors and modulation of P2X₇ function by the inflammatory cytokine IFN γ in human epithelial cells

Lynn Welter-Stahl^{a,1}, Camila Marques da Silva^{b,1}, Julieta Schachter^b, Pedro Muanis Persechini^b, Heitor S. Souza^c, David M. Ojcius^{a,d}, Robson Coutinho-Silva^{b,*}

^a Université Paris – Denis Diderot, Paris, France

^b Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

^c Departamento de Clínica Médica, Hospital Universitário Clementino Fraga Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

^d School of Natural Sciences, University of California, Merced, CA, USA

ARTICLE INFO

Article history:

Received 11 November 2008

Received in revised form 10 March 2009

Accepted 13 March 2009

Available online 21 March 2009

Keywords:

P2X and P2Y receptors

Inflammation

Pro-inflammatory cytokine

Cervical epithelial cell

Apoptosis

ABSTRACT

The cervical epithelial cell line, HeLa, is one of the oldest and most commonly used cell lines in cell biology laboratories. Although a truncated P2X₇ receptor has recently been identified in HeLa cells, the expression of other purinergic receptors or the function of the P2X₇ protein has not been characterized. We here show that HeLa cells express transcripts for most P2X and P2Y purinergic receptors. Treatment of cells with ATP or other P2X₇ agonists does not stimulate cell death, but can induce atypical calcium fluxes and ion currents. Cervical epithelial cells represent an important target for sexually-transmitted pathogens and are commonly exposed to pro-inflammatory cytokines such as IFN γ . Stimulation of HeLa cells with IFN γ upregulates expression of P2X₇ mRNA and full-length protein, modifies ATP-dependent calcium fluxes, and renders the cells sensitive to ATP-induced apoptosis, which can be blocked by a P2X₇ antagonist. IFN γ treatment also increased dramatically the sensitivity of the intestinal epithelial cell line, HCT8, to ATP-induced apoptosis. Significantly, IFN γ also stimulated P2X₇ expression on human intestinal tissues. Responses to other purinergic receptor ligands suggest that HeLa cells may also express functional P2Y₁, P2Y₂ and P2Y₆ receptors, which could be relevant for modulating ion homeostasis in the cells.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

Extracellular nucleotides are signalling molecules that exert effects on a variety of tissues, mainly via interaction with specific membrane-bound receptors [1]. The target-cell responses are mediated by P2 receptors named either P2X or P2Y. P2X receptors are ligand-gated ion channels, and P2Y receptors are coupled to trimeric G-proteins [2]. Seven members of the P2X family in mammals have been cloned (P2X_{1–7}) [3], and eight P2Y receptors (P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁, P2Y₁₂, P2Y₁₃ and P2Y₁₄) [4,5].

Activation of the P2X₇ receptors leads to opening of a plasma membrane pore, inducing the permeabilization of the cells to molecules up to 800 Da; although not all cells display pore formation in response to P2X₇ ligation [3]. The P2X₇ receptor is responsible for

the release of pro-inflammatory cytokines such as IL-1 β and subsequently IL-6 in murine and human macrophages and mast cells [6,7], induction of apoptosis in different cell types [6,8], and elimination of intracellular parasites in macrophages [9–12].

Although most studies on purinergic receptors have been performed with cells from the central nervous system or macrophages, various P2X and P2Y receptors have also been described recently in epithelial cells from different sources [13,14]. Thus, human intestinal epithelial cells express mRNA for P2X₁, P2X₃, P2X₄, P2X₅, P2X₆, P2X₇, P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁ and P2Y₁₂, and the functional presence of P2X₇, P2Y₁, P2Y₂ and P2Y₄ has been confirmed based on the effect of extracellular nucleotides on apoptosis, cell proliferation, and calcium fluxes [15]. The P2X₇ receptor is also expressed in the murine gut epithelium, where its expression is associated with cell death [16]. While parotid acinar epithelial cells do not express functional P2X₇, duct cells express functional receptor [17], showing a restricted distribution of some receptors even within the same organ. The P2Y₁ and P2Y₂ receptors are also involved with ATP-induced calcium signaling in airway epithelial cells [18,19], and P2Y₂ has been linked with inhibition of growth of human esophageal squamous and colorectal carcinoma cells [20,21]. Finally, the P2Y₄ and P2Y₆ receptors control chloride transport in jejunum and colon epithelial cells, respectively [22,23].

Abbreviations: DMEM, Dulbecco's modified Eagle's Medium; IFN γ , interferon- γ ; IL-1 β , interleukin-1 β ; IL-6, interleukin-6; PLC, phospholipase C; TNF α , tumor necrosis factor α ; α , β -meATP, α , β -methylene adenosine 5'-triphosphate

* Corresponding author. Instituto de Biofísica Carlos Chagas Filho – UFRJ, Av. Carlos Chagas Filho n 373, Bloco G do CCS, Cidade Universitária, 21941-902; Ilha do Fundão, Rio de Janeiro, Brazil. Tel.: +55 21 2562 6565; fax: +55 21 2280 8193.

E-mail address: rcsilva@biof.ufrj.br (R. Coutinho-Silva).

¹ Both authors contributed equally to this work.

Nonetheless, the presence of the family of P2 receptors has not been fully characterized on the cervical epithelial cell line, HeLa, which represents the “work horse” of many cell biology and microbiology laboratories [24]. The uterine cervical epithelium is maintained by a balance between proliferation of a basal layer of cells, and death of cells in the upper layer. They regulate lubrication of the genital tract, represent a barrier against invasion by pathogens, and provide the necessary conditions for reproduction. Recently it has been shown that extracellular ATP mediates apoptosis via P2X₇ in hECE and CaSki, two human cervical epithelial cell lines; and that estrogen, epinephrine, and epidermal growth factor, combined with epinephrine, inhibit P2X₇-receptor-mediated pore formation and apoptosis in these cells [25–27]. HeLa cells, a human cervical carcinoma cell line that is used as a model host cell for infection by pathogens that infect the cervical epithelium but also other tissues [28–32], express P2Y₂ and P2Y₆ receptors [33,34] and a truncated form of P2X₇ [35]. However, given the presence of the truncated P2X₇ [35] and the effect of a PLC inhibitor on calcium signaling, it was assumed that HeLa cells do not express functional P2X purinergic receptors [33]. In addition, the functional expression of other P2 receptors on this cell line has never been investigated fully. We therefore examined whether other P2X and P2Y receptors may be expressed in epithelial HeLa cells, have begun to identify their cellular behavior, and have re-addressed the question of whether P2X₇ on HeLa cells may be functional. As cervical epithelial cells are also a common site of infection with sexually-transmitted pathogens, most of which initiate a strong inflammatory response [36–38], we also investigated whether the pro-inflammatory cytokines, IFN γ and TNF α , may modulate the expression or activity of P2X₇ receptors on HeLa cells.

2. Materials and methods

2.1. Cells and materials

The human cervical adenocarcinoma cell line, HeLa229, was from the American Type Culture Collection (ATCC; Manassas, VA). The cells were cultured in a humidified incubator at 37 °C with 5% CO₂ in Dulbecco's modified minimal essential medium (DMEM) with Glutamax-1 (Life Technologies, Inc.; Rockville, MD) supplemented with 10% heat-inactivated fetal calf serum (FCS) and 25 μ g/ml gentamicin. Recombinant human IFN- γ and TNF α were from BD Biosciences Pharmingen (San Jose, CA). All nucleotides were purchased from Sigma-Aldrich (St. Louis, MO). Fura-2 was obtained from Molecular Probes (Eugene, OR). Unless specified otherwise, all other reagents were purchased from Sigma. The nucleotides were prepared as 100 mM stock solution in PBS and stored at –20 °C until use.

2.2. RT-PCR detection of nucleotide receptors

RNA from HeLa cells was isolated using an RNeasy kit (Qiagen), following the manufacturer's instructions. Total RNA was converted into cDNA by standard reverse transcription with Superscript II RNase H[–] Reverse Transcriptase in the manufacturer's buffer (Invitrogen, Life Technologies; Carlsbad, CA). cDNAs were amplified using the iCycler (BIO-RAD laboratories, Richmond, CA) in a 50 μ l reaction mixture containing one-fifteenth of the cDNA generated from reverse transcription reaction, 1 \times PCR buffer, 2.5 mM MgCl₂, 0.25 mM (each) dNTPs, 0.5 μ M forward and reverse primers, and 1 U HotGoldstar DNA polymerase (Eurogentec, Seraing, Belgium). Amplification was done with the following sequences: GAPDH: 5'-AACGGATTGGTCCG-TATTGGGC-3', 5'-CTTGACGGTGCCATGGAATTG-3'; P2X₁: 5'-GCTGGTGCCTAATAAGAAGGTG-3', 5'-ATGAGCCGCTCGAGGTCTG-3'; P2X₂: 5'-AGGTTTGCCTAATACTACAAGATC-3', 5'-GCTGA-ACCTCCCGCTGTC-3'; P2X₃: 5'-CTTCACCTATGAGACCACCA-3', 5'-CGGTATTCTCTCACTCTCTG-3'; P2X₄: 5'-GATACCAGCTCAGGAG-GAAAC-3', 5'-GCATCATAAATGCACGACTTGAG-3'; P2X₅: 5'-GGCA-

TTCTGATGGCGCGTG-3', 5'-GGCACCAGGCAAAGATCTCAC-3'; P2X₆: 5'-AGCACTGCCGCTATGAACCAC-3', 5'-AGTGAGGCCAGCAGCCAGAG-3'; P2X₇: 5'-TGATAAAAAGTCTTCGGGATCCGT-3', 5'-TGGACA-AATCTGTGAAGTCCATC-3'; P2Y₁: 5'-CTTGGTGTGATTCTGGGCTG-3', 5'-GCTCGGGAGAGTCTCTTCTG-3'; P2Y₂: 5'-CCGCTCGTGGACCT-CAGCTG-3', 5'-CTCACTGCTGCCAACACATC-3'; P2Y₄: 5'-CGTGCCCACTCATCTAC-3', 5'-CCGAGTGGTGTATGGACAG-3'; P2Y₅: 5'-TGTTCACTCTACCACTCTCAG-3', 5'-CTTACTGCTGCCACTACT-GAGC-3'; P2Y₆: 5'-AACCTTGCTCTGGTGCACCTG-3', 5'-GCAGG-CACTGGGTGTGTCAGC-3'; P2Y₈: 5'-CTCTGGCGCACATCGTGAG-3', 5'-GGAGCGCAGCGACGTGGTC-3'; P2Y₁₀: 5'-GGTGTTCATGTGTGCTG-CAGTC-3', 5'-ATGGCGGGATAGTTGGTCAGC-3'; P2Y₁₁: 5'-GAGGCTG-CATCAAGTGTCTG-3', 5'-ACGTTGAGCACCCGATGATG-3'; P2Y₁₂: 5'-ACCGTTCATACGTAAGAACAG-3', 5'-GCAGAATTGGGGCACTTCAGC-3'; P2Y₁₃: 5'-GGAAGCAACACCATCTGTGTG-3', 5'-GACTGTGAGTA-TATGGAAGTCTG-3'; and P2Y₁₄: 5'-CCGCAACATATTCAGCATCTG-3', 5'-ACATTTGCAGCAGATAGTAGCAG-3'. The PCR cycling protocol for all primers was 94 °C at 45 s, 60 °C at 45 s, 72 °C at 45 s. The protocol was conducted for 40 cycles and included an initial 10-min enzyme activation step at 95 °C and a final 10-min extension step at 72 °C. PCR products were separated by electrophoresis on a 2% agarose gel and visualized by ethidium bromide staining. Peripheral blood lymphocytes were used as a positive control for expression of all the P2X receptors. The human laryngeal epithelial carcinoma cell line Hep2 was used to verify expression of all P2Y receptors.

2.3. Real-time PCR for P2X₇ mRNA from HeLa cells

RNA from HeLa cells was isolated using an RNeasy kit (Qiagen) following the manufacturer's instructions. Total RNA was converted into cDNA by standard reverse transcription with Superscript II RNase H[–] Reverse Transcriptase in the manufacturer's buffer (Invitrogen). Quantitative PCR was performed with one-fifteenth of the cDNA preparation in an Mx3000P (Stratagene; La Jolla, CA) in 25 μ l final volume with Brilliant QPCR Master Mix (Stratagene). cDNA was amplified using 300 nM of P2X₇ forward primer, 100 nM of P2X₇ reverse primer or 400 nM of GAPDH forward and reverse primers. We also used 300 nM of the fluorogenic oligonucleotides specific for the gene segments in which a reporter fluorescent dye on 5' (FAM[®]) and a quencher dye on 3' (TAMRA[®]) were attached. Real-time PCR was carried out at 95 °C for 10 min, followed by 40 cycles at 95 °C for 30 s, 60 °C for 1 min. The specific activities of cDNA from P2X₇ were compared with GAPDH and normalized to untreated control HeLa cells by the comparative C_T method as described by the manufacturer. For human GAPDH, the primers used were as follows: sense primer, 5'-GAGAAGGCTGGGGCTCAT-3'; antisense primer, 5'-TGCTGATGATCTT-GAGGCTG-3'; probe, 5'-CTCTGCTGATGCCCCATGTTCTG-3'. For P2X₇, the primers used were as follows: sense primer, 5'-CTTTCTCAAAA-CAGAAGGCCAAGA-3'; antisense primer, 5'-CAACCTCGGTCAGAGGAA-CAGA-3'; probe, 5'-TGTGTCCCGAGTATCCACCCGC-3'.

2.4. Measurement of cell death by cytofluorimetry

HeLa cells were incubated in the presence or absence of IFN γ (0.5, 1, or 2 ng/ml) for 3 days. HeLa cells were then treated with the indicated concentration of ATP (3 or 5 mM) in serum-free DMEM medium for 2 h, after which the medium was replaced with DMEM medium containing 10% heat-inactivated FCS, 25 μ g/ml gentamicin, in the absence of nucleotides. After an additional 24 h in the incubator at 37 °C, both adherent cells and cells in suspension were collected, washed twice in PBS, and analyzed by cytofluorimetry. Phosphatidylserine exposure on dying cells was measured by labeling with Annexin V-FITC (R&D Systems; Minneapolis, MN), and loss of plasma membrane integrity was measured by PI staining, as described [39]. Data from 10,000 HeLa cells were collected on a CyAn LX flow cytometer (Dako Cytomation) with a laser tuned to 488 nm.

In addition to analysis of phosphatidylserine exposure, in some experiments the degree of apoptosis was determined by counting the number of hypo-diploid nuclei after ATP treatment. In brief, HeLa and HCT8 cells were treated as described above. The cells were incubated with 300 μ M oATP for 2 h before ATP treatment. After the final 24 h in the incubator at 37 °C, both adherent cells and cells in suspension were collected, washed twice in PBS, spun for 10 min (200 \times g), and resuspended in an apoptotic buffer containing 50 μ g/ml ethidium bromide, 0.01 g sodium citrate, and 0.1% Triton X-100. The samples were transferred into 12 \times 75 mm FALCON 2052 FACS tubes (Becton Dickinson, San Jose, CA) and kept on ice until use. Data from 10,000 cells were collected on a FACScan flow cytometer (Becton Dickinson) equipped with a 488 nm argon laser. The cells acquired for analysis were gated to eliminate cell aggregates using WinMDI software. The cells with sub-diploid amounts of DNA were considered apoptotic.

2.5. Western blot analysis

HeLa cells were incubated in the presence or absence of IFN γ (0.2, 0.5, 2 or 5 ng/ml), or in a combination of IFN γ (0.5 or 2 ng/ml) plus TNF α (10 or 100 ng/ml) for 2 days. After an additional 24 h in the incubator at 37 °C in the absence of any cytokines, the cells were collected, transferred to microfuge tubes, and washed once in cold PBS, and the pellets were stored at –80 °C until use. Samples were lysed in buffer containing 50 mM Tris–HCl, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100 and protease inhibitor cocktail (5 μ l/ml, Sigma), and left for 30 min on ice. The protein concentration was determined using the micro BCA protein assay (Pierce), and 35 μ g of protein were separated on a 10% SDS-PAGE and transferred onto nitrocellulose membranes (Amersham Biosciences; Piscataway, NJ). After blocking in Tris-buffered saline containing 0.05% Tween 20 (TBST) and 5% BSA for 1 h, the membranes were washed extensively in TBST and immunostained with a rabbit polyclonal anti-P2X $_7$ receptor antibody (Calbiochem) diluted at 1:5000 in blocking buffer. Following three washes with TBST, membranes were incubated with an anti-rabbit polyclonal antibody conjugated to HRP diluted at 1:10,000 (Amersham Biosciences) in TBST supplemented with 5% nonfat dry milk. After subsequent washes, specific bands were visualized by enhanced chemiluminescence (Amersham Biosciences).

2.6. Statistics

The effect of ATP on the viability of HeLa cells was statistically evaluated using the Student *t* test. Effects of IFN γ on P2X $_7$ expression in the normal colonic epithelium were analyzed as the difference of expression using the Wilcoxon signed rank test.

2.7. Organ culture and immunohistochemistry

Colonic mucosal explants were cultured for 24 h at 37 °C with 95% O $_2$ and 5% CO $_2$ by standard procedures. Briefly, biopsy specimens were placed on iron grids in an organ culture dish in the presence of medium alone, consisting of RPMI 1640 medium (Life Technologies), 10% fetal calf serum, and penicillin and streptomycin (100 U/ml), or with the addition of IFN γ (2 ng/ml). After 24 h, mucosal explants were immediately embedded in Tissue-Tek O.C.T. compound (Miles Scientific Laboratories Ltd, Naperville, IL) and snap-frozen in isopentane in a liquid nitrogen bath. Samples were then stored at –80 °C until processing, and cut into 5 μ m sections in a cryostat maintained at –20 °C. Tissue sections were air-dried and fixed for 10 min in acetone. P2X $_7$ expression was evaluated using the indirect immunoperoxidase technique. Immunohistochemical staining was performed using the rabbit anti-P2X $_7$ antibody (1:300 dilution; Alomone, Jerusalem, Israel). Briefly, frozen sections were incubated at room temperature with normal horse serum for 30 min and then with the anti-P2X $_7$ antibody for 1 h. Negative control experiments were performed by pre-absorbing the

P2X $_7$ receptor antibodies with homologous peptide antigens and/or by omission of the primary P2X $_7$ antibodies. After rinsing in PBS for 10 min, tissue sections were incubated for 30 min with a goat anti-rabbit peroxidase conjugate IgG (Zymed Laboratories Inc.). Additional rinsing was followed by development with a solution containing hydrogen peroxide and diaminobenzidine, dehydrated, lightly counterstained in Harris's hematoxylin, dehydrated again, and mounted in histological mounting medium. Sections were examined by two independent observers using light microscopy at \times 400 magnification.

2.8. Intracellular calcium measurements

HeLa cells treated with 0, 1, or 2 ng/ml IFN γ were grown on glass coverslips for 48 h to subconfluence. The cells were then loaded with 2.5 μ M Fura-2-AM (Molecular Probes) for 40 min at room temperature in culture medium. The coverslips were washed in PBS and mounted in a three-compartment superfusion chamber attached to the stage of an inverted microscope (NIKON DIAPHOT 300 TMD) whose base was formed by a coverslip containing the cells. The central chamber containing the cells had a volume of 200 μ l, and was perfused with Ca $^{2+}$ -containing saline (PBS supplemented with 1 mM CaCl $_2$) or Ca $^{2+}$ -free saline (PBS supplemented with 1 mM EGTA) at room temperature at a rate of 1 ml min $^{-1}$. The intracellular calcium concentration of groups of 15–30 cells was monitored continuously with the use of a fluorescence photometer (Photon Technology; Princeton, NJ). Fura-2 was excited alternately at 340 and 380 nm, and the emission at 510 nm was measured. The ratio measurement, which is proportional to the intracellular calcium concentration, was determined every 100 ms. The nucleotides were injected in bolus, allowing the drugs to persist in the presence of the cells for 1 min, or the ATP was perfused continuously while the temperature was kept constant at 37 °C.

2.9. Electrophysiology

Cells were grown in 35 mm plastic culture dishes (Falcon) and incubated in the presence or absence of 1 or 2 ng/ml IFN γ for 24–48 h. The cells were then transferred to an inverted microscope (Zeiss Axiovert 100; Carl Zeiss; Jena, Germany). The culture medium was exchanged for an extracellular solution containing, in mM: 135 NaCl, 5 KCl, 1 MgCl $_2$, and 10 Na-HEPES, pH 7.4. Unless specified in the text, the experiments were performed at 23–25 °C. ATP and UTP were applied locally to the cell of interest using a microperfusion device. Recordings were carried out using the conventional whole cell patch-clamp technique [40]. Ionic currents were recorded in whole cell configuration, using an EPC-7 amplifier (List Electronic, Darmstadt, Germany) according to standard patch-clamping techniques [40]. Gigaohm seals were formed after offset potential compensation, using heat-polished micropipettes of 5–10 M Ω filled with an intrapipette solution (in mM: 135 KCl, 5 NaCl, 2 MgCl $_2$, 0.1 K-EGTA, and 10 K-HEPES, pH 7.4). ATP and UTP were applied to the cell surface by pneumatic injection, using a second micropipette filled with a different concentration of nucleotides dissolved in extracellular solution and connected to a PPM-2 pneumatic pump (NeuroPhore BH-2 system, List). Data were collected using pClamp and Fetchex software, version 6.0, and a Digidata 1200 interface (Axon Instruments) and plotted using Origin software version 4.0 (Microcal Inc.).

3. Results

3.1. Expression of P2X and P2Y mRNA on HeLa cells

In order to determine which P2X and P2Y isoforms are expressed by cervical epithelial cells, mRNA was isolated from HeLa cells and was analyzed by RT-PCR. Amplified PCR products of the expected sizes were obtained for P2X $_3$ (324 bp), P2X $_4$ (393 bp), P2X $_5$ (142 bp), P2X $_6$ (176 bp) and P2X $_7$ (396 bp) (Fig. 1A). The expression was faint for




Fig. 1. P2X and P2Y mRNA is expressed in human cervical epithelial cells. (A) P2X RT-PCR products in HeLa cells. (B) P2Y RT-PCR products in HeLa cells. Peripheral blood lymphocytes were used as positive controls for expression of the P2X receptors. All the PCR amplifications were performed at least 3 times.

P2X₃ mRNA, while P2X₁ and P2X₂ were completely absent. RT-PCR for the housekeeping gene, GAPDH, was performed to control for the quantity of cDNA loaded on the gel. Peripheral blood lymphocytes, used as a positive control for expression of all the P2X receptors, showed expression of P2X₁ and P2X₂ (data not shown), confirming the quality of the P2X₁ and P2X₂ PCR probes used for the epithelial cells. Amplification of P2X₄ and P2X₆ mRNA resulted in two fragments, one of the expected size and a second band. We had previously observed additional fragments for P2X₂ and P2X₃ in macrophages, which were ascribed to non-specific products of PCR amplification and alternative splicing [46]. Similarly, amplified PCR products of the expected sizes were obtained in HeLa cells for P2Y₁ (327 bp), P2Y₂ (259 bp), P2Y₄ (311 bp), P2Y₆ (283 bp), P2Y₁₁ (260 bp), P2Y₁₂ (307 bp), P2Y₁₃ (293 bp) and P2Y₁₄ (168 bp), and also the orphans genes P2Y₅ (351 bp), P2Y₈ (237 bp) and P2Y₁₀ (227 bp) (Fig. 1B).

3.2. IFN γ modulates P2X₇ receptor expression in HeLa cells

The synthesis of P2X₇ receptor protein was then assessed with and without pre-treatment with IFN γ , and IFN γ plus TNF α , using the Western blot assay. As shown in Fig. 2 the 70 kDa band characteristic of the full-length P2X₇ receptor [41] was present on control untreated HeLa cells (Fig. 2A lane 1), besides an additional 40 kDa band, likely corresponding to truncated P2X₇ protein. We observed a dose-dependent increase in levels of the full-length P2X₇ receptor band following IFN γ treatment, with the concomitant disappearance of the 40 kDa band (Fig. 2A lanes 2–5). It has been shown that the pro-inflammatory cytokine TNF α has a synergistic effect on P2X₇ receptor modulation in macrophages [42,43]. The HeLa cells were thus co-treated with different concentrations of IFN γ plus TNF α , as described in the Materials and methods. However, the presence of TNF α did not have an additional effect over IFN γ alone (data not shown).

3.3. Effect of IFN γ treatment on transcription of the P2X₇ gene

Since IFN γ treatment had a positive effect on P2X₇ receptor protein expression, we evaluated whether IFN γ could affect P2X₇ at the

transcriptional level. We thus incubated the cells with 0, 0.5, 1 or 2 ng/ml of IFN γ for 72 h and measured P2X₇ transcription by real-time PCR. There was a large dose-dependent increase in the amount of P2X₇ mRNA after IFN γ treatment (Fig. 2B). A noticeable effect was observed with 0.5 ng/ml IFN γ , but became much larger after treatment with 2 ng/ml IFN γ .

3.4. HeLa cells become sensitive to ATP-induced lysis after IFN γ treatment

HeLa cells are known to be more resistant to ATP-induced lysis than primary cervical cells [26]. But after treatment with IFN γ , the HeLa cells became sensitive to ATP as a function of the IFN γ dose (Fig. 3A). The mode of cell death was next analyzed by double-staining with Annexin V and PI. We observed that untreated cells are completely resistant to apoptosis induced by ATP, but the IFN γ -treated cells became sensitive to ATP-mediated cell death, with characteristics of both apoptosis and necrosis (Fig. 3B). We next investigated the effect of the irreversible P2X₇ antagonist, oATP, on apoptosis in the presence of ATP. We pre-treated HeLa cells that were incubated with IFN γ with oATP before incubation with ATP. Pre-treatment with oATP inhibited ATP-induced apoptosis, further confirming the IFN γ -mediated modulation of P2X₇ receptor expression (Fig. 3C).

A similar effect was observed with another human epithelial cell line, the intestinal epithelial HCT8 cells, after treatment with IFN γ . We observed that HCT8 cells were 50% more sensitive to ATP-mediated apoptosis than HeLa cells in the absence of IFN γ treatment, but became very sensitive to ATP-induced apoptosis after pre-treatment with increasing concentrations of IFN γ (Fig. 3D).

To determine whether IFN γ could also modulate P2X₇ receptor expression in primary human epithelial cells, we incubated explants of human intestinal epithelial tissue with IFN γ for 24 h and then




Fig. 2. The pro-inflammatory cytokine IFN γ positively modulates protein and mRNA expression of the P2X₇ receptor. (A) Western blot analysis of total protein extracts from untreated HeLa cells and HeLa cells treated with IFN γ (0.2, 0.5, 2 and 5 ng/ml) for 2 days. Lane 1: untreated HeLa cells. Lanes 2–5: cells treated with IFN γ at 0.2, 0.5, 2 and 5 ng/ml, respectively. Lane 6: the J774 macrophage cell line used as a positive control. Upper lower arrows indicate the 70 kDa and 40 kDa bands. The results shown are representative of two experiments performed on different days. (B) Cells were treated with control buffer or increasing concentrations of IFN γ for 3 days. Total mRNA was measured by real-time PCR, as described in the Materials and methods. The experiment was performed on two separate days, with similar results.




Fig. 3. IFN γ increases the sensitivity to ATP-mediated cell death. HeLa cells were incubated in the presence or absence of IFN γ (0.5, 1, or 2 ng/ml) for 3 days. The cells were then treated with increasing concentrations of ATP for 2 h, and maintained in cell culture for an additional 24 h. (A) The loss of viability induced by ATP after IFN γ incubation, measured by cytofluorimetry (viable cells = cells negative for both Annexin V and PI). B shows the increase in the number of apoptotic cells after incubation with IFN γ (2 ng/ml). Phosphatidylserine exposure on dying cells was measured by labeling with Annexin V-FITC, and loss of plasma membrane integrity was measured by PI staining. C and D show the formation of hypo-diploid nuclei after ATP treatment in HeLa cells (C) and HCT8 cells (D). Note that the pre-treatment with 300 μ M ATP inhibited ATP-induced apoptosis. The data shown in A and B are from a representative experiment performed in triplicate repeated on two separate occasions. The data shown in C and D are mean \pm SEM of three and four independent experiments performed in triplicate (* P < 0.05 for 3 mM ATP compared with 0 mM ATP, cells treated with 1 ng/ml IFN γ ; ** P < 0.001 for 3 and 5 mM ATP compared with 0 mM ATP, cells treated with 2 ng/ml IFN γ).

examined the samples by immunohistochemistry for the presence of the P2X $_7$ receptor. We observed an increase in P2X $_7$ -specific immunostaining on epithelial cells from explants incubated with IFN γ (Fig. 4A–B) compared with untreated ones (Fig. 4C–D), demonstrating that the effects of IFN γ on P2X $_7$ expression are not limited to epithelial cell lines. Quantification of the effect of IFN γ on P2X $_7$ expression is shown in Fig. 4E.

3.5. ATP and UTP elicit calcium fluxes in HeLa cells

To identify the P2-receptor subtypes that may be functional in HeLa cells, we measured changes in $[Ca^{2+}]_i$ induced by nucleotides. In Ca^{2+} -containing PBS solution, successive application of ATP and UTP (100 μ M) induced a rapid transient increase of $[Ca^{2+}]_i$, regardless of which nucleotide was added first (Fig. 5A, B).




Fig. 4. Effect of IFN γ on P2X $_7$ expression in the normal colonic epithelium. Human colonic samples from endoscopic biopsies were treated in organ cultures with or without the addition of 2 ng/ml IFN γ for 24 h. Frozen sections were stained by immunoperoxidase using an antibody against the P2X $_7$ receptor. A–D shows immunostaining for P2X $_7$ receptors on human epithelial tissues from untreated (C and D) and IFN γ -treated (A and B) samples. Sections were examined using light microscopy at $\times 100$ (A and C) and $\times 400$ magnification (B and D). E, The density of P2X $_7$ -positive cells is given as the percentage of immunoreactive cells among epithelial cells in the crypts and in the surface epithelium. Numbers are representative of five different experiments. Horizontal bars represent medians, boxes represent the 25th and 75th percentiles, and vertical bars represent ranges. Differences were analyzed using the Wilcoxon signed rank test.

Removing Ca^{2+} from the extracellular bath barely reduced the ATP responses, suggesting that the main cytosolic $[\text{Ca}^{2+}]_i$ increase was due to release from intracellular stores (Fig. 5C). In addition, the ATP response was only partially blocked by pre-treatment with the P2X $_7$ antagonist, oATP (Fig. 5D), and the UTP response was completely blocked by suramin treatment (data not shown). To investigate possible cross-desensitization between ATP and UTP in nucleotide-induced $[\text{Ca}^{2+}]_i$ changes, HeLa cells were perfused with 100 μM ATP or UTP, until the $[\text{Ca}^{2+}]_i$ response returned to basal levels, and the cells were then exposed to 100 μM of a second agonist (ATP or UTP) in the continued presence of the first agonist. As shown in Fig. 5E, UTP pre-treatment markedly, but not completely, reduced the $[\text{Ca}^{2+}]_i$ response to ATP, whereas ATP

pre-treatment completely abolished the UTP-induced $[\text{Ca}^{2+}]_i$ changes (Fig. 5F). UTP and ATP promoted dose-dependent $[\text{Ca}^{2+}]_i$ fluxes transiently, with responses starting at 100 nM for both nucleotides, and an EC $_{50}$ of 17 ± 4 and 27 ± 5 μM for UTP and ATP, respectively (Fig. 5G). The nucleotides UDP, ADP and $\alpha\beta$ -meATP also elicited changes in $[\text{Ca}^{2+}]_i$, beginning at 5 μM for ADP and UDP and 25–100 μM for $\alpha\beta$ -meATP (Fig. 5H).

3.6. IFN- γ treatment modifies the pattern of ATP and UTP-induced $[\text{Ca}^{2+}]_i$ responses

HeLa cells were perfused with buffer saline containing 1 mM ATP at 37 °C. Under these conditions, we observed a transient rise




Fig. 5. Effect of different nucleotides on intracellular Ca^{2+} fluxes. HeLa cells were loaded with Fura-2, and nucleotides were added in a bolus of 1-min duration in a continuous perfusion chamber. (A) Effects of 100 μM ATP on $[\text{Ca}^{2+}]_i$, followed by stimulation with 100 μM UTP. (B) Same as in A, but UTP was applied first. (C) 100 μM ATP was added in Ca^{2+} -containing saline (first bar) and in calcium-free saline (second bar). (D) Effect of 100 μM ATP on $[\text{Ca}^{2+}]_i$ in cells pre-treated with oATP. (E and F) The effect of ATP/UTP on cross-desensitization of Ca^{2+} responses. (E) Addition of 100 μM ATP during prolonged stimulation with 100 μM UTP, showing a small remaining response to ATP. (F) Addition of 100 μM UTP during prolonged stimulation with 100 μM ATP, showing that pre-stimulation with ATP blocks UTP-induced calcium responses. (G) Dose response curves for ATP- and UTP-induced $[\text{Ca}^{2+}]_i$ responses. Cells were exposed to the indicated concentration of ATP (open circles) or UTP (filled triangles). (H) Changes in $[\text{Ca}^{2+}]_i$ in response to 1, 5, 10, 25, 50, 75 and 100 μM of the indicated nucleotides. The $[\text{Ca}^{2+}]_i$ is given as the fluorescence ratio (F340/F380 nm), where fluorescence is proportional to the intracellular calcium concentration. In A–F, representative data of a single experiment with 20 cells per incubation are shown. The experiments were repeated at least three times with similar results. Values in G and H represent mean \pm SEM of five (G) and six (H) independent experiments.

in $[\text{Ca}^{2+}]_i$, which returned to basal levels after ~ 2 min (125 ± 16 s, $n = 4$) (Fig. 6A, C). The response was longer but comparable to the response observed with a small bolus application at room temperature. However, cells pre-incubated with 1–2 ng/ml IFN γ for 24 h responded to ATP perfusion with a rapid biphasic

increase of $[\text{Ca}^{2+}]_i$, followed by a sustained plateau that slowly (4–5 min) (267 ± 28 s, $n = 5$) returned to $[\text{Ca}^{2+}]_i$ basal levels (Fig. 6B, C), compatible with the opening of a non-selective pore on the plasma membrane, as demonstrated for P2X $_7$ in other cell types [6,8,41,44]. The pre-treatment with the P2X $_7$ receptor




Fig. 6. Effect of IFN γ on ATP- and UTP-induced Ca^{2+} responses. (A) Untreated cells under perfusion with 1 mM ATP at 37 °C, showing a fast peak and gradual return to basal levels. (B) Cells pre-treated with 2 ng/ml IFN γ for 24 h, under perfusion with 1 mM ATP, showing a fast peak that remained at high levels for several minutes. (C) Graph showing the time needed for $[Ca^{2+}]_i$ changes shown in A and B to return to basal levels. (D) same in B, with cells pre-treated for 2 h with oATP. The $[Ca^{2+}]_i$ is given as the fluorescence ratio (F340/F380 nm), where fluorescence is proportional to the intracellular calcium concentration. (E and F) Dose response curves for UTP-induced $[Ca^{2+}]_i$ responses in untreated and IFN γ -treated cells, respectively. (G) UTP was added in a bolus of 1-min duration, followed by a second 1-min bolus 5 min later, in calcium-free saline and a continuous perfusion chamber at room temperature. Values in C represent mean \pm SEM of four (A) and five (B) independent experiments. $n = 3$ and $n = 5$ for curves shown in E and F, respectively. Values in G represent mean \pm SEM of 3 independent experiments.




Fig. 7. Profile of ATP-activated currents in epithelial cells that were not treated with IFN γ . Whole cell currents were measured in HeLa cells kept at $V_H = -40$ mV. (A) Transient outward current after application of 1 mM ATP. (B and C) Transient outward currents after application of 100 μ M UTP, showing a delay between stimulation with the nucleotide and initiation of the outward current. None of the cells tested responded with inward currents ($n = 11$). (D and E) Outward current desensitization after successive stimulation with 100 μ M UTP. The first and second arrows indicate the fourth and fifth application of UTP.

antagonist, oATP, strongly inhibited the calcium response observed in IFN γ -incubated cells (Fig. 6D), further suggesting a role for P2X $_7$ in the ATP-dependent effects.

In order to investigate the possibility that IFN γ could be affecting the activity of receptors other than P2X $_7$, we also characterized the effect of IFN γ on UTP-induced fluxes. In untreated cells, UTP promoted dose-dependent $[Ca^{2+}]_i$ fluxes transiently, with an EC $_{50}$ of 14 μ M (Fig. 6E). The EC $_{50}$ decreased

slightly (≈ 9 μ M) in IFN γ -incubated cells, but the UTP-induced response clearly did not fit a typical sigmoid curve (Fig. 6F), suggesting the presence of more than one functional P2Y receptor on IFN γ -incubated cells. Comparing the amplitude of the UTP-induced calcium responses in untreated and IFN γ -treated cells, we observed that two successive applications of 100 μ M UTP induced a rapid transient increase of $[Ca^{2+}]_i$ in untreated cells, with similar amplitudes both times (Fig. 6G). However, in IFN γ -incubated cells,




Fig. 8. IFN γ pre-treatment modifies ATP-mediated currents. Whole cell currents were measured in cells treated with 2 ng/ml IFN γ for 24 h and kept at a $V_H = -40$ mV. (A) A biphasic current induced immediately after ATP application. (B) Same cell as in A, showing that while the outward component of the response disappeared, the inward current does not desensitize. Note that the inward current starts immediately after ATP stimulation. Two of six cells responded as shown in A and B, while four other cells responded with outward currents, as shown for cells not treated with IFN γ .

the first response was bigger than in untreated cells, and the second response was smaller than the first one (Fig. 6G).

3.7. ATP induces ion currents in HeLa cells

Application of 1 mM ATP induced an outward current in HeLa cells after 10–15 s (Fig. 7A). The same transient outward currents were observed in response to 100 μ M UTP (Fig. 7B). Occasionally the outward currents were biphasic or took place in multiples steps, and displayed clear desensitization (Fig. 7C). We observed almost complete outward current desensitization after four or five successive applications of UTP (Fig. 7D). None of the cells tested responded with an inward current. The behavior of the responses was compatible with the activation of a Ca^{2+} -activated K^+ current, similarly to the $P2Y_2$ -dependent response previously observed in other cell types [45,46].

A very different response was observed in HeLa cells pre-treated for 24 h with IFN γ . ATP application elicited biphasic responses composed of a fast inward current followed by an outward current (Fig. 8A). In some cells, we observed only the inward or the outward component of the current (data not shown). After continued ATP stimulation, the outward current disappeared, and only the inward component remained (Fig. 8B).

4. Discussion

Purinergic signalling is an important component of host defence against intracellular pathogens, mainly through activation of P2 receptors and particularly the $P2X_7$ receptor [47]. The $P2X_7$ receptor is involved in elimination of mycobacteria and chlamydiae in infected macrophages [9,11], and secretion of pro-inflammatory cytokines such as IL-1 β and IL-18 [48]. Extracellular ATP is released from infected or

stressed cells, and is now viewed as a “danger signal” that, acting through the $P2X_7$ receptor, can stimulate secretion of pro-inflammatory cytokines [49,50]. Ligation of the $P2X_7$ receptor has also been associated with homeostasis of T regulatory cells and regulation of autoimmune disease [51–54].

Our experiments suggest that, in the absence of IFN γ stimulation, HeLa cells express functional $P2X_7$ at a low level. Although $P2X_7$ mRNA and $P2X_7$ protein was detected in HeLa cells, the functional experiments demonstrate that ATP treatment does not elicit pore opening nor induce apoptosis, the two hallmarks of functional $P2X_7$ expression [1,55]. It has been proposed that a low level of expression of this receptor makes it unresponsive to ATP [56]. Similarly, it has recently been suggested that hetero-oligomerization of $P2X_7$ with a naturally occurring truncated form of the receptor could block the receptor's function in HeLa cells, as described by Feng et al. [35]. In support of this hypothesis, we observed the presence of a 40 kDa band on gels from HeLa cells that was not present in macrophage extracts, which could correspond to the truncated form of $P2X_7$. Another possibility is that our antibody cross-reacts with a different splice variant, since under our experimental conditions [56] we might not expect to observe the 40 kDa band described by Feng et al. Regardless, the new band disappears after IFN γ treatment, which is associated with a change in the functional state of the $P2X_7$ receptor.

Furthermore, IFN γ treatment clearly induces up-regulation of $P2X_7$ receptor and activity, as revealed by an increase in the levels of $P2X_7$ mRNA and protein, $P2X_7$ -dependent calcium fluxes, and the presence of non-desensitizing inward currents. Interestingly, we could not detect the 40 kDa band in extracts of IFN γ -treated cells.

Most if not all cells express more than one subtype of P2 receptors [57]. But the observation that UTP-induced calcium signaling in HeLa cells did not completely eliminate subsequent ATP responses suggests that the cells express ATP receptors that are not sensitive to UTP. We could not detect calcium signaling using $\alpha\beta$ -meATP at concentrations that activate $P2X_{1-3}$ receptors, implying that those P2X receptors are not functional in HeLa cells. On the other hand, the outward currents observed after ATP and UTP addition confirm the functional presence of $P2Y$, possibly $P2Y_2$ receptors, on the cells [58]. Our data on calcium fluxes suggest the presence of UDP-sensitive and ADP-sensitive $P2Y$ receptors, possibly $P2Y_6$ and $P2Y_1$ subtypes, but this conclusion will require further studies. It has been proposed previously that $P2Y_2$ receptors on HeLa cells inhibit the activity of the Na^+/K^+ -ATPase [33], but unlike the present study, the authors were able to detect mRNA for only the $P2Y_4$ and $P2Y_6$ subtypes on HeLa cells. Since $P2Y_4$ in humans is activated only by UTP [59], it should not be functional in HeLa cells, despite being present at the transcriptional level, as there were no UTP-induced calcium transients after perfusion with ATP. Our measurement of calcium fluxes in the presence of the $P2Y_2$ antagonist, suramin, confirms the functional expression of $P2Y_2$ receptors on HeLa cells. The UTP dose–response curve and sequential UTP pulses after IFN γ treatment suggest that this pro-inflammatory cytokine upregulates additional $P2Y$ receptors on epithelial cells.

In conclusion, our study demonstrates that the most popular cell type in cell biology laboratories expresses a functional $P2X_7$ receptor, which can be upregulated under the inflammatory conditions found during microbial infection, thus amplifying inflammatory responses requiring stimulation of a cytosolic “inflammasome” by extracellular danger signals [60,61]. Even though mRNA for a large number of purinergic receptors is present, HeLa cells appear to express a smaller subset of functional receptors, most likely for $P2Y_1$, $P2Y_2$ and $P2Y_6$. These receptors may also play a role in regulating essential functions of the epithelial cells, such as maintenance of the Na^+ and K^+ gradients across the plasma membrane and controlling infection by intracellular pathogens.

Acknowledgements

We thank Vandir da Costa, Thomas Jungas, and Anneliese F. A. F. da Costa for technical and expert assistance with cell culture. This work was supported by funds from the Conselho Nacional de Desenvolvimento Científico e Tecnológico do Brasil (CNPq), Programa de Núcleos de Excelência (Pronex), Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), and National Institutes of Health.

References

- [1] B.S. Khakh, R.A. North, P2X receptors as cell-surface ATP sensors in health and disease, *Nature* 442 (2006) 527–532.
- [2] V. Ralevic, G. Burnstock, Receptors for purines and pyrimidines, *Pharmacol. Rev.* 50 (1998) 413–492.
- [3] R.A. North, Molecular physiology of P2X receptors, *Physiol. Rev.* 82 (2002) 1013–1067.
- [4] L. Erb, Z. Liao, C.I. Seye, G.A. Weisman, P2 receptors: intracellular signaling, *Pflügers Arch.* 452 (2006) 552–562.
- [5] M.P. Abbracchio, J.M. Boeynaems, E.A. Barnard, J.L. Boyer, C. Kennedy, M.T. Miras-Portugal, B.F. King, C. Gachet, K.A. Jacobson, G.A. Weisman, G. Burnstock, Characterization of the UDP-glucose receptor (re-named here the P2Y14 receptor) adds diversity to the P2Y receptor family, *Trends Pharmacol. Sci.* 24 (2003) 52–55.
- [6] F. Di Virgilio, P. Chiozzi, D. Ferrari, S. Falzoni, J.M. Sanz, A. Morelli, M. Torboli, G. Bolognesi, O.R. Baricordi, Nucleotide receptors: an emerging family of regulatory molecules in blood cells, *Blood* 97 (2001) 587–600.
- [7] E. Bulanova, V. Budagian, Z. Orinska, M. Hein, F. Petersen, L. Thon, D. Adam, S. Bulfone-Paus, Extracellular ATP induces cytokine expression and apoptosis through P2X7 receptor in murine mast cells, *J. Immunol.* 174 (2005) 3880–3890.
- [8] R. Coutinho-Silva, P.M. Persechini, R.C. Bisaggio, J.-L. Perfettini, A.C.T. Sá-Neto, J. M. Kanellopoulos, I. Motta-Ly, A. Dautry-Varsat, D.M. Ojcius, P2Z/P2X7 receptor-dependent apoptosis of dendritic cells, *Am. J. Physiol.* 276 (1999) C1139–C1147.
- [9] R. Coutinho-Silva, L. Stahl, M.-N. Raymond, T. Jungas, P. Verbeke, G. Burnstock, T. Darville, D.M. Ojcius, Inhibition of chlamydial infectious activity due to P2X7R-dependent phospholipase D activation, *Immunity* 19 (2003) 403–412.
- [10] R. Coutinho-Silva, J.-L. Perfettini, P.M. Persechini, A. Dautry-Varsat, D.M. Ojcius, Modulation of P2Z/P2X7 receptor activity in macrophages infected with *Chlamydia psittaci*, *Am. J. Physiol.* 280 (2001) C81–C89.
- [11] D.A. Lammas, C. Stober, C.J. Harvey, N. Kendrick, S. Panchalingam, D.S. Kumararatne, ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2X7) receptors, *Immunity* 7 (1997) 433–444.
- [12] I.P. Fairbairn, C.B. Stober, D.S. Kumararatne, D.A. Lammas, ATP-mediated killing of intracellular mycobacteria by macrophages is a P2X(7)-dependent process inducing bacterial death by phagosome-lysosome fusion, *J. Immunol.* 167 (2001) 3300–3307.
- [13] J. Leipziger, Control of epithelial transport via luminal P2 receptors, *Am. J. Physiol., Renal Physiol.* 284 (2003) F419–F432.
- [14] E.M. Schwiebert, A. Zsembery, Extracellular ATP as a signaling molecule for epithelial cells, *Biochim. Biophys. Acta* 1615 (2003) 7–32.
- [15] R. Coutinho-Silva, L. Stahl, K.K. Cheung, N.E. de Campos, S.C. de Oliveira, D.M. Ojcius, G. Burnstock, P2X and P2Y purinergic receptors on human intestinal epithelial carcinoma cells: effects of extracellular nucleotides on apoptosis and cell proliferation, *Am. J. Physiol., Gastrointest. Liver Physiol.* 288 (2005) G1024–G1035.
- [16] U. Groschel-Stewart, M. Bardini, T. Robson, G. Burnstock, P2X receptors in the rat duodenal villus, *Cell Tissue Res.* 297 (1999) 111–117.
- [17] Q. Li, X. Luo, W. Zeng, S. Muallem, Cell-specific behavior of P2X7 receptors in mouse parotid acinar and duct cells, *J. Biol. Chem.* 278 (2003) 47554–47561.
- [18] L. Homolya, W.C. Watt, E.R. Lazarowski, B.H. Koller, R.C. Boucher, Nucleotide-regulated calcium signaling in lung fibroblasts and epithelial cells from normal and P2Y(2) receptor (—/—) mice, *J. Biol. Chem.* 274 (1999) 26454–26460.
- [19] A. Marino, Y. Rodrig, M. Metioui, L. Lagneaux, E. Alzola, M. Fernandez, D.J. Fogarty, C. Matute, A. Moran, J.P. Dehaye, Regulation by P2 agonists of the intracellular calcium concentration in epithelial cells freshly isolated from rat trachea, *Biochim. Biophys. Acta* 1439 (1999) 395–405.
- [20] M. Hopfner, K. Maaser, B. Barthel, B. von Lampe, C. Hanski, E.O. Riecken, M. Zeit, H. Scherubl, Growth inhibition and apoptosis induced by P2Y2 receptors in human colorectal carcinoma cells: involvement of intracellular calcium and cyclic adenosine monophosphate, *Int. J. Colorectal Dis.* 16 (2001) 154–166.
- [21] K. Maaser, M. Hopfner, H. Kap, A.P. Sutter, B. Barthel, B. von Lampe, M. Zeit, H. Scherubl, Extracellular nucleotides inhibit growth of human oesophageal cancer cells via P2Y(2)-receptors, *Br. J. Cancer* 86 (2002) 636–644.
- [22] M. Kottgen, T. Löffler, C. Jacobi, R. Nitschke, H. Pavenstadt, R. Schreiber, S. Frische, S. Nielsen, J. Leipziger, P2Y6 receptor mediates colonic NaCl secretion via differential activation of cAMP-mediated transport, *J. Clin. Invest.* 111 (2003) 371–379.
- [23] B. Robaye, E. Ghanem, F. Wilkin, D. Fokan, W. Van Driessche, S. Schurmans, J.M. Boeynaems, R. Beauwens, Loss of nucleotide regulation of epithelial chloride transport in the jejunum of P2Y(4)-null mice, *Mol. Pharmacol.* 63 (2003) 777–783.
- [24] S. Grimm, The art and design of genetic screens: mammalian culture cells, *Nat. Rev., Genet.* 5 (2004) 179–189.
- [25] Q. Wang, L. Wang, Y.H. Feng, X. Li, R. Zeng, G.I. Gorodeski, P2X7 receptor-mediated apoptosis of human cervical epithelial cells, *Am. J. Physiol., Cell Physiol.* 287 (2004) C1349–C1358.
- [26] Q. Wang, X. Li, L. Wang, Y.H. Feng, R. Zeng, G. Gorodeski, Antiapoptotic effects of estrogen in normal and cancer human cervical epithelial cells, *Endocrinology* 145 (2004) 5568–5579.
- [27] L. Wang, Y.H. Feng, G.I. Gorodeski, Epidermal growth factor facilitates epinephrine inhibition of P2X7-receptor-mediated pore formation and apoptosis: a novel signaling network, *Endocrinology* 146 (2005) 164–174.
- [28] S.F. Fischer, J. Vier, S. Kirschnek, A. Klos, S. Hess, S. Ying, G. Hacker, Chlamydia inhibit host cell apoptosis by degradation of proapoptotic BH3-only proteins, *J. Exp. Med.* 200 (2004) 905–916.
- [29] P. Tang, C.L. Sutherland, M.R. Gold, B.B. Finlay, *Listeria monocytogenes* invasion of epithelial cells requires the MEK-1/ERK-2 mitogen-activated protein kinase pathway, *Infect. Immun.* 66 (1998) 1106–1112.
- [30] B.K. Coombes, M.E. Wickham, M.J. Lowden, N.F. Brown, B.B. Finlay, Negative regulation of *Salmonella* pathogenicity island 2 is required for contextual control of virulence during typhoid, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 17460–17465.
- [31] J.J. Martinez, S. Seveau, E. Veiga, S. Matsuyama, P. Cossart, Ku70, a component of DNA-dependent protein kinase, is a mammalian receptor for *Rickettsia conorii*, *Cell* 123 (2005) 1013–1023.
- [32] E. Veiga, P. Cossart, *Listeria* hijacks the clathrin-dependent endocytic machinery to invade mammalian cells, *Nat. Cell Biol.* 7 (2005) 894–900.
- [33] A. Muscella, M.G. Elia, S. Greco, C. Storelli, S. Marsigliante, Activation of P2Y2 purinoceptor inhibits the activity of the Na⁺/K⁺-ATPase in HeLa cells, *Cell. Signal.* 15 (2003) 115–121.
- [34] A. Muscella, S. Greco, M.G. Elia, C. Storelli, S. Marsigliante, Differential signalling of purinoceptors in HeLa cells through the extracellular signal-regulated kinase and protein kinase C pathways, *J. Cell. Physiol.* 200 (2004) 428–439.
- [35] Y.H. Feng, X. Li, L. Wang, L. Zhou, G.I. Gorodeski, A truncated P2X7 receptor variant (P2X7-j) endogenously expressed in cervical cancer cells antagonizes the full-length P2X7 receptor through hetero-oligomerization, *J. Biol. Chem.* 281 (2006) 17228–17237.
- [36] S.P. Lad, E.Y. Fukuda, J. Li, L.M. de la Maza, E. Li, Up-regulation of the JAK/STAT1 signal pathway during *Chlamydia trachomatis* infection, *J. Immunol.* 174 (2005) 7186–7193.
- [37] R.N. Fichorova, P.J. Desai, F.C. Gibson III, C.A. Genco, Distinct proinflammatory host responses to *Neisseria gonorrhoeae* infection in immortalized human cervical and vaginal epithelial cells, *Infect. Immun.* 69 (2001) 5840–5848.
- [38] S.J. Rasmussen, L. Eckmann, A.J. Quayle, L. Shen, Y.X. Zhang, D.J. Anderson, J. Fierer, R.S. Stephens, M.F. Kagnoff, Secretion of proinflammatory cytokines by epithelial cells in response to Chlamydia infection suggests a central role for epithelial cells in chlamydial pathogenesis, *J. Clin. Invest.* 99 (1997) 77–87.
- [39] J.L. Perfettini, M. Gissot, P. Souque, D.M. Ojcius, Modulation of apoptosis during infection with Chlamydia, *Methods Enzymol.* 358 (2002) 334–344.
- [40] O.P. Hamill, A. Marty, E. Neher, B. Sakmann, F.J. Sigworth, Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches, *Eur. J. Physiol.* 391 (1981) 85–100.
- [41] A. Morelli, D. Ferrari, G. Bolognesi, R. Rizzuto, F. Di Virgilio, Proapoptotic plasma membrane pore: P2X7 receptor, *Drug Dev. Res.* 52 (2001) 571–578.
- [42] B.D. Humphreys, G.R. Dubyak, Modulation of P2X7 nucleotide receptor expression by pro- and anti-inflammatory stimuli in THP-1 monocytes, *J. Leukoc. Biol.* 64 (1998) 265–273.
- [43] B.D. Humphreys, G.R. Dubyak, Induction of the P2Z/P2X₇ nucleotide receptor and associated phospholipase D activity by lipopolysaccharide and IFN-γ in the human THP-1 monocytic cell line, *J. Immunol.* 157 (1996) 5627–5637.
- [44] P.M. Persechini, R.C. Bisaggio, J.L. Alves-Neto, R. Coutinho-Silva, Extracellular ATP in the lymphohematopoietic system: P2Z purinoceptors and membrane permeabilization, *Braz. J. Med. Biol. Res.* 31 (1998) 25–34.
- [45] C. Albuquerque, S.M. Oliveira, R. Coutinho-Silva, G.M. Oliveira-Castro, P.M. Persechini, ATP- and UTP-induced currents in macrophages and macrophage-polykaryons, *Am. J. Physiol.* 265 (1993) C1663–C1673.
- [46] R. Coutinho-Silva, D.M. Ojcius, D.C. Gorecki, P.M. Persechini, R.C. Bisaggio, A.N. Mendes, J. Marks, G. Burnstock, P.M. Dunn, Multiple P2X and P2Y receptor subtypes in mouse J774, spleen and peritoneal macrophages, *Biochem. Pharmacol.* 69 (2005) 641–655.
- [47] R. Coutinho-Silva, C.M. da Cruz, P.M. Persechini, D.M. Ojcius, The role of P2 receptors in controlling infections by intracellular pathogens, *Purinergic Signaling* 3 (2007) 83–90.
- [48] D. Ferrari, C. Pizzirani, E. Adinolfi, R.M. Lemoli, A. Curti, M. Idzko, E. Panther, F. Di Virgilio, The P2X7 receptor: a key player in IL-1 processing and release, *J. Immunol.* 176 (2006) 3877–3883.
- [49] M.J. Bours, E.L. Swennen, F. Di Virgilio, B.N. Cronstein, P.C. Dagnelie, Adenosine 5'-triphosphate and adenosine as endogenous signaling molecules in immunity and inflammation, *Pharmacol. Ther.* 112 (2006) 358–404.
- [50] S. Gallucci, P. Matzinger, Danger signals: SOS to the immune system, *Curr. Opin. Immunol.* 13 (2001) 114–119.
- [51] F. Aswad, H. Kawamura, G. Dennert, High sensitivity of CD4⁺CD25⁺ regulatory T cells to extracellular metabolites nicotinamide adenine dinucleotide and ATP: a role for P2X7 receptors, *J. Immunol.* 175 (2005) 3075–3083.
- [52] J.M. Labasi, N. Petrushova, C. Donovan, S. McCurdy, P. Lira, M.M. Payette, W. Brissette, J.R. Wicks, L. Audoly, C.A. Gabel, Absence of the P2X(7) receptor alters

- leukocyte function and attenuates an inflammatory response, *J. Immunol.* 168 (2002) 6436–6445.
- [53] H. Kawamura, F. Aswad, M. Minagawa, S. Govindarajan, G. Dennert, P2X7 receptors regulate NKT cells in autoimmune hepatitis, *J. Immunol.* 176 (2006) 2152–2160.
- [54] L. Chen, C.F. Brosnan, Exacerbation of experimental autoimmune encephalomyelitis in P2X7R^{-/-} mice: evidence for loss of apoptotic activity in lymphocytes, *J. Immunol.* 176 (2006) 3115–3126.
- [55] E. Adinolfi, C. Pizzirani, M. Idzko, E. Panther, J. Norgauer, F. Di Virgilio, D. Ferrari, P2X7 receptor: death or life? *Purinergic Signalling* 1 (2005) 219–227.
- [56] L. Gudipaty, B.D. Humphreys, G. Buell, G.R. Dubyak, Regulation of P2X(7) nucleotide receptor function in human monocytes by extracellular ions and receptor density, *Am. J. Physiol.* 280 (2001) C943–C953.
- [57] G. Burnstock, G.E. Knight, Cellular distribution and functions of P2 receptor subtypes in different systems, *Int. Rev. Cyt.* 240 (2004) 31–304.
- [58] A. Muscella, M.G. Elia, S. Greco, C. Storelli, S. Marsigliante, Activation of P2Y2 receptor induces c-FOS protein through a pathway involving mitogen-activated protein kinases and phosphoinositide 3-kinases in HeLa cells, *J. Cell. Physiol.* 195 (2003) 234–240.
- [59] C. Kennedy, A.D. Qi, C.L. Herold, T.K. Harden, R.A. Nicholas, ATP, an agonist at the rat P2Y(4) receptor, is an antagonist at the human P2Y(4) receptor, *Mol. Pharmacol.* 57 (2000) 926–931.
- [60] J.D. Lich, J.C. Arthur, J.P. Ting, Cryopyrin: in from the cold, *Immunity* 24 (2006) 241–243.
- [61] E. Meylan, J. Tschopp, M. Karin, Intracellular pattern recognition receptors in the host response, *Nature* 442 (2006) 39–44.