



Thermodynamic stability of bacteriorhodopsin mutants measured relative to the bacterioopsin unfolded state[☆]

Zheng Cao^a, Jonathan P. Schleich^b, Chiwook Park^b, James U. Bowie^{a,*}

^a Department of Chemistry and Biochemistry, UCLA DOE Institute of Genomics and Proteomics, Molecular Biology Institute, University of California, Los Angeles, Boyer Hall, 611 Charles E. Young Dr. E, Los Angeles, CA 90095-1570, USA

^b Department of Medicinal Chemistry and Molecular Pharmacology, Purdue University, 575 Stadium Mall Drive, West Lafayette, IN 47907-2091, USA

ARTICLE INFO

Article history:

Received 25 May 2011

Received in revised form 8 July 2011

Accepted 15 August 2011

Available online 22 August 2011

Keywords:

Membrane protein

Protein folding

Folding kinetics

SDS

ABSTRACT

The stability of bacteriorhodopsin (bR) has often been assessed using SDS unfolding assays that monitor the transition of folded bR (bR_f) to unfolded (bR_u). While many criteria suggest that the unfolding curves reflect thermodynamic stability, slow retinal (RET) hydrolysis during refolding makes it impossible to perform the most rigorous test for equilibrium, *i.e.*, superimposable unfolding and refolding curves. Here we made a new equilibrium test by asking whether the refolding rate in the transition zone is faster than RET hydrolysis. We find that under conditions we have used previously, refolding is in fact slower than hydrolysis, strongly suggesting that equilibrium is not achieved. Instead, the apparent free energy values reported previously are dominated by unfolding rates. To assess how different the true equilibrium values are, we employed an alternative method by measuring the transition of bR_f to unfolded bacterioopsin (bO_u), the RET-free form of unfolded protein. The bR_f -to- bO_u transition is fully reversible, particular when we add excess RET. We compared the difference in unfolding free energies for 13 bR mutants measured by both assays. For 12 of the 13 mutants with a wide range of stabilities, the results are essentially the same within experimental error. The congruence of the results is fortuitous and suggests the energetic effects of most mutations may be focused on the folded state. The bR_f -to- bO_u reaction is inconvenient because many days are required to reach equilibrium, but it is the preferable measure of thermodynamic stability. This article is part of a Special Issue entitled: Protein Folding in Membranes.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Measuring protein stability requires a method to drive the folding equilibrium in favor of the unfolded state. For water-soluble proteins this is typically accomplished using urea, Gdn-HCl or temperature. To our knowledge, thermal unfolding has not been found to be reversible for any helical membrane proteins [1–6]. Urea and Gdn-HCl have been particularly effective for beta-barrel membrane proteins [7] and have been found to reversibly unfold a few large helical membrane proteins [7,8]. An alternative to urea and Gdn-HCl is to use a denaturing detergent [9,10]. SDS has the advantage that the protein is maintained in a micelle environment, which leads to maintenance of considerable secondary structure and may therefore be a better mimic of the more structured unfolded state in a bilayer [9,11–14].

Bacteriorhodopsin can be completely refolded from an SDS unfolded state [11,15,16]. Titration of bR with SDS from a DMPC/CHAPSO or DMPC/CHAPS solution, leads to bR_f -to- bR_u unfolding curves that are

reasonably stable over the course of an hour (see below). Moreover, kinetic constants in DMPC/CHAPS conditions indicate that unfolding to bR_u should reach equilibrium in a matter of minutes [17–19]. The unfolding curves are also well modeled by a two-state equilibrium and the extrapolated kinetic constants of the unfolding and refolding reactions are consistent with extracted equilibrium constants within the transition zones [17,18]. Thus, we have assumed that the unfolding transitions reflect a folding equilibrium, in spite of the fact that the most rigorous test of equilibrium is not possible, *i.e.*, overlap of the unfolding and refolding curves. We have now come to the conclusion, however, that true equilibrium cannot be achieved under the DMPC/CHAPSO conditions we have used in the past (16 mM DMPC, 6 mM CHAPSO) [20–25], because RET hydrolysis exceeds the rate of refolding near the midpoint of the observed unfolding transition (see below).

Instead of measuring the bR_f -to- bR_u transition that is accessed from unfolding at short time scales, it is also possible to measure the bR_f -to- bO_u transition if the reaction is allowed to proceed for many days. The advantage of the bR_f -to- bO_u transition is that it can be rigorously shown that the reaction is at equilibrium in the transition zones because the folding and unfolding curves are essentially the same [26]. To measure the error associated with the equilibrium

[☆] This article is part of a Special Issue entitled: Protein Folding in Membranes.

* Corresponding author. Tel. +1 310 206 4747; fax: +1 310 206 4749.

E-mail address: bowie@mbi.ucla.edu (J.U. Bowie).

assumption for the bR_f -to- bR_u transition, we compared the effects of mutations on both reactions. Fortunately, the results are remarkably similar for both measurements.

2. Materials and methods

2.1. Materials

All bR variants were prepared as described [20,27,28]. 1,2-Dimyristoyl-sn-glycerol-3-phosphocholine (DMPC) was purchased from Avanti Polar Lipids. 3((3-cholamidopropyl)dimethylammonio)-2-hydroxy-1-propanesulphonate (CHAPSO) was purchased from Anatrace. Sodium dodecyl sulfate (SDS) and all-*trans* retinal (RET) were purchased from Sigma-Aldrich.

2.2. Methods

2.2.1. bR_f -to- bR_u unfolding and bR_u -to- bR_f refolding assays

The bR_f -to- bR_u unfolding assays were performed as described by Faham S. et al. [20]. Purple membrane was dissolved in a 2500 μ L solution containing 15 mM DMPC, 6 mM CHAPSO and 10 mM sodium phosphate [pH 6.3]. The final concentration of the bR protein was in a range of 3–7 μ M. After equilibration in dark for 1 h, the solution was titrated by using an SDS titrant at room temperature in a 1-cm cuvette stirred with a magnetic stir bar. The SDS titrant contained 20% (w/v) SDS, 15 mM DMPC, 6 mM CHAPSO and 10 mM sodium phosphate [pH 6.3]. 10 μ L of the SDS titrant was added every 3 min during the titration.

To test the effect of RET hydrolysis on the time evolution of the bR_f -to- bR_u unfolding curve, a series of 200 μ L solutions containing 3.7 μ M wild-type bR, 15 mM DMPC, 6 mM CHAPSO, 10 mM sodium phosphate [pH 6.3] and different amounts of SDS varying from 5 to 71 mM were made at the same time and then loaded on a 96-well micro-plate (Thermo Scientific). The absorbance of each solution at 560 nm was monitored using a SpectraMax M5 plate reader (Molecular Devices).

To examine the bR_u -to- bR_f refolding reactions, purple membrane was dissolved to make a final solution containing 37 μ M bR, 15 mM DMPC, 6 mM CHAPSO, 10 mM sodium phosphate [pH 6.3] and 66.5 mM SDS. 3 min later, the absorbance at 560 nm disappeared and absorbance at 440 nm reached the maximal value. 20 μ L of the unfolded bR solution was then mixed with a series of 180 μ L solutions. The final solutions contained 3.7 μ M wild-type bR, 15 mM DMPC, 6 mM CHAPSO, 10 mM sodium phosphate [pH 6.3] and different amounts of SDS varying from 5.2 to 66.5 mM.

2.2.2. bR_f -to- bO_u unfolding and bO_u -to- bR_f refolding assays

The bR_f -to- bO_u unfolding and bO_u -to- bR_f refolding assays were performed similarly to Chen and Gouaux [26]. The main difference from their experiments was that we equilibrated each sample for much longer times.

For the unfolding assays, a stock solution was prepared by dissolving purple membrane in a stock solution containing 20.625 mM DMPC, 22 mM CHAPSO and 13.75 mM sodium phosphate [pH 6.3]. For experiments with added RET, 15.4 μ M all-*trans* RET was included in the stock solution. 160 μ L aliquots of the stock solution were combined with 60 μ L SDS solutions at various concentrations. The final solutions contained 1.5–3 μ M bR, 15 mM DMPC, 16 mM CHAPSO, 10 mM sodium phosphate [pH 6.3], 0 or 11.2 μ M all-*trans* RET and SDS varying from 8 to 138 mM.

For the refolding assays, bR was first unfolded in a buffer containing 15 mM DMPC, 16 mM CHAPSO, 10 mM sodium phosphate [pH 6.3] and 104 mM SDS. After equilibration for 1 h in dark, the absorbance at 560 nm disappeared and the absorbance at 390 nm reached the maximal value. Then, the unfolded bR solution was mixed with a series of DMPC/CHAPSO/SDS/RET/sodium phosphate [pH 6.3] solutions with varying SDS concentrations. The final solutions contained 1.5–3 μ M bR, 15 mM DMPC, 16 mM CHAPSO, 10 mM sodium phosphate [pH 6.3], 0 or

11.2 μ M all-*trans* RET and different amounts of SDS varying from 13 to 79 mM.

After the samples for unfolding and refolding assays were prepared, all the samples without added RET were equilibrated in dark at room temperature for 212 h (~9 days) and those with added RET were equilibrated in dark at room temperature for 96 h (~4 days). 200 μ L of each unfolding and refolding sample was loaded on a 96-well UV-star micro-plate (Greiner Bio-One). Absorbance at 560 nm and fluorescence emission at 335 nm (excitation at 290 nm) of each solution were measured by SpectraMax M5 plate reader (Molecular Devices).

For unfolding experiments without added RET, the unfolding curves can be described by

$$\text{Abs}_{560} = \text{Abs}_{560}^0 \cdot F_f, \quad (1)$$

where Abs_{560}^0 is the absorbance of the subdenaturant line and its extension over the experimental X_{SDS} range, which represents the theoretical absorbance if all the bR is folded and is assumed to be linearly dependent on X_{SDS} :

$$\text{Abs}_{560}^0 = aX_{\text{SDS}} + b, \quad (2)$$

and F_f is the fraction of the folded state in each bR variant, i.e. $F_f = [bR_f]/c$, where c is the total concentration of each bR variant in units of μ M. Since the unfolding free energy calculated for a standard state of 1 μ M is defined as

$$\Delta G_U = -RT \ln([bO_u] \cdot [\text{RET}] / [bR_f]) = -RT \ln[c(1 - F_f)^2 / F_f],$$

where RT is 0.592 kcal mol⁻¹, F_f can be written as a function of ΔG_U and c :

$$F_f = \frac{2 + \frac{\exp[-\Delta G_U/(RT)]}{c} - \sqrt{\frac{\exp[-2\Delta G_U/(RT)]}{c^2} + \frac{4\exp[-\Delta G_U/(RT)]}{c}}}{2}. \quad (3)$$

We assumed that ΔG_U has a linear relationship with X_{SDS} :

$$\Delta G_U = m(X_{\text{SDS}} - C_m) - RT \ln(0.5c).$$

Then Eq. (3) can be re-written as

$$F_f = \frac{2 + \exp\left[\frac{-[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right] - \sqrt{\frac{\exp\left[\frac{-2[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right]}{2} + 4 \cdot \exp\left[\frac{-[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right]}}{2}. \quad (4)$$

By plugging Eqs. (2) and (4) into Eq. (1), we derived the equation of Abs_{560} in the expression of X_{SDS} and used this equation to fit each unfolding curve without added RET:

$$\text{Abs}_{560} = (aX_{\text{SDS}} + b) \cdot \frac{2 + \exp\left[\frac{-[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right] - \sqrt{\frac{\exp\left[\frac{-2[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right]}{2} + 4 \cdot \exp\left[\frac{-[m(X_{\text{SDS}} - C_m) + \ln 2]}{RT}\right]}}{2}. \quad (5)$$

Parameters a , b , m and C_m were fit using Kaleidagraph 4.1. The unfolding free energies for wild-type and mutant were compared by subtracting ΔG_U^{WT} from $\Delta G_U^{\text{mutant}}$ at the midpoint of the wild-type transition, $X_{\text{SDS}} = C_m^{\text{WT}} = 0.572$.

For unfolding experiments with added RET, the RET concentration was held constant, so the RET concentration was combined with the equilibrium constant, giving the unfolding reaction a pseudo 1:1 stoichiometry, i.e. $\Delta G_U = -RT \ln([bO_u]/[bR_f]) = -RT \ln[(1 - F_f)/F_f]$.

F_f can be written as

$$F_f = \frac{1}{1 + \exp[-\Delta G_U/(RT)]}. \quad (6)$$

We assume that ΔG_U has a linear relationship with X_{SDS} :

$$\Delta G_U = m(X_{SDS} - C_m).$$

Then F_f can be re-written as

$$F_f = \frac{1}{1 + \exp[-m \cdot (X_{SDS} - C_m)/(RT)]}. \quad (7)$$

By plugging Eqs. (2) and (7) into Eq. (1), we obtained the equation of Abs_{560} in terms of X_{SDS} and used this equation to fit each unfolding curve with added RET:


$$Abs_{560} = \frac{a \cdot X_{SDS} + b}{1 + \exp[-m \cdot (X_{SDS} - C_m)/(RT)]}.$$

Parameters a , b , m and C_m were fit using Kaleidagraph 4.1. The unfolding free energies for wild-type and mutant were compared by subtracting ΔG_U^{WT} from ΔG_U^{mutant} at the midpoint of the wild-type transition, $X_{SDS} = C_m^{WT} = 0.611$.

3. Results

The bR_f -to- bR_u reaction is complicated by the slow hydrolysis of RET, which occurs from the fully unfolded protein with a half life of around 12 min [17]. Because loss of RET is slow until there is a significant fraction of unfolded protein, the unfolding curves are relatively stable over a modest time scale. Fig. 1a shows the change in unfolding curves over time. Minimal change is seen over the first 30 min. These results, combined with measurements of unfolding rates under similar conditions [17,18], which indicate that unfolding should be achieved in a matter of minutes, suggest that the bR_f -to- bR_u unfolding curves can be used to extract equilibrium constants over a useable time scale. Nevertheless, due to RET hydrolysis during refolding, the refolding curves do not superimpose with the unfolding curves (Fig. 1b). At low X_{SDS} , where refolding is rapid, the protein refolds to near completion, but in the transition zones where refolding is slow, RET hydrolysis becomes a significant factor, complicating the interpretation.

We therefore sought a further test of the equilibrium assumption. A simplified reaction scheme for unfolding followed by essentially irreversible RET hydrolysis is given by:



Where k_u is the unfolding rate constant, k_f is the refolding rate constant and k_{hyd} is the RET hydrolysis rate constant. For equilibrium between bR_f and bR_u to be achieved, k_f must be greater than k_{hyd} . To test whether this is true in the transition zone, we rapidly unfolded the protein to bR_u at a high SDS concentration ($X_{SDS} = 0.78$), then jumped back an SDS concentration at the midpoint of the observed transition ($X_{SDS} = 0.67$) and monitored both refolding at 560 nm and hydrolysis at 390 nm. As shown in Fig. 1c, we essentially saw no refolding before hydrolysis was complete. Thus, under the solution conditions we have used for measuring stability, equilibrium cannot be established near the midpoint of the transition and the extent of the observed unfolding must be limited by the unfolding rate.

We therefore measured the unfolding free energy differences for a variety of mutants using an alternative thermodynamic stability




Fig. 1. Effect of RET hydrolysis on the SDS-induced unfolding of wild-type bR. (a) Change of the wild-type bR_f -to- bR_u unfolding curve in time. (b) Plot of normalized absorbance at 560 nm against the SDS mole fraction concentration, X_{SDS} , for the wild-type bR_f -to- bR_u unfolding and bR_u -to- bR_f refolding experiments. (c) Change of absorbances at 390, 440 and 560 nm in time when the wild-type protein was refolded from bR_u at the apparent C_m of the bR_f -to- bR_u transition ($X_{SDS} = 0.673$).

measurement. If the unfolding reactions are left to incubate, a bR_f -to- bO_u equilibrium is attained as described by Chen and Gouaux [26]: $bR_f \leftrightarrow bO_u + RET$. In our hands it took 9 days to reach equilibrium at room temperature as judged by the nearly superimposable folding and unfolding curves monitored by absorbance at 560 nm and fluorescence at 335 nm (Fig. 2a and b). By adding excess RET in the samples, we shortened the equilibration time to 4 days and the folding and unfolding curves monitored by absorbance at 560 nm and fluorescence at 335 nm (Fig. 2c and d) were more superimposable. The SDS concentration at the midpoint of the transition zone, C_m , was much higher for the bR_f -to- bR_u reaction [17,18] than for the bR_f -to- bO_u reaction, which is consistent with the spontaneous hydrolysis of the RET Schiff base in SDS solutions at neutral pH [29]. In addition, the only peaks observed in the absorbance spectra throughout the transition were at 560 nm, corresponding to bR_f , and at 390 nm, corresponding to free RET (Fig. 3). No indications of a contribution from bR_u at 440 nm were evident, which indicates that the product of the unfolding experiment was bO and free RET but not bR_u [15,17,26]. The unfolding curves derived from Abs_{560} and Flu_{335} were essentially the same (Fig. 4), and the absorbance spectra exhibited an isosbestic point (Fig. 3), suggesting that the unfolding of bR_f to bO_u can be described by a two-state model. As the bR_f -to- bO_u reaction




Fig. 2. Equilibrium unfolding (bR_F -to- bO_U) and refolding (bO_U -to- bR_F) of wild-type protein. (a) The unfolding and refolding samples without added RET monitored by absorbance at 560 nm after incubated for ~9 days. (b) The unfolding and refolding samples without added RET monitored by fluorescence at 335 nm after incubated for ~9 days. (c) The unfolding and refolding samples with added RET monitored by absorbance at 560 nm after incubated for ~4 days. (d) The unfolding and refolding samples with added RET monitored by fluorescence at 335 nm after incubation for ~4 days.

should release RET, we expected that the addition of free RET would drive the equilibrium to higher SDS concentrations. Indeed, this is what we observed (Fig. 4), and is consistent with a thermodynamic equilibrium process. The addition of free RET has two advantages: (1) the concentration of RET becomes a constant, simplifying analysis of the unfolding curves and (2) the time it took to reach equilibrium decreases from 9 days to 4 days.

The $\Delta\Delta G_U$ values obtained using the bR_F -to- bO_U transition and $\Delta\Delta G_U^{app}$ values using the bR_F -to- bR_U transition were measured for 13 mutants and are listed in Table 1. We compare measurements at the center of the transition zones for the wild-type protein to minimize extrapolations outside of the observable range of fraction unfolded ($X_{SDS}=0.572$ for the bR_F -to- bO_U transition without added RET, $X_{SDS}=0.611$ for the bR_F -to- bO_U transition in the presence of 11.2 μM RET, and $X_{SDS}=0.673$ for the bR_F -to- bR_U transition). For 12 of the 13 mutants, the $\Delta\Delta G_U^{app}$ and $\Delta\Delta G_U$ values measured using the two transitions bR_F -to- bR_U and bR_F -to- bO_U , respectively, were generally within the experimental error. The correlation between the different

measures of $\Delta\Delta G_U$ for the 12 mutants is shown in Fig. 5, illustrating the close correspondence between the two methods. An ideal correlation would have an intercept at 0 kcal mol⁻¹ and a slope of 1.0. For the fitted line, the intercept is (-0.12 ± 0.09) kcal mol⁻¹ and the slope is 1.08 ± 0.06 .

4. Discussion

While the bR_F -to- bR_U unfolding transition apparently reaches a steady state rapidly, RET hydrolysis and slow refolding preclude establishment of a true equilibrium under the conditions we have used. In contrast, the bR_F -to- bO_U equilibrium is stable. Nevertheless, when we compare the effects of 12 of the 13 mutations on the two reactions, very similar results are obtained for most mutations. The origin of the fortuitous congruence of the two measures is unclear. One possibility is that mutations largely alter k_U , but not k_F . This could




Fig. 3. Absorbance spectra of the wild-type protein as a function of SDS concentration. The spectra were taken at $X_{SDS}=0.315$, 0.572 and 0.734, where the protein was completely folded, 50% unfolded and completely unfolded, respectively.




Fig. 4. Wild-type bR_F -to- bO_U unfolding curve. Plot of the fraction of the native-state protein, F_N , derived by monitoring absorbance and fluorescence for the reactions with and without added RET. Superposition of the curves is consistent with a two-state model for SDS-induced unfolding of bR_F to bO_U . The transition shifts to higher SDS concentration in the presence of added RET, consistent with an equilibrium reaction that releases RET. Under both conditions with and without added RET, 3 μM of wild-type protein was included in the samples.

Table 1 $\Delta\Delta G_U$ of bR variants tested from the bR_F-to-bO_u and bR_F-to-bR_u reactions at certain X_{SDS} .

Single mutants	$\Delta\Delta G_U^a$ (bR _F -to-bO _u without added RET)	$\Delta\Delta G_U^a$ (bR _F -to-bO _u with 11.2 μ M RET)	$\Delta\Delta G_U^a$ (bR _F -to-bR _u)		
X_{SDS}	0.572 ^b	0.611 ^b	0.572	0.611	0.673 ^b
K41A	–	-1.2 ± 0.3	-1.4 ± 0.3	-1.4 ± 0.3	-1.4 ± 0.2
F42A	-2.6 ± 0.5	-2.2 ± 0.3	-2.1 ± 0.3	-2.1 ± 0.2	-2.0 ± 0.2
Y43A	-1.9 ± 0.3	-2.1 ± 0.3	-1.9 ± 0.4	-1.9 ± 0.3	-1.6 ± 0.2
I45A	-2.2 ± 0.4	-2.1 ± 0.2	-2.1 ± 0.3	-2.1 ± 0.3	-1.9 ± 0.3
T46A	–	-2.8 ± 0.3	-2.3 ± 0.3	-2.3 ± 0.3	-2.2 ± 0.3
T47A	-1.1 ± 0.3	-0.9 ± 0.2	-1.6 ± 0.3	-1.4 ± 0.2	-1.1 ± 0.2
V49A	-0.3 ± 0.4	-0.3 ± 0.3	-0.7 ± 0.2	-0.6 ± 0.2	-0.3 ± 0.2
I52A	-2.0 ± 0.3	-1.6 ± 0.2	-2.0 ± 0.2	-1.9 ± 0.2	-1.5 ± 0.1
F54A	–	-0.9 ± 0.3	-0.5 ± 0.2	-0.4 ± 0.2	-0.4 ± 0.1
M56A	–	1.6 ± 0.3	2.0 ± 0.2	1.9 ± 0.2	1.6 ± 0.1
S59A	–	-0.1 ± 0.2	-0.1 ± 0.2	-0.1 ± 0.2	-0.1 ± 0.1
M60A	–	-1.3 ± 0.3	-1.1 ± 0.3	-1.1 ± 0.3	-1.0 ± 0.2
P50A	-1.1 ± 0.3	-1.0 ± 0.2	0.7 ± 0.4	0.7 ± 0.3	0.6 ± 0.1

^a Unit of $\Delta\Delta G_U$: kcal mol^{–1}.^b C_m of the wild-type bR for each reaction.

happen if the mutations have large effects on the free energy of the native state, but minimal effects on the transition state or unfolded state free energies. The effects of one mutant, P50A, were very different when measured by the two methods. It was found to be significantly stabilizing in the bR_F-to-bR_u transition, but destabilizing in the bR_F-to-bO_u transition. P50 is in the center of a kinked helix in the folded bR structure, so it is perhaps not surprising that there might be long-range effects imparted by this mutation that differentially alter unfolding transition state.

Booth and co-workers have found excellent correspondence between equilibrium and kinetic measurements [17,18]. Their conditions are slightly different than the ones we have used, however, employing CHAPS instead of CHAPSO and the CHAPS is used at a higher concentration. We have observed that increasing CHAPSO concentrations from 6 to 16 mM increases refolding rates (unpublished result), which could then lead to equilibrium for the bR_F-to-bR_u transition. Nevertheless, the

bR_F-to-bO_u transition is the more reliable way to measure the effects of mutations on thermodynamic stability.

Acknowledgements

This work was supported by NIH grant R01GM063919 to JUB.

References

- [1] P.E. Morin, D. Diggs, E. Freire, Thermal stability of membrane-reconstituted yeast cytochrome c oxidase, *Biochemistry* 29 (1990) 781–788.
- [2] L.K. Thompson, A.F. Miller, C.A. Buser, J.C. de Paula, G.W. Brudvig, Characterization of the multiple forms of cytochrome b559 in photosystem II, *Biochemistry* 28 (1989) 8048–8056.
- [3] L.R. Maneri, P.S. Low, Structural stability of the erythrocyte anion transporter, band 3, in different lipid environments. A differential scanning calorimetric study, *J. Biol. Chem.* 263 (1988) 16170–16178.
- [4] G.Q. Chen, E. Gouaux, Reduction of membrane protein hydrophobicity by site-directed mutagenesis: introduction of multiple polar residues in helix D of bacteriorhodopsin, *Protein Eng.* 10 (1997) 1061–1066.
- [5] T.W. Kahn, J.M. Sturtevant, D.M. Engelman, Thermodynamic measurements of the contributions of helix-connecting loops and of retinal to the stability of bacteriorhodopsin, *Biochemistry* 31 (1992) 8829–8839.
- [6] T. Haltia, E. Freire, Forces and factors that contribute to the structural stability of membrane proteins, *Biochim. Biophys. Acta* 1241 (1995) 295–322.
- [7] E.A. Roman, J.M. Argüello, F.L. González Flecha, Reversible unfolding of a thermophilic membrane protein in phospholipid/detergent mixed micelles, *J. Mol. Biol.* 397 (2010) 550–559.
- [8] H.E. Findlay, N.G. Rutherford, P.J.F. Henderson, P.J. Booth, Unfolding free energy of a two-domain transmembrane sugar transport protein, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 18451–18456.
- [9] F.W. Lau, J.U. Bowie, A method for assessing the stability of a membrane protein, *Biochemistry* 36 (1997) 5884–5892.
- [10] R. Renthal, An unfolding story of helical transmembrane proteins, *Biochemistry* 45 (2006) 14559–14566.
- [11] K.S. Huang, H. Bayley, M.J. Liao, E. London, H.G. Khorana, Refolding of an integral membrane protein. Denaturation, renaturation, and reconstitution of intact bacteriorhodopsin and two proteolytic fragments, *J. Biol. Chem.* 256 (1981) 3802–3809.
- [12] B. Jirgensons, Factors determining the reconstructive denaturation of proteins in sodium dodecyl sulfate solutions. Further circular dichroism studies on structural reorganization of seven proteins, *J. Protein Chem.* 1 (1982) 71–84.
- [13] B. Jirgensons, S. Capetillo, Effect of sodium dodecyl sulfate on circular dichroism of some nonhelical proteins, *Biochim. Biophys. Acta* 214 (1970) 1–5.
- [14] M.M. Snel, R. Kaptein, B. de Kruijff, Interaction of apocytochrome c and derived polypeptide fragments with sodium dodecyl sulfate micelles monitored by photochemically induced dynamic nuclear polarization 1H NMR and fluorescence spectroscopy, *Biochemistry* 30 (1991) 3387–3395.
- [15] E. London, H.G. Khorana, Denaturation and renaturation of bacteriorhodopsin in detergents and lipid-detergent mixtures, *J. Biol. Chem.* 257 (1982) 7003–7011.
- [16] T. Marti, Refolding of bacteriorhodopsin from expressed polypeptide fragments, *J. Biol. Chem.* 273 (1998) 9312–9322.
- [17] P. Curnow, P.J. Booth, Combined kinetic and thermodynamic analysis of alpha-helical membrane protein unfolding, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 18970–18975.
- [18] P. Curnow, P.J. Booth, The transition state for integral membrane protein folding, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 773–778.




Fig. 5. Correlation between $\Delta\Delta G_U^{app}$ values measured for the bR_F-to-bR_u transition and the $\Delta\Delta G_U$ values measured for the bR_F-to-bO_u transition. Square symbols refer to values measured without added RET and circle symbols refer to values measured with added RET. The open symbols refer to values for the P50A mutation, which is an outlier. The least squares fit line through all the filled symbols has a slope of 1.08 ± 0.06 and an intercept of -0.12 ± 0.09 kcal mol^{–1}. The correlation coefficient is 0.97.

- [19] P.J. Booth, S.L. Flitsch, L.J. Stern, D.A. Greenhalgh, P.S. Kim, H.G. Khorana, Intermediates in the folding of the membrane protein bacteriorhodopsin, *Nat. Struct. Biol.* 2 (1995) 139–143.
- [20] S. Faham, D. Yang, E. Bare, S. Yohannan, J.P. Whitelegge, J.U. Bowie, Side-chain contributions to membrane protein structure and stability, *J. Mol. Biol.* 335 (2004) 297–305.
- [21] S. Yohannan, S. Faham, D. Yang, D. Grosfeld, A.K. Chamberlain, J.U. Bowie, A C alpha-H...O hydrogen bond in a membrane protein is not stabilizing, *J. Am. Chem. Soc.* 126 (2004) 2284–2285.
- [22] S. Yohannan, S. Faham, D. Yang, J.P. Whitelegge, J.U. Bowie, The evolution of trans-membrane helix kinks and the structural diversity of G protein-coupled receptors, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 959–963.
- [23] S. Yohannan, D. Yang, S. Faham, G. Boulting, J. Whitelegge, J.U. Bowie, Proline substitutions are not easily accommodated in a membrane protein, *J. Mol. Biol.* 341 (2004) 1–6.
- [24] N.H. Joh, A. Min, S. Faham, J.P. Whitelegge, D. Yang, V.L. Woods, et al., Modest stabilization by most hydrogen-bonded side-chain interactions in membrane proteins, *Nature* 453 (2008) 1266–1270.
- [25] N.H. Joh, A. Oberai, D. Yang, J.P. Whitelegge, J.U. Bowie, Similar energetic contributions of packing in the core of membrane and water-soluble proteins, *J. Am. Chem. Soc.* 131 (2009) 10846–10847.
- [26] G.Q. Chen, E. Gouaux, Probing the folding and unfolding of wild-type and mutant forms of bacteriorhodopsin in micellar solutions: evaluation of reversible unfolding conditions, *Biochemistry* 38 (1999) 15380–15387.
- [27] S.W. Cline, W.F. Doolittle, Efficient transfection of the archaeobacterium *Halobacterium halobium*, *J. Bacteriol.* 169 (1987) 1341–1344.
- [28] D. Oesterhelt, W. Stoerkenius, Isolation of the cell membrane of *Halobacterium halobium* and its fractionation into red and purple membrane, *Methods Enzymol.* 31 (1974) 667–678.
- [29] A. Cooper, S.F. Dixon, M.A. Nutley, J.L. Robb, Mechanism of retinal Schiff base formation and hydrolysis in relation to visual pigment photolysis and regeneration: resonance Raman spectroscopy of a tetrahedral carbinolamine intermediate and oxygen-18 labeling of retinal at the metarhodopsin stage in photoreceptor membranes, *J. Am. Chem. Soc.* 109 (1987) 7254–7263.