

Erratum

Erratum to ‘Calorimetrically obtained information about the efficiency of ectoine synthesis from glucose in *Halomonas elongata*’ [Biochimica et Biophysica Acta, 1527 (2001) 4–10][☆]

T. Maskow *, W. Babel

UFZ Centre for Environmental Research Leipzig-Halle GmbH, Department of Environmental Microbiology, Permoserstrasse 15, D-04318 Leipzig, Germany

The term t^2 was missing in the middle equation of row 3 in Table 1. For completeness, Table 1 is reproduced below.

Table 1
 Models for the evolution of the formation rates

Evolution of rate of ectoine formation	Solutions of Eq. 10	Number of parameters needed to adjust
$r_M = \text{const.} \neq r_M^0$	$\Phi_M = \Phi_M^0 e^{-Dt + (r_M/D)(1 - e^{-Dt})}$	1
$r_M = r_M^0 + \dot{r}_M t$	$\Phi_M = \Phi_M^0 + (\dot{r}_M/D^2)(e^{-Dt} + Dt - 1)$	1
$r_M = r_M^0 + \dot{r}_M t + \ddot{r}_M t^2$	$\Phi_M = \Phi_M^0 + (\ddot{r}_M/D)t^2 + ((\dot{r}_M/D^2) - (2\ddot{r}_M/D^3))(e^{-Dt} + Dt - 1)$	2

Φ_M^0 , r_M^0 , concentration (mol l⁻¹) and formation rate (mol l⁻¹ h⁻¹) during steady state phase, respectively; $\dot{r}_M$, $\ddot{r}_M$, first (mol l⁻¹ h⁻²) and second derivation (mol l⁻¹ h⁻³) of formation rate versus time, respectively; t , time (h); M , biomass or ectoine.

[☆] PII of original article S0304-4165(01)00115-5.

* Corresponding author. Fax: +49-341-235-2247.

E-mail address: maskow@umb.ufz.de (T. Maskow).