

Review

Matrix regulators in neural stem cell functions[☆]

Anna Wade^a, Andrew McKinney^a, Joanna J. Phillips^{a,b,c,*}
^a Department of Neurological Surgery, UCSF, San Francisco, CA, USA

^b Brain Tumor Research Center, UCSF, San Francisco, CA, USA

^c Neuropathology, Department of Pathology, UCSF, San Francisco, CA, USA


ARTICLE INFO

Article history:

Received 16 November 2013

Received in revised form 9 January 2014

Accepted 10 January 2014

Available online 18 January 2014

Keywords:

Extracellular matrix

Proteoglycan

HSPG

Sulfs

Neural stem cell

NSC

ABSTRACT

Background: Neural stem/progenitor cells (NSPCs) reside within a complex and dynamic extracellular microenvironment, or niche. This niche regulates fundamental aspects of their behavior during normal neural development and repair. Precise yet dynamic regulation of NSPC self-renewal, migration, and differentiation is critical and must persist over the life of an organism.

Scope of review: In this review, we summarize some of the major components of the NSPC niche and provide examples of how cues from the extracellular matrix regulate NSPC behaviors. We use proteoglycans to illustrate the many diverse roles of the niche in providing temporal and spatial regulation of cellular behavior.

Major conclusions: The NSPC niche is comprised of multiple components that include; soluble ligands, such as growth factors, morphogens, chemokines, and neurotransmitters, the extracellular matrix, and cellular components. As illustrated by proteoglycans, a major component of the extracellular matrix, the NSPC, niche provides temporal and spatial regulation of NSPC behaviors.

General significance: The factors that control NSPC behavior are vital to understand as we attempt to modulate normal neural development and repair. Furthermore, an improved understanding of how these factors regulate cell proliferation, migration, and differentiation, crucial for malignancy, may reveal novel anti-tumor strategies. This article is part of a Special Issue entitled Matrix-mediated cell behaviour and properties.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

The patterning and development of the central nervous system (CNS) are a complex and highly organized process. In successive waves of proliferation, migration, and differentiation neural stem cells populate the CNS and give rise to neurons, astrocytes, and oligodendrocytes. During cortical neurogenesis, neural stem cells (neuroepithelial stem cells and subsequently radial glia cells slightly later in development) in the ventricular zone initially expand via symmetric division and then shift toward an asymmetric division mode to give rise to intermediate progenitor cells [47,94]. While the process of cortical gliogenesis is less well understood (see review [91]), the progenitor cells must also undergo tightly regulated self-renewal, migration, and differentiation. In the adult mammalian brain, neural stem and progenitor cells persist and can participate in neurogenesis and gliogenesis

under both normal and disease conditions [32,103,104]. Neural stem cells exist in the ventricular–subventricular zone, in the walls of the lateral ventricle, and in the subgranular zone, at the junction of the hilus and dentate gyrus [68,125]. In addition, more differentiated oligodendroglial precursor cells (OPCs) reside in the brain parenchyma and can rapidly respond to injury [96,123]. Throughout life, the behavior of neural precursor cells requires exquisite regulation.

Neural precursors reside within a complex and dynamic extracellular microenvironment, or niche. This niche regulates fundamental aspects of their behavior [5] and includes the brain extracellular matrix, blood vessels, microglia, and more differentiated neural cells (Fig. 1). Extracellular factors such as growth factors, morphogens, chemokines, neurotransmitters, and the extracellular matrix are critical components of this niche. Indeed, extracellular cues are responsible for patterning of the nervous system, and they generate critical domain-specific signals to radial glial cells that extend from the ventricular surface to the cortical pial surface during embryogenesis [94]. In this review, we focus on the factors present in the cellular microenvironment that regulate neural stem cells and their derived intermediate progenitor cells, collectively referred to as neural stem/progenitor cells (NSPCs). An understanding of how extrinsic cues regulate the spatial, temporal, and even subcellular signaling in NSPCs will improve our understanding of normal neural development and, as dysregulation of this process can lead to disease, may contribute to the development of novel therapeutic approaches to neurologic disease.

Abbreviations: NSPC, neural stem/progenitor cell; HSPG, heparan sulfate proteoglycan; CSPG, chondroitin sulfate proteoglycan; EGF, epidermal growth factor; FGF, fibroblast growth factor; Shh, sonic hedgehog; BMP, bone morphogenic protein; OPC, oligodendroglial precursor cells

[☆] This article is part of a Special Issue entitled Matrix-mediated cell behaviour and properties.

* Corresponding author at: Neurological Surgery, 1450 Third Street, UCSF, San Francisco, CA 94115, USA. Tel.: +1 415 514 4929; fax: +1 415 514 9729.

E-mail address: Joanna.phillips@ucsf.edu (J.J. Phillips).




Fig. 1. Extracellular matrix components within the neural stem/progenitor cell (NSPC) niche. In the adult NSPC niche in the wall of the lateral ventricle, NSPCs (grey) are in close proximity with ependymal cells (yellow), cerebrospinal fluid, blood vessels (pink) and more differentiated neural cells (green). There is also a diverse extracellular matrix (inset). Heparan sulfate and chondroitin sulfate proteoglycans (1,2) are present either at the plasma membrane, attached via a transmembrane domain or a GPI-link, or are secreted. At the plasma membrane, proteoglycans can sequester ligands (3) to alter local ligand concentrations or prevent degradation, promote RTK signaling by acting as co-receptors, promote cell adhesion by interacting with the extracellular matrix and with integrins (5), and facilitate cell migration. In the extracellular environment proteoglycans can be enzymatically modified to generate soluble, biologically active fragments and the SULFs (4) remove 6O-sulfates from HS. Integrins (5) facilitate cell–extracellular matrix adhesion in the NSPC niche through direct interactions with extracellular components including laminin (6), glycan binding proteins (7) and glycoproteins such as tenascin C (8). Figure is not to scale.

2. Components of the extracellular matrix of neural stem/progenitor cells

Diverse components make up the NSPC microenvironment, or niche, and many have the potential to regulate cell behaviors. Some of these components are reviewed below.

Table 1
Extracellular factors implicated in NSPC lineage specification and signaling.

Extracellular factor	Commonly reported role in NSPCs	Reference
BMP	Differentiation, dorsal morphogen; quiescence	[41,77,89]
EGF	Proliferation at onset of gliogenesis	[21,76]
FGF1 and FGF2	Proliferation	[76,113]
GABA	Differentiation, decreased proliferation	[80,110,117]
Noggin	Proliferation, blocks BMP	[77]
Notch	Inhibit neuronal differentiation; maintenance	[2,24,25,30,55]
PDGF-AA	Proliferation of NSPC early and OPCs later in development	[50]
SDF1	Recruitment to vasculature, adhesion and motility	[64,105]
SHH	Proliferation; ventral specification; determinant of cell fate	[18,51,54,71,124]
VEGF	Proliferation; neurogenesis	[22,59,74]
VCAM	Adhesion, maintenance	[65]
Wnt	Self-renewal	[26,61,75]

2.1. Growth factors, morphogens, chemokines, and neurotransmitters

Extracellular ligands, many of which are soluble, are a critical component of the regulatory network necessary for NSPC maintenance of self-renewal, differentiation, cell adhesion, and migration. Some of the many extracellular factors implicated in NSPC lineage specification and signaling include fibroblast growth factor (FGF), epidermal growth factor (EGF), stromal derived factor 1 (SDF1), sonic hedgehog (SHH), bone morphogenic protein (BMP), Notch, and GABA (see Table 1 and associated references).

2.2. Proteoglycans

Proteoglycans, consisting of a core protein and covalently attached glycosaminoglycan (GAG) chains, are present in the extracellular microenvironment and on the cell surface, often as GPI-linked or transmembrane proteins. Due to their structural complexity, they can bind diverse extracellular factors, including signaling molecules, membrane proteins, and components of the extracellular matrix (ECM) [13]. As a result, proteoglycans play a vital role in cell–cell and cell–ECM signaling in the CNS. The most common types of proteoglycans in the brain contain heparan sulfate (HS), chondroitin sulfate (CS) or dermatan sulfate (DS) side chains. Another major constituent of the brain ECM is hyaluronan (or hyaluronic acid, HA). Composed of GAG chains without a protein core, HA is released from the cell where it interacts with ligands and matrix components and helps to regulate cell signaling

[90]. On NSPCs, HA binds a number of proteoglycans including neurocan (NCAN), aggrecan (ACAN), and versican (VCAN) [1].

2.3. Glycoproteins

Glycoproteins are a diverse group of proteins that contain one or more oligosaccharide chains (glycans), and they include integral membrane proteins, secreted extracellular proteins, and cytosolic proteins. Tenascin C (TNC) is an extracellular matrix glycoprotein that is highly expressed by NSPCs located in the brain and spinal cord during development and in the adult [42,43,63]. In the spinal cord, genetic ablation of TNC mediates deficits in the NSPC compartment resulting in sustained generation and delayed migration of astrocytes due to altered responsiveness to FGF2 and EGF [63]. Consistent with the dynamic interplay of factors within the NSPC niche, the TNC-mediated alterations in growth factor responsiveness may be due, in part, to secondary alterations in HSPGs [63].

2.4. Glycan binding proteins

Glycan binding proteins, divided broadly into lectins or glycosaminoglycan binding proteins, are thought to mediate many of the biologic functions of glycans [114]. For example, galectin-1, a lectin, promotes the proliferation of NSPCs *in vivo*, and this phenotype is thought to be mediated via galectin-1 binding to glycans on β 1 integrin [57,100,101].

2.5. Integrins

Cell adhesion to ECM is a critical component of the NSPC niche. The major cell surface receptors that mediate this interaction are the integrins. Due to their heterodimeric structure, consisting of α and β subunits, integrins are able to interact with a vast array of extracellular ligands, including collagens, laminin, fibronectin, vitronectin, osteopontin, and transmembrane components such as E-cadherin. Many of these interactions are known to influence NSPC adhesion, migration, cell–cell communication, and cell survival (see review [108]).

2.6. Basal lamina

The basal lamina is an important component of the NSPC niche. Enriched in laminin and collagen 1 [88], the basal lamina helps to regulate adhesion of NSPCs to the extracellular matrix. In the adult, the basal lamina surrounds blood vessels, and blocking α 6 β 1 integrin from binding the laminin receptor resulted in decreased NSPC adhesion and altered proliferation [105]. Laminin also appears to regulate postnatal oligodendrocyte production in the ventricular–subventricular zone [99].

3. Proteoglycan regulation of neural stem and progenitor cell populations

Many NSPC behaviors are dependent on growth factor signaling (Table 1). HSPGs and CSPGs are highly expressed in the NSPC niche, and they are known to modulate growth factor-mediated signaling [53,70,87]. Proteoglycan binding of extracellular ligands can have multiple effects on ligand bioavailability. Binding can sequester the ligand from degradation, establish morphogen gradients, and promote receptor activation by acting as a co-receptor [34]. Indeed, HSPGs stabilize the FGF ligand–receptor complex to promote FGF2 signaling [27,35,45,67,98,122]. Proteoglycans can also regulate cell adhesion and cell migration through interactions with membrane proteins and other components of the ECM, including integrins, laminin, collagen, and fibronectin [70,82]. In addition, proteoglycan intracellular functions are likely to contribute to NSPC regulation as they link extracellular signals with cytoskeletal organization and intracellular signaling [12,33,48,81]. In the following discussion, we use proteoglycans to

illustrate how components of the cellular microenvironment regulate NSPC behavior. In addition, we use human disease phenotypes and data from experimental models to illustrate how a failure in the NSPC niche can lead to disease.

3.1. Heparan sulfate proteoglycans

HSPGs consist of a family of proteoglycans that include members that are primarily transmembrane (e.g. syndecans), GPI-linked to the membrane (e.g. glypicans), secreted from the cell (e.g. HSPG2), or reside in secretory vesicles (e.g. serglycin) [102]. Their ability to regulate growth factor signaling has been demonstrated in multiple systems and for multiple extracellular ligands. In murine embryonic stem cells (mESCs), HS chains help to drive neural precursor cell differentiation via promotion of FGF and BMP signaling [66]. Interestingly, promotion of BMP signaling is associated with an HS-mediated decrease in BMP degradation suggesting that HS binding stabilizes or protects BMP ligand [66]. The syndecans (SDCs) include four members in mammals: SDC1–4. During cortical neurogenesis, SDC1 is highly expressed in the NSPC niche where it promotes NSPC proliferation and progenitor cell maintenance, in part via its ability to promote Wnt signaling in NSPCs [118]. In other studies, SDC3 and SDC4 have been implicated in promoting neural cell migration [14,83].

Perlecan (HSPG2) is an HSPG highly expressed in the basal lamina of the developing neuroepithelium and basal lamina surrounding blood vessels [92]. Mutations in the *Drosophila* homolog of HSPG2 cause defects in FGF and Hedgehog signaling to cause cell cycle arrest and prevents the onset of stem cell proliferation during early development in the fly [Park, 2003 #352, [120]. Similarly, genetic ablation of HSPG2 in the developing murine brain results in decreased Shh signaling, delayed cell cycle progression, and decreased progenitor cell differentiation [44]. Consistent with a role for HSPG2 in neural development, alterations in HSPG2 are associated with cephalic defects in mouse [6,44] and in human with poor development of the forebrain [7]. The growth factor binding ability of HSPG2, including to FGFs [8,115] and Shh [95], is thought to be the primary mechanism by which HSPG2 regulates NSPC behavior.

The specificity and the affinity of HSPG–ligand interactions are largely determined by the HS chains and their sulfation pattern, particularly the 6-O-sulfate (6OS) of glucosamine [15,19,34,49,62]. HS 6OS levels are regulated during biosynthesis and post-synthetically by the extracellular sulfatases (SULFs) [49,93]. The SULFs were first identified in the quail embryo based on their ability to promote Wnt signaling in myogenic progenitor cells [31]. Subsequent studies have demonstrated SULF regulation of diverse signaling factors, including Wnts, FGF2, vascular endothelial growth factor (VEGF), glial cell line–derived neurotrophic (GDNF), and stromal cell–derived factor 1 (SDF1) [3,4,28,29,31,38,39,72,112,116]. Thus, it is likely that HS sulfation status in the NSPC niche helps to regulate crucial progenitor cell behaviors.

Indeed, alterations in HS sulfation can regulate NSPC differentiation. In the ventral spinal cord decreased HS sulfation, mediated by the SULFs, is necessary for the Shh-mediated switch from generation of neural progenitors to oligodendroglial progenitors [17,29,109]. In contrast, embryonic stem cells with undersulfation of HS exhibit restricted differentiation potential and blocked neural lineage maturation [36]. The striking difference in cell response to decreased HS sulfation in these two systems can be attributed to differences in HS binding specificity and function for different signaling molecules. In the spinal cord, Shh signaling is required for differentiation and, decreased HS sulfation promotes Shh signaling potentially via release of sequestered ligand [109]. In embryonic stem cells, differentiation into neural cells requires FGF signaling, and this signaling is promoted by sulfated HS [27,35,45,67,98,122]. The 6-O-sulfation levels of HS are also important in regulating signaling after neural injury. In a recent study, increased HS sulfation was shown to promote the formation of the glial scar

[52], a process that inhibits CNS repair, and involves concomitant up-regulation of CSPGs [16,73].

Consistent with HS chains playing an important role in regulating ligand-mediated signaling in the brain, genetic alteration of HS chain synthesis results in abnormal brain phenotypes in mice [34,82]. EXT1 (exostosin 1) and EXT2 (exostosin 2) encode glycosyltransferases which elongate HS chains and are essential for HS biosynthesis [78,79,85]. Conditional knockout of Ext1 in nestin-positive neural stem cells resulted in defects in neural patterning and cortical neurogenesis [56]. While multiple signaling pathways may be involved, the FGF pathway may be particularly important as Ext1-deficient neural stem cells exhibited reduced proliferation in response to FGF2 and FGF8 [56,121].

In addition to their ability to regulate ligand-mediated signaling, HSPGs also play fundamental roles in cell–cell and cell–matrix interactions, including binding to integrins as described in the previous section, which in turn have important roles in the NSPC niche [9–11,86].

3.2. Chondroitin sulfate proteoglycans

CSPGs are also abundantly expressed by NSPCs and have many known functions in ligand-mediated signaling, cell adhesion, and cell migration. Indeed, enzymatic removal of CS from NSPCs is associated with decreased proliferation in response to FGF2 and altered differentiation in response to EGF [106]. In embryonic OPCs, PTPRZ1, a transmembrane proteoglycan that can be modified by CS and DS, acts to maintain self-renewal and suppress differentiation [84]. Another proteoglycan, which has been widely studied to have many functions in OPCs, is the transmembrane proteoglycan CSPG4/NG2. Highly expressed on OPCs, CSPG4/NG2 promotes OPC proliferation and migration [40,46,69] (for review see [111]). Using mice lacking the *cspg4* gene, it has been shown that CSPG4/NG2 promotes proliferation of platelet-derived growth factor receptor alpha (PDGFR α)-positive OPCs and its absence confers delays in mature oligodendrocyte production [69]. In combination with binding studies demonstrating high affinity binding sites for FGF2 and PDGFAA on CSPG4/NG2 [46], it has been suggested that CSPG4/NG2 may act as a reservoir or co-receptor for these growth factors. In addition, direct interactions between CSPG4/NG2 and the receptor tyrosine kinase (RTK) itself can promote mitogenic signaling, as has been observed for FGFR1 and FGFR3 in pericytes and smooth muscle cells [23].

In OPCs, CSPG4/NG2 is also a marker of polarity and regulates EGF-dependent proliferation and self-renewal [107]. As CSPG4/NG2 is required to set up OPC polarity, CSPG4/NG2 may actively participate in regulating asymmetric progenitor divisions, a fundamental process to maintaining progenitor populations in the brain [107]. In addition, CSPG4/NG2 can functionally interact with diverse ECM components [20], including $\alpha 3 \beta 1$ integrins [40] and carbohydrate binding proteins (lectins) [40,119], which, as already described, are important components of the NPSC niche. As with HSPGs, sulfation of CS chains is a critical determinant of function, and knockdown of CSPG biosynthetic enzymes have demonstrated defects in cell migration from the ventricular zone into the cortical plate [58].

Another major constituent of the developing brain and of the adult NSPC niche is hyaluronan (or hyaluronic acid, HA). The physiologic role for HA is diverse due to both its range in size, from a small number of disaccharide units to an extensive high molecular weight polysaccharide, and its ability to interact with multiple extracellular molecules, including hyalactins, neurocan, aggrecan, versican, and lectican present in the NSPC niche [1,60]. Hyaluronan can also block the differentiation of progenitor cells in the brain [37] and promote activation of RTK signaling pathways including ERBB2 and PDGFR β [90]. Its function in the NSPC niche has been recently reviewed [97].

4. Conclusions

The development and repair of the central nervous system require both precise and dynamic regulation of NSPCs that must persist over the life of the organism. The extracellular microenvironment, or niche, is complex. There are many cellular components, such as endothelial cells, ependymal cells, more differentiated neural cells, astrocytes, and microglia. There are soluble ligands, including growth factors, morphogens, chemokines, and neurotransmitters, and there are the many components of the extracellular matrix. Together these components make up the NSPC niche and act to regulate fundamental behaviors of NSPCs. In this review, we use proteoglycans to illustrate ways in which the niche can regulate NSPC behavior. An improved understanding of how extrinsic cues regulate NSPC behavior is critical and may contribute to advances in stem/progenitor cell-based therapies, improved repair from CNS injury, and potentially novel therapies for neoplastic diseases in the brain including glioblastoma.

Acknowledgments

This work was supported by the National Institutes of Health (R01 NS081117 to JJP) and the James S. McDonnell Foundation (J.J.P.). We apologize to the many investigators whose articles we did not cite due to space constraints.

References

- [1] M. Abaskharoun, M. Bellemare, E. Lau, R.U. Margolis, Expression of hyaluronan and the hyaluronan-binding proteoglycans neurocan, aggrecan, and versican by neural stem cells and neural cells derived from embryonic stem cells, *Brain Res.* 1327 (2010) 6–15.
- [2] A. Aguirre, M.E. Rubio, V. Gallo, Notch and EGFR pathway interaction regulates neural stem cell number and self-renewal, *Nature* 467 (2010) 323–327.
- [3] X. Ai, A.T. Do, O. Lozynska, et al., Qsulf1 remodels the 6-O sulfation states of cell surface heparan sulfate proteoglycans to promote Wnt signaling, *J. Cell Biol.* 162 (2003) 341–351.
- [4] X. Ai, T. Kitazawa, A.T. Do, et al., SULF1 and SULF2 regulate heparan sulfate-mediated GDNF signaling for esophageal innervation, *Development* 134 (2007) 3327–3338.
- [5] A. Alvarez-Buylla, D.A. Lim, For the long run: maintaining germinal niches in the adult brain, *Neuron* 41 (2004) 683–686.
- [6] E. Arikawa-Hirasawa, H. Watanabe, H. Takami, J.R. Hassell, Y. Yamada, Perlecan is essential for cartilage and cephalic development, *Nat. Genet.* 23 (1999) 354–358.
- [7] E. Arikawa-Hirasawa, W.R. Wilcox, A.H. Le, et al., Dyssegmental dysplasia, Silverman-Handmaker type, is caused by functional null mutations of the perlecan gene, *Nat. Genet.* 27 (2001) 431–434.
- [8] D. Aviezer, E. Levy, M. Safran, et al., Differential structural requirements of heparin and heparan sulfate proteoglycans that promote binding of basic fibroblast growth factor to its receptor, *J. Biol. Chem.* 269 (1994) 114–121.
- [9] M.D. Bass, K.A. Roach, M.R. Morgan, et al., Syndecan-4-dependent Rac1 regulation determines directional migration in response to the extracellular matrix, *J. Cell Biol.* 177 (2007) 527–538.
- [10] D.M. Beauvais, B.J. Burbach, A.C. Rapraeger, The syndecan-1 ectodomain regulates alphavbeta3 integrin activity in human mammary carcinoma cells, *J. Cell Biol.* 167 (2004) 171–181.
- [11] D.M. Beauvais, B.J. Ell, A.R. McWhorter, A.C. Rapraeger, Syndecan-1 regulates alphavbeta3 and alphavbeta5 integrin activation during angiogenesis and is blocked by synstatin, a novel peptide inhibitor, *J. Exp. Med.* 206 (2009) 691–705.
- [12] D.M. Beauvais, A.C. Rapraeger, Syndecan-1 couples the insulin-like growth factor-1 receptor to inside-out integrin activation, *J. Cell Sci.* 123 (2010) 3796–3807.
- [13] M. Bernfield, M. Gotte, P.W. Park, et al., Functions of cell surface heparan sulfate proteoglycans, *Annu. Rev. Biochem.* 68 (1999) 729–777.
- [14] M.M. Beshpalov, Y.A. Sidorova, S. Tumova, et al., Heparan sulfate proteoglycan syndecan-3 is a novel receptor for GDNF, neurturin, and artemin, *J. Cell Biol.* 192 (2011) 153–169.
- [15] R.J. Bink, H. Habuchi, Z. Lele, et al., Heparan sulfate 6-O-sulfotransferase is essential for muscle development in zebrafish, *J. Biol. Chem.* 278 (2003) 31118–31127.
- [16] E.J. Bradbury, L.D. Moon, R.J. Popat, et al., Chondroitinase ABC promotes functional recovery after spinal cord injury, *Nature* 416 (2002) 636–640.
- [17] C. Braquart-Varnier, C. Danesin, C. Cloucard-Martinato, et al., A subtractive approach to characterize genes with regionalized expression in the gliogenic ventral neuroepithelium: identification of chick sulfatase 1 as a new oligodendrocyte lineage gene, *Mol. Cell. Neurosci.* 25 (2004) 612–628.
- [18] J.J. Breunig, M.R. Sarkisian, J.I. Arellano, et al., Primary cilia regulate hippocampal neurogenesis by mediating sonic hedgehog signaling, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 13127–13132.
- [19] H.E. Bulow, O. Hobert, Differential sulfations and epimerization define heparan sulfate specificity in nervous system development, *Neuron* 41 (2004) 723–736.

- [20] M.A. Burg, E. Tillet, R. Timpl, W.B. Stallcup, Binding of the NG2 proteoglycan to type VI collagen and other extracellular matrix molecules, *J. Biol. Chem.* 271 (1996) 26110–26116.
- [21] R.C. Burrows, D. Wancio, P. Levitt, L. Lillien, Response diversity and the timing of progenitor cell maturation are regulated by developmental changes in EGFR expression in the cortex, *Neuron* 19 (1997) 251–267.
- [22] L. Cao, X. Jiao, D.S. Zuzga, et al., VEGF links hippocampal activity with neurogenesis, learning and memory, *Nat. Genet.* 36 (2004) 827–835.
- [23] S. Cattaruzza, U. Ozerdem, M. Denzel, et al., Multivalent proteoglycan modulation of FGF mitogenic responses in perivascular cells, *Angiogenesis* 16 (2013) 309–327.
- [24] C.B. Chambers, Y. Peng, H. Nguyen, et al., Spatiotemporal selectivity of response to Notch1 signals in mammalian forebrain precursors, *Development* 128 (2001) 689–702.
- [25] P. Chapouton, P. Skupien, B. Hesl, et al., Notch activity levels control the balance between quiescence and recruitment of adult neural stem cells, *J. Neurosci.* 30 (2010) 7961–7974.
- [26] A. Chenn, C.A. Walsh, Regulation of cerebral cortical size by control of cell cycle exit in neural precursors, *Science* 297 (2002) 365–369.
- [27] C.Y. Chuang, M.S. Lord, J. Melrose, et al., Heparan sulfate-dependent signaling of fibroblast growth factor 18 by chondrocyte-derived perlecan, *Biochemistry* 49 (2010) 5524–5532.
- [28] Y. Dai, Y. Yang, V. MacLeod, et al., HSulf-1 and HSulf-2 are potent inhibitors of myeloma tumor growth in vivo, *J. Biol. Chem.* 280 (2005) 40066–40073.
- [29] C. Danesin, E. Agius, N. Escalas, et al., Ventral neural progenitors switch toward an oligodendroglial fate in response to increased Sonic hedgehog (Shh) activity: involvement of Sulfatase 1 in modulating Shh signaling in the ventral spinal cord, *J. Neurosci.* 26 (2006) 5037–5048.
- [30] L. Dang, K. Yoon, M. Wang, N. Gaiano, Notch3 signaling promotes radial glial/progenitor character in the mammalian telencephalon, *Dev. Neurosci.* 28 (2006) 58–69.
- [31] G.K. Dhoot, M.K. Gustafsson, X. Ai, et al., Regulation of Wnt signaling and embryo patterning by an extracellular sulfatase, *Science* 293 (2001) 1663–1666.
- [32] F. Doetsch, I. Caille, D.A. Lim, J.M. Garcia-Verdugo, A. Alvarez-Buylla, Subventricular zone astrocytes are neural stem cells in the adult mammalian brain, *Cell* 97 (1999) 703–716.
- [33] F. Echtermeyer, P.C. Baci, S. Saoncella, Y. Ge, P.F. Goetinck, Syndecan-4 core protein is sufficient for the assembly of focal adhesions and actin stress fibers, *J. Cell Sci.* 112 (Pt 20) (1999) 3433–3441.
- [34] J.D. Esko, U. Lindahl, Molecular diversity of heparan sulfate, *J. Clin. Invest.* 108 (2001) 169–173.
- [35] M. Ford-Perriss, S.E. Guimond, U. Greferath, et al., Variant heparan sulfates synthesized in developing mouse brain differentially regulate FGF signaling, *Glycobiology* 12 (2002) 721–727.
- [36] M. Forsberg, K. Holmborn, S. Kundu, et al., Undersulfation of heparan sulfate restricts differentiation potential of mouse embryonic stem cells, *J. Biol. Chem.* 287 (2012) 10853–10862.
- [37] J.R. Fraser, T.C. Laurent, U.B. Laurent, Hyaluronan: its nature, distribution, functions and turnover, *J. Intern. Med.* 242 (1997) 27–33.
- [38] S.D. Freeman, W.M. Moore, E.C. Guiral, et al., Extracellular regulation of developmental cell signaling by Xtsulf1, *Dev. Biol.* 320 (2008) 436–445.
- [39] K. Fujita, E. Takechi, N. Sakamoto, et al., HpSulf, a heparan sulfate 6-O-endosulfatase, is involved in the regulation of VEGF signaling during sea urchin development, *Mech. Dev.* 127 (2010) 235–245.
- [40] J. Fukushi, I.T. Makagiansar, W.B. Stallcup, NG2 proteoglycan promotes endothelial cell motility and angiogenesis via engagement of galectin-3 and alpha3beta1 integrin, *Mol. Biol. Cell* 15 (2004) 3580–3590.
- [41] C.R. Gajera, H. Emich, O. Lioubinski, et al., LRP2 in ependymal cells regulates BMP signaling in the adult neurogenic niche, *J. Cell Sci.* 123 (2010) 1922–1930.
- [42] E. Garcion, A. Faissner, C. French-Constant, Knockout mice reveal a contribution of the extracellular matrix molecule tenascin-C to neural precursor proliferation and migration, *Development* 128 (2001) 2485–2496.
- [43] E. Garcion, A. Halilagic, A. Faissner, C. French-Constant, Generation of an environmental niche for neural stem cell development by the extracellular matrix molecule tenascin C, *Development* 131 (2004) 3423–3432.
- [44] A. Giros, J. Morante, C. Gil-Sanz, A. Fairen, M. Costell, Perlecan controls neurogenesis in the developing telencephalon, *BMC Dev. Biol.* 7 (2007) 29.
- [45] S.J. Goodger, C.J. Robinson, K.J. Murphy, et al., Evidence that heparin saccharides promote FGF2 mitogenesis through two distinct mechanisms, *J. Biol. Chem.* 283 (2008) 13001–13008.
- [46] L. Goretzki, M.A. Burg, K.A. Grako, W.B. Stallcup, High-affinity binding of basic fibroblast growth factor and platelet-derived growth factor-AA to the core protein of the NG2 proteoglycan, *J. Biol. Chem.* 274 (1999) 16831–16837.
- [47] M. Gotz, W.B. Huttner, The cell biology of neurogenesis, *Nat. Rev. Mol. Cell Biol.* 6 (2005) 777–788.
- [48] D.K. Greene, S. Tumova, J.R. Couchman, A. Woods, Syndecan-4 associates with alpha-actinin, *J. Biol. Chem.* 278 (2003) 7617–7623.
- [49] H. Habuchi, O. Habuchi, K. Kimata, Sulfation pattern in glycosaminoglycan: does it have a code? *Glycoconj. J.* 21 (2004) 47–52.
- [50] A. Hall, N.A. Giese, W.D. Richardson, Spinal cord oligodendrocytes develop from ventrally derived progenitor cells that express PDGF alpha-receptors, *Development* 122 (1996) 4085–4094.
- [51] Y.G. Han, N. Spassky, M. Romaguera-Ros, et al., Hedgehog signaling and primary cilia are required for the formation of adult neural stem cells, *Nat. Neurosci.* 11 (2008) 277–284.
- [52] J.R. Higginson, S.M. Thompson, A. Santos-Silva, et al., Differential sulfation remodeling of heparan sulfate by extracellular 6-O-sulfatases regulates fibroblast growth factor-induced boundary formation by glial cells: implications for glial cell transplantation, *J. Neurosci.* 32 (2012) 15902–15912.
- [53] M. Iida, T. Shuo, K. Hirano, et al., Identification and functions of chondroitin sulfate in the milieu of neural stem cells, *J. Biol. Chem.* 281 (2006) 5982–5991.
- [54] R.A. Ihrie, J.K. Shah, C.C. Harwell, et al., Persistent sonic hedgehog signaling in adult brain determines neural stem cell positional identity, *Neuron* 71 (2011) 250–262.
- [55] I. Imayoshi, M. Sakamoto, M. Yamaguchi, K. Mori, R. Kageyama, Essential roles of Notch signaling in maintenance of neural stem cells in developing and adult brains, *J. Neurosci.* 30 (2010) 3489–3498.
- [56] M. Inatani, F. Irie, A.S. Plump, M. Tessier-Lavigne, Y. Yamaguchi, Mammalian brain morphogenesis and midline axon guidance require heparan sulfate, *Science* 302 (2003) 1044–1046.
- [57] S. Ishibashi, T. Kuroiwa, M. Sakaguchi, et al., Galectin-1 regulates neurogenesis in the subventricular zone and promotes functional recovery after stroke, *Exp. Neurol.* 207 (2007) 302–313.
- [58] M. Ishii, N. Maeda, Oversulfated chondroitin sulfate plays critical roles in the neuronal migration in the cerebral cortex, *J. Biol. Chem.* 283 (2008) 32610–32620.
- [59] K. Jin, Y. Zhu, Y. Sun, et al., Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 11946–11950.
- [60] P. Kabos, H. Matundan, M. Zandian, et al., Neural precursors express multiple chondroitin sulfate proteoglycans, including the lectican family, *Biochem. Biophys. Res. Commun.* 318 (2004) 955–963.
- [61] M.Y. Kalani, S.H. Cheshier, B.J. Cord, et al., Wnt-mediated self-renewal of neural stem/progenitor cells, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 16970–16975.
- [62] K. Kamimura, M. Fujise, F. Villa, et al., Drosophila heparan sulfate 6-O-sulfotransferase (dHS6ST) gene. Structure, expression, and function in the formation of the tracheal system, *J. Biol. Chem.* 276 (2001) 17014–17021.
- [63] M. Karus, B. Denecke, C. French-Constant, S. Wiese, A. Faissner, The extracellular matrix molecule tenascin C modulates expression levels and territories of key patterning genes during spinal cord astrocyte specification, *Development* 138 (2011) 5321–5331.
- [64] E. Kokovay, S. Goderie, Y. Wang, et al., Adult SVZ lineage cells home to and leave the vascular niche via differential responses to SDF1/CXCR4 signaling, *Cell Stem Cell* 7 (2010) 163–173.
- [65] E. Kokovay, Y. Wang, G. Kusek, et al., VCAM1 is essential to maintain the structure of the SVZ niche and acts as an environmental sensor to regulate SVZ lineage progression, *Cell Stem Cell* 11 (2012) 220–230.
- [66] D.C. Kraushaar, S. Rai, E. Condac, et al., Heparan sulfate facilitates FGF and BMP signaling to drive mesoderm differentiation of mouse embryonic stem cells, *J. Biol. Chem.* 287 (2012) 22691–22700.
- [67] J. Kreuger, M. Salmivirta, L. Sturiale, G. Gimenez-Gallego, U. Lindahl, Sequence analysis of heparan sulfate epitopes with graded affinities for fibroblast growth factors 1 and 2, *J. Biol. Chem.* 276 (2001) 30744–30752.
- [68] A. Kriegstein, A. Alvarez-Buylla, The glial nature of embryonic and adult neural stem cells, *Annu. Rev. Neurosci.* 32 (2009) 149–184.
- [69] K. Kucharova, W.B. Stallcup, The NG2 proteoglycan promotes oligodendrocyte progenitor proliferation and developmental myelination, *Neuroscience* 166 (2010) 185–194.
- [70] J.C. Kwok, P. Warren, J.W. Fawcett, Chondroitin sulfate: a key molecule in the brain matrix, *Int. J. Biochem. Cell Biol.* 44 (2012) 582–586.
- [71] J. Lai, J. Chien, J. Staub, et al., Loss of HSulf-1 up-regulates heparin-binding growth factor signaling in cancer, *J. Biol. Chem.* 278 (2003) 23107–23117.
- [72] W.C. Lamanna, M.A. Frese, M. Balleininger, T. Dierks, Sulf loss influences N-, 2-O-, and 6-O-sulfation of multiple heparan sulfate proteoglycans and modulates fibroblast growth factor signaling, *J. Biol. Chem.* 283 (2008) 27724–27735.
- [73] L.W. Lau, M.B. Keough, S. Haylock-Jacobs, et al., Chondroitin sulfate proteoglycans in demyelinated lesions impair remyelination, *Ann. Neurol.* 72 (2012) 419–432.
- [74] T. Licht, I. Goshen, A. Avital, et al., Reversible modulations of neuronal plasticity by VEGF, *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 5081–5086.
- [75] D.C. Lie, S.A. Colamarino, H.J. Song, et al., Wnt signalling regulates adult hippocampal neurogenesis, *Nature* 437 (2005) 1370–1375.
- [76] L. Lillien, H. Raphael, BMP and FGF regulate the development of EGF-responsive neural progenitor cells, *Development* 127 (2000) 4993–5005.
- [77] D.A. Lim, A.D. Tramontin, J.M. Trevejo, et al., Noggin antagonizes BMP signaling to create a niche for adult neurogenesis, *Neuron* 28 (2000) 713–726.
- [78] X. Lin, G. Wei, Z. Shi, et al., Disruption of gastrulation and heparan sulfate biosynthesis in EXT1-deficient mice, *Dev. Biol.* 224 (2000) 299–311.
- [79] T. Lind, F. Tufaro, C. McCormick, U. Lindahl, K. Lidholt, The putative tumor suppressors EXT1 and EXT2 are glycosyltransferases required for the biosynthesis of heparan sulfate, *J. Biol. Chem.* 273 (1998) 26265–26268.
- [80] X. Liu, Q. Wang, T.F. Haydar, A. Bordey, Nonsynaptic GABA signaling in postnatal subventricular zone controls proliferation of GFAP-expressing progenitors, *Nat. Neurosci.* 8 (2005) 1179–1187.
- [81] R.L. Longley, A. Woods, A. Fleetwood, et al., Control of morphology, cytoskeleton and migration by syndecan-4, *J. Cell Sci.* 112 (Pt 20) (1999) 3421–3431.
- [82] N. Maeda, M. Ishii, K. Nishimura, K. Kamimura, Functions of chondroitin sulfate and heparan sulfate in the developing brain, *Neurochem. Res.* 36 (2011) 1228–1240.
- [83] H.K. Matthews, L. Marchant, C. Carmona-Fontaine, et al., Directional migration of neural crest cells in vivo is regulated by Syndecan-4/Rac1 and non-canonical Wnt signaling/RhoA, *Development* 135 (2008) 1771–1780.
- [84] C.R. McClain, F.J. Sim, S.A. Goldman, Pleiotrophin suppression of receptor protein tyrosine phosphatase-beta/zeta maintains the self-renewal competence of fetal human oligodendrocyte progenitor cells, *J. Neurosci.* 32 (2012) 15066–15075.

- [85] C. McCormick, Y. Leduc, D. Martindale, et al., The putative tumour suppressor EXT1 alters the expression of cell-surface heparan sulfate, *Nat. Genet.* 19 (1998) 158–161.
- [86] K.J. McQuade, D.M. Beauvais, B.J. Burbach, A.C. Rapraeger, Syndecan-1 regulates alphavbeta5 integrin activity in B82L fibroblasts, *J. Cell Sci.* 119 (2006) 2445–2456.
- [87] F. Mercier, E. Arikawa-Hirasawa, Heparan sulfate niche for cell proliferation in the adult brain, *Neurosci. Lett.* 510 (2012) 67–72.
- [88] F. Mercier, J.T. Kitasako, G.I. Hattton, Anatomy of the brain neurogenic zones revisited: fractones and the fibroblast/macrophage network, *J. Comp. Neurol.* 451 (2002) 170–188.
- [89] H. Mira, Z. Andreu, H. Suh, et al., Signaling through BMPR-IA regulates quiescence and long-term activity of neural stem cells in the adult hippocampus, *Cell Stem Cell* 7 (2010) 78–89.
- [90] S. Misra, B.P. Toole, S. Ghatak, Hyaluronan constitutively regulates activation of multiple receptor tyrosine kinases in epithelial and carcinoma cells, *J. Biol. Chem.* 281 (2006) 34936–34941.
- [91] A.V. Molofsky, R. Krencik, E.M. Ullian, et al., Astrocytes and disease: a neurodevelopmental perspective, *Genes Dev.* 26 (2012) 891–907.
- [92] M. Mongiat, S.M. Sweeney, J.D. San Antonio, J. Fu, R.V. Iozzo, Endorepellin, a novel inhibitor of angiogenesis derived from the C terminus of perlecan, *J. Biol. Chem.* 278 (2003) 4238–4249.
- [93] M. Morimoto-Tomita, K. Uchimura, Z. Werb, S. Hemmerich, S.D. Rosen, Cloning and characterization of two extracellular heparin-degrading endosulfatases in mice and humans, *J. Biol. Chem.* 277 (2002) 49175–49185.
- [94] S.C. Noctor, A.C. Flint, T.A. Weissman, R.S. Dammerman, A.R. Kriegstein, Neurons derived from radial glial cells establish radial units in neocortex, *Nature* 409 (2001) 714–720.
- [95] Y. Park, C. Rangel, M.M. Reynolds, et al., Drosophila perlecan modulates FGF and hedgehog signals to activate neural stem cell division, *Dev. Biol.* 253 (2003) 247–257.
- [96] A. Polito, R. Reynolds, NG2-expressing cells as oligodendrocyte progenitors in the normal and demyelinated adult central nervous system, *J. Anat.* 207 (2005) 707–716.
- [97] M. Preston, L.S. Sherman, Neural stem cell niches: roles for the hyaluronan-based extracellular matrix, *Front. Biosci.* 3 (2011) 1165–1179.
- [98] A.C. Rapraeger, A. Krufka, B.B. Olwin, Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation, *Science* 252 (1991) 1705–1708.
- [99] J. Relucio, M.J. Menezes, Y. Miyagoe-Suzuki, S. Takeda, H. Colognato, Laminin regulates postnatal oligodendrocyte production by promoting oligodendrocyte progenitor survival in the subventricular zone, *Glia* 60 (2012) 1451–1467.
- [100] M. Sakaguchi, Y. Imaizumi, T. Shingo, et al., Regulation of adult neural progenitor cells by Galectin-1/beta1 Integrin interaction, *J. Neurochem.* 113 (2010) 1516–1524.
- [101] M. Sakaguchi, T. Shingo, T. Shimazaki, et al., A carbohydrate-binding protein, Galectin-1, promotes proliferation of adult neural stem cells, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 7112–7117.
- [102] S. Sarrazin, W.C. Lamanna, J.D. Esko, Heparan sulfate proteoglycans, *Cold Spring Harb. Perspect. Biol.* 3 (2011).
- [103] B. Seri, J.M. Garcia-Verdugo, L. Collado-Morente, B.S. McEwen, A. Alvarez-Buylla, Cell types, lineage, and architecture of the germinal zone in the adult dentate gyrus, *J. Comp. Neurol.* 478 (2004) 359–378.
- [104] B. Seri, J.M. Garcia-Verdugo, B.S. McEwen, A. Alvarez-Buylla, Astrocytes give rise to new neurons in the adult mammalian hippocampus, *J. Neurosci.* 21 (2001) 7153–7160.
- [105] Q. Shen, Y. Wang, E. Kokovay, et al., Adult SVZ stem cells lie in a vascular niche: a quantitative analysis of niche cell-cell interactions, *Cell Stem Cell* 3 (2008) 289–300.
- [106] S. Sirko, A. von Holst, A. Weber, et al., Chondroitin sulfates are required for fibroblast growth factor-2-dependent proliferation and maintenance in neural stem cells and for epidermal growth factor-dependent migration of their progeny, *Stem Cells* 28 (2010) 775–787.
- [107] S. Sugiarto, A.I. Persson, E.G. Munoz, et al., Asymmetry-defective oligodendrocyte progenitors are glioma precursors, *Cancer Cell* 20 (2011) 328–340.
- [108] J. Takagi, Structural basis for ligand recognition by integrins, *Curr. Opin. Cell Biol.* 19 (2007) 557–564.
- [109] Y. Touahri, N. Escalas, B. Benazeraf, et al., Sulfatase 1 promotes the motor neuron-to-oligodendrocyte fate switch by activating Shh signaling in Olig2 progenitors of the embryonic ventral spinal cord, *J. Neurosci.* 32 (2012) 18018–18034.
- [110] Y. Tozuka, S. Fukuda, T. Namba, T. Seki, T. Hisatsune, GABAergic excitation promotes neuronal differentiation in adult hippocampal progenitor cells, *Neuron* 47 (2005) 803–815.
- [111] J. Trotter, K. Karram, A. Nishiyama, NG2 cells: properties, progeny and origin, *Brain Res. Rev.* 63 (2010) 72–82.
- [112] K. Uchimura, M. Morimoto-Tomita, A. Bistrup, et al., HSulf-2, an extracellular endoglucosaminase-6-sulfatase, selectively mobilizes heparin-bound growth factors and chemokines: effects on VEGF, FGF-1, and SDF-1, *BMC Biochem.* 7 (2006) 2.
- [113] F.M. Vaccarino, M.L. Schwartz, R. Raballo, et al., Changes in cerebral cortex size are governed by fibroblast growth factor during embryogenesis, *Nat. Neurosci.* 2 (1999) 848.
- [114] A. Varki, M.E. Etzler, R.D. Cummings, J.D. Esko, Discovery and Classification of Glycan-Binding Proteins, in: A. Varki, R.D. Cummings, J.D. Esko, H.H. Freeze, P. Stanley, C.R. Bertozzi, G.W. Hart, M.E. Etzler (Eds.), *Cold Spring Harbor, NY*, 2009, (2009, pp.).
- [115] M. Vigny, M.P. Ollier-Hartmann, M. Lavigne, et al., Specific binding of basic fibroblast growth factor to basement membrane-like structures and to purified heparan sulfate proteoglycan of the EHS tumor, *J. Cell. Physiol.* 137 (1988) 321–328.
- [116] B.L. Viviano, S. Paine-Saunders, N. Gasunas, J. Gallagher, S. Saunders, Domain-specific modification of heparan sulfate by Qsulf1 modulates the binding of the bone morphogenetic protein antagonist Noggin, *J. Biol. Chem.* 279 (2004) 5604–5611.
- [117] L.P. Wang, G. Kempermann, H. Kettenmann, A subpopulation of precursor cells in the mouse dentate gyrus receives synaptic GABAergic input, *Mol. Cell. Neurosci.* 29 (2005) 181–189.
- [118] Q. Wang, L. Yang, C. Alexander, S. Temple, The niche factor syndecan-1 regulates the maintenance and proliferation of neural progenitor cells during mammalian cortical development, *PLoS One* 7 (2012) e42883.
- [119] Y. Wen, I.T. Makagiansar, J. Fukushi, et al., Molecular basis of interaction between NG2 proteoglycan and galectin-3, *J. Cell. Biochem.* 98 (2006) 115–127.
- [120] J.M. Whitelock, J. Melrose, R.V. Iozzo, Diverse cell signaling events modulated by perlecan, *Biochemistry* 47 (2008) 11174–11183.
- [121] Y. Yamaguchi, M. Inatani, Y. Matsumoto, J. Ogawa, F. Irie, Roles of heparan sulfate in mammalian brain development current views based on the findings from Ext1 conditional knockout studies, *Prog. Mol. Biol. Transl. Sci.* 93 (2010) 133–152.
- [122] A. Yayon, M. Klagsbrun, J.D. Esko, P. Leder, D.M. Ornitz, Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor, *Cell* 64 (1991) 841–848.
- [123] K.M. Young, K. Psachoulia, R.B. Tripathi, et al., Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling, *Neuron* 77 (2013) 873–885.
- [124] K. Yu, S. McGlynn, M.P. Matisse, Floor plate-derived sonic hedgehog regulates glial and ependymal cell fates in the developing spinal cord, *Development* 140 (2013) 1594–1604.
- [125] C. Zhao, W. Deng, F.H. Gage, Mechanisms and functional implications of adult neurogenesis, *Cell* 132 (2008) 645–660.