



Review

Mucins and tumor resistance to chemotherapeutic drugs


Nicolas Jonckheere^{*,1}, Nicolas Skrypek¹, Isabelle Van Seuning

Inserm, UMR837, Jean Pierre Aubert Research Center, Team #5 "Mucins, Epithelial Differentiation and Carcinogenesis", rue Polonovski, 59045 Lille Cedex, France
 Université Lille Nord de France, Lille, France
 Centre Hospitalier Régional et Universitaire de Lille, Place de Verdun, 59037 Lille Cedex, France

ARTICLE INFO

Article history:

Received 3 March 2014

Received in revised form 22 April 2014

Accepted 23 April 2014

Available online 29 April 2014

Keywords:

Mucin
 Cancer
 Resistance
 Chemotherapeutic drug
 Apoptosis
 Prognosis biomarker

ABSTRACT

Epithelial cancer patients not considered eligible for surgical resection frequently benefit from chemotherapy. Chemotherapy is the treatment of cancer with one or combination of cytotoxic or cytostatic drugs. Recent advances in chemotherapy allowed a great number of cancer patients to receive treatment with significant results. Unfortunately, resistance to chemotherapeutic drug treatment is a major challenge for clinicians in the majority of epithelial cancers because it is responsible for the inefficiency of therapies.

Mucins belong to a heterogeneous group of large O-glycoproteins that can be either secreted or membrane-bound. Implications of mucins have been described in relation to cancer cell behavior and cell signaling pathways associated with epithelial tumorigenesis. Because of the frequent alteration of the pattern of mucin expression in cancers as well as their structural and functional characteristics, mucins are thought to also be involved in response to therapies. In this report, we review the roles of mucins in chemoresistance and the associated underlying molecular mechanisms (physical barrier, resistance to apoptosis, drug metabolism, cell stemness, epithelial–mesenchymal transition) and discuss the therapeutic tools/strategies and/or prognosis biomarkers for personalized chemotherapy that could be proposed from these studies.

© 2014 Elsevier B.V. All rights reserved.

Contents

1. Introduction	143
2. Mucins and chemoresistance <i>in vitro</i>	143
3. Mucins and clinical response to chemotherapy	144
3.1. Mucinous tumors	144
3.2. Non-mucinous tumors	144
4. Mucins form a physical barrier	144
5. Mucins and resistance to apoptosis	144
5.1. MUC1	144
5.2. MUC4	146
6. Mucins and alteration of drug metabolism	146
7. Mucins and cancer stem cells	147
8. Mucins and epithelial–mesenchymal transition	147
9. Mucins and polymorphisms associated with chemosensitivity?	148
10. Outlook to the future: Using mucins as a therapeutic target to sensitize cancer cells to chemotherapeutic drugs or as biomarkers of chemoresistance?	148
11. Conclusion	148
Acknowledgements	149
References	149

Abbreviations: PDAC, Pancreatic ductal adenocarcinoma; CRC, Colorectal cancer; EMT, Epithelial–mesenchymal transition; MUC1-CT, MUC1 cytoplasmic tail; N-t, Amino-terminal; C-t, Carboxy-terminal; PTS, Proline Threonine Serine; 5-FU, 5-fluoro-uracile; ROS, Reactive oxygen species; VNTR, Variable number tandem repeat; SNP, single nucleotide polymorphisms

* Corresponding author at: Inserm UMR837/JPARC, rue Polonovski, 59045 Lille Cedex, France. Tel.: +33 3 20 29 88 65; fax: +33 3 20 53 85 62.

E-mail address: nicolas.jonckheere@inserm.fr (N. Jonckheere).

¹ Authors have equally contributed to this manuscript.

1. Introduction

Mucins belong to a heterogeneous group of large O-glycoproteins composed of a long peptidic chain (called apomucin) on which are linked hundreds of oligosaccharidic chains. Initially, the mucin word designated glycoproteins secreted by specialized epithelial cells, the goblet cells, as part of the mucus gel. Mucins were biochemically characterized as massive molecules with high molecular weight able to form viscoelastic gels and responsible for the rheological properties of mucus [1]. The molecular era that led to genome sequencing allowed the classification of two sub-groups of mucins: (i) secreted mucins that mostly complied with this definition and (ii) membrane-bound or transmembrane mucins that did not fit in. Despite this dichotomy, mucins were all included in the MUC family with the approval of Human Genome Organization Gene Nomenclature Committee (HUGO/GNC) [2].

Secreted mucins are the major components of viscoelastic mucus gels and form a tridimensional network that protects the epithelia against various aggressions (inflammation, bacteria, virus, pollutants, pH, etc.). This subgroup mainly includes: MUC2, MUC5AC, MUC5B, MUC6 (clustered on the p15 arm of chromosome 11) and MUC19. MUC7 and MUC9 are smaller secreted mucins that do not oligomerize and are secreted by specialized cells as monomers [3–5]. Secreted mucins comprise amino- (N-t) and carboxy-terminal (C-t) regions sharing structural domains with von Willebrand (vW) factor. The central part is enriched in Pro, Thr and Ser amino acid residues forming the variable PTS domain that is O-glycosylated [4]. The O-glycosylation process is crucial for mucin secretion, stability, processing, and functions during both development and pathophysiological conditions [6–8]. The adjacent CYS domains are highly hydrophobic and are believed to cause the aggregation of mucins [9]. By forming disulfide bonds, the main intestinal mucin MUC2 dimerizes via its cysteine-knot (CK) domain and also trimerizes via N-t vWD domains building a complex molecular network [10,11]. On the contrary MUC5AC and MUC5B are linear disulfide-linked polymers that polydisperse and behave as random coils in solution [4].

The membrane-bound mucins are type I membrane-anchored proteins including MUC1, MUC3, MUC4, MUC12, MUC13, MUC15, MUC16, MUC17, MUC20, MUC21 and MUC22 [12–14]. Typically, membrane-bound mucins contain a long extracellular domain, a hydrophobic transmembrane domain, and a short cytoplasmic tail. Analysis of the peptidic sequences of mucins allowed description of their modular organization. The PTS domain, the only domain not conserved at the genomic level, is the common feature between mucins. Membrane-bound mucins share conserved domains such as epidermal growth factor-like (EGF) or Sea urchin sperm protein Enterokinase and Agrin (SEA) domains [13,15,16]. Based on their structure and localization at the cell surface they were shown to act in cell–cell, cell–extracellular matrix interactions and in cell signaling.

Mucins have a cell- and tissue-specific pattern of expression profoundly altered in epithelial cancers (loss of expression, overexpression, aberrant expression, neo-expression, glycosylation alterations) [17–21]. Because of their specific pattern of expression during the different steps of tumor progression toward adenocarcinoma, mucins stay under intense investigation as both potent new biomarkers and therapeutic targets in epithelial cancers.

Numerous reviews in the literature describe the roles of mucins in relation to cancer cell behavior and cell signaling pathways associated with tumorigenesis. Among them, membrane-bound mucins MUC1 and MUC4 have been extensively studied [14,22–25]. MUC1 and MUC4 govern both cellular differentiation and proliferation. They are also involved in metastasis and tumor proliferation. Secreted mucins MUC5B and MUC5AC and membrane-bound mucins MUC13 and MUC16 have also been associated with aggressive behavior of cancer cells [26–31]. On the contrary, Muc2 is involved in the suppression of colorectal cancer (CRC) since Muc2^{KO} mice develop adenoma progressing with age to invasive adenocarcinoma [32].

The dramatic outcome of epithelial cancers of the gastrointestinal tract is often related to a lack of both efficient therapeutic tools and early diagnostic markers. Patients not considered eligible for surgical resection frequently benefit from chemotherapy. However, chemoresistance is a common feature of epithelial cancers. Lately, mucins have been proposed as actors of this phenomenon. In this review, we discuss their role in chemoresistance and the associated cellular mechanisms to propose them as therapeutic tools and/or prognosis biomarkers of chemotherapy response.

2. Mucins and chemoresistance *in vitro*

In our laboratory and others, initial *in vitro* studies showing the relationship between mucins and chemoresistance came from colorectal carcinoma cells (HT29) stably resistant to 5-fluorouracil (5-FU) or methotrexate [33–35]. These cells were characterized by the overexpression of secreted mucins when they became resistant cells. This observation pointed out to the potential of mucins as actors of chemoresistance (Table 1).

In breast cancer cells, the overexpression of MUC1 is involved in cell sensitivity to Herceptin® via the increase of the cleavage of this mucin. These cells are also resistant to paclitaxel (Taxol®), doxorubicin and cyclophosphamide, suggesting a broader involvement of membrane-bound mucins [36]. Similarly, silencing the MUC1 or MUC4 gene can reverse resistance to trastuzumab in HER2/ErbB2-positive gastric cancers [37,38].

Xenograft tumors of estrogen receptor positive (ER)/ErbB2-overexpressing breast cancer cells, that are developing resistance to lapatinib and trastuzumab, harbor an increase in mucin-filled vacuoles and upregulation of several mucins including MUC4 [39] suggesting a role of MUC4 in acquired resistance to chemotherapy. MUC4 influence on chemosensitivity has mainly been studied in pancreatic cancer cells using gain or loss of function strategies. Several reports showed that MUC4 protects pancreatic cancer cells from gemcitabine-induced cytotoxicity [40–43]. Similar observation was made regarding another cytidine analog, the cytarabine/aracytin ARA-C [42] or 5-FU (unpublished data). Overexpression of rat ortholog MUC4/SMC in melanoma cells also reverts antiproliferative effect of taxol, doxorubicin, vinblastine, rhodamine-123 or 2-deoxyglucose and cell death induced by doxorubicin [44]. Finally MUC4 expression was shown to reduce the mitochondrial damage in pancreatic cancer cells induced by the inhibition

Table 1
Mucins and drug chemoresistance in epithelial cancer cells.

Mucin	Drug	Tumor type	Refs
<i>Membrane-bound mucins</i>			
MUC1	5-FU/methotrexate	CRC	[34,35]
	5-FU	PDAC	[58,60]
	Cisplatin	Ovarian	[52]
	Cisplatin	CRC	[67]
	Taxane/platinum compound	Ovarian	[53]
MUC3	Trastuzumab/paclitaxel/doxorubicin/cyclophosphamide	Breast cancer	[36]
	Gemcitabine/etoposide	PDAC	[90]
	Methotrexate	CRC	[34]
	5-FU	CRC	[34]
	Lapatinib/trastuzumab	Breast cancer	[39]
MUC4	Trastuzumab	Melanoma/ breast cancer	[65]
	Cytarabine/aracytin	PDAC	[42]
	Paclitaxel/doxorubicin/vinblastine/ rhodamine-123/2-deoxyglucose	Melanoma	[44]
	Bortezomib	PDAC	[43]
	Gemcitabine	PDAC	[40–42]
<i>Secreted mucins</i>			
MUC5AC	5-FU/methotrexate	CRC	[34,35]
MUC5B	5-FU/methotrexate	CRC	[34,35]
MUC2	5-FU	CRC	[34]

of the proteasome when treated by the boronic acid derivative bortezomib [43].

3. Mucins and clinical response to chemotherapy

3.1. Mucinous tumors

Retrospective clinical studies indicate that mucinous carcinoma or adenocarcinoma (uterine, colorectal or breast cancer), characterized by an abundant mucus gel covering the tumor, is less sensitive to neoadjuvant chemotherapies and radiotherapies. These data support that a link exist between the mucus and chemotherapeutic efficiency in the tumor context [45–48]. In advanced gastric cancer, the signet ring cell (SRC) histologic subtype, characterized by cells containing a large mucus vacuole, is an independent predictor of poor prognosis. Moreover, perioperative chemotherapy provides no benefit on survival [49]. However, the direct involvement of secreted mucins in these clinical observations as well as the associated underlying cellular mechanisms remains to be determined.

3.2. Non-mucinous tumors

On the contrary to mucinous tumors, a fair number of studies have shown that tumors expressing MUC1 exhibit an increased capacity in resisting the effects of chemotherapeutic drugs in breast and ovarian cancers. Frequency analysis of MUC1 expression in an important cohort of 691 breast cancer biopsies showed that high expression of both MUC1 mRNA and protein was associated with lower probability of complete response to neoadjuvant chemotherapy [50]. MUC1 regulates cholesterol and fatty acid metabolism in human breast cancer. Activation of these pathways in estrogen receptors (ER)+ breast cancers predicts failure to tamoxifen treatment with a higher risk for death and recurrence/distant metastasis [51].

MUC1 protein level is significantly increased in platinum-resistant ovarian tumors compared to those in sensitive-group because of chromosomal amplifications of 1q21–q22 [52]. A follow-up analysis of 92 ovarian cancer patient cohort also showed that the increase of MUC1 expression is a significant risk factor for chemoresistance to taxane combined platinum-based drugs in patients with ovarian epithelial cancer [53].

4. Mucins form a physical barrier

The outstanding web formed by the secreted mucus as well as the extraordinary size of heavily glycosylated membrane-bound mucins such as MUC1, MUC4 or MUC16 ($>10^6$ Da) suggest that mucins are capable of limiting (i) drug intracellular entrance and/or (ii) the accessibility of the plasma membrane and its tumor cell epitopes against immune recognition or antibody-based therapy (Fig. 1A).

Because of the viscosity/rheology of the mucus, secreted mucins may stop poorly soluble drug diffusion through the cell membrane. In diseases characterized by dense mucus such as chronic inflammatory cystic fibrosis, the mucus permeability is also decreased making the diffusion more difficult for therapeutics molecules [54]. Two functions are hypothesized: Mucins either act as size filters and allow entrance of drugs smaller than the mucus network porosity, or they act as interaction filters *via* electrostatic or hydrophobic forces accordingly to their surface properties as ionization of the drug was shown to influence its diffusion through mucus [55,56]. The size, the electric charge and the hydrophilic properties of mucin molecules composing the mucus are critical for the permeability. The electronegative potential of mucins, mostly carried by sialic acids on *O*-glycosidic chains, could alter the interaction properties of mucus [20] and may create electrostatic interactions with positively charged molecules (such as amikacin or gemcitabine) thereby decreasing their diffusion [57].

The steric hindrance of membrane-bound mucins is closely linked to the number and length of *O*-glycosidic chains they bare. The benzyl-2-acetamido-2-deoxy- α -D-galactopyranoside (benzyl- α -GalNAc) is a reagent used to inhibit the synthesis of mucin-type *O*-glycosylation. The inhibition of *O*-glycosylation in MUC1-expressing CAPAN-1 and HPAF-II pancreatic tumor cells resulted in significant 5-FU antiproliferative activity [58]. Benzyl- α -GalNAc that leads to a storage phenotype and abnormal intracellular localization of apical glycoproteins, does not alter directly mucin expression in CAPAN-1 cells since it was previously shown that MUC1 keeps its normal localization [59]. Therefore, Kalra and Campbell speculated that the inhibition of mucin glycosylation may reduce the formidable mucin *O*-glycosylation mesh and facilitate the entrance of chemotherapeutic drug [58]. In this model, the mucin network is thought to limit the intracellular uptake of 5-FU and to attenuate its chemotherapeutic effect independently of the sialic acid moiety [60]. The mucus layer on the surface of normal epithelial cells was also shown to limit the diffusion of nutrients and small molecules depending on their size and physicochemical properties [61].

The overexpression of membrane-bound mucins on the target tumor cell surface may mask the surface antigens, and thereby decrease their accessibility and the cytotoxic response induced by antibody-based therapy as well as the tumor cell killing mediated by immune cells. It has been shown that overexpression of rMuc4, MUC1 or MUC16 can create an immunosuppressive barrier by decreasing the accessibility of immune cells (as lymphokine-activated killer cell) to antigenic epitope dependent on *O*-glycosylation length [62–64]. The overexpression of membrane-bound mucins blocks drug accessibility to therapeutic targets such as oncogenic receptors EGFR, ErbB2 and VEGFR targeted by Erbitux/Cetuximab, Herceptin/trastuzumab and Avastin/Bevacizumab, respectively. The overexpression of rMuc4 in pancreatic cancer cell has been shown to decrease the recognition and thus the efficiency of the monoclonal antibody trastuzumab (Herceptin) which targets the oncogenic receptor ErbB2 [65]. A converse correlation has been established between the expression of MUC4 and Herceptin sensitivity in mammary cancer cells. The JIMT-1 cell line expressing the highest level of MUC4 is highly resistant to this monoclonal antibody [66].

Collectively, these data suggest that the steric hindrance induced by both secreted and membrane-bound mucins leads to limited drug intracellular entrance and reduced accessibility to the cell surface for targeted therapies.

5. Mucins and resistance to apoptosis

The induction of apoptosis is the main goal of all cytotoxic therapy. Resistance to programmed cell death is a major cause of chemoresistance in which expression of mucins may reduce the sensitivity of cancer cells to genotoxic drugs by decreasing the apoptotic effect following DNA damage or physiological stress (Fig. 1B).

5.1. MUC1

MUC1 role as an apoptotic inhibitor in cancer cells is well-documented (Fig. 2). Indeed, MUC1 decreases apoptotic mitochondrial factor release, caspase-3 activation, and subsequent apoptosis induction in CRC cells treated with the genotoxic agent cisplatin [67]. MUC1 cytoplasmic tail (MUC1-CT) inhibits the activation of a large array of intrinsic apoptotic pathways by regulating different signaling pathways such as p53, FOXO3a, c-Abl, I κ B complex, FADD or Bax. MUC1-CT interacts with p53 tumor suppressor on its responsive elements to coactivate p21 at the transcriptional level. On the contrary, MUC1 attenuates Bax transcription [68]. MUC1-CT also associates with Bax in the cytoplasm and the mitochondria *via* their CQC motif and BH3 domains, respectively, and blocks Bax dimerisation and subsequent cytochrome c release that normally activates the mitochondrial death pathway [69]. MUC1-CT anti-apoptotic function is also mediated *via* c-Abl sequestration




Fig. 1. Mucins and chemoresistance in epithelial cells. (A) Secreted and membrane-bound mucins form a physical barrier blocking cell accessibility to therapeutic drugs. Mucin overexpression is associated with (B) resistance to apoptosis, (C) drug metabolism (exclusion or reduction of drug intake), (D) cell stemness and (E) epithelial–mesenchymal transition (EMT). Cancer stem cells (CSC) belong to the small subset of cells (side population) commonly expressing CSC markers such as CD133 visualized here by flow cytometry. EMT is characterized by the loss of cell polarity, decreased expression of epithelial markers and increased expression of mesenchymal markers (e.g. vimentin).




Fig. 2. MUC1 overexpression and associated cellular mechanisms of chemoresistance in epithelial cancer cells.

within the cytoplasm and blocking of c-Abl and cytosolic 14-3-3 interaction [70]. Mutation of MUC1-CT at Tyr 60 (Y60F) disrupts the MUC1–Abl interaction inducing ARF expression and inhibiting MDM2–p53 pathway [71]. MUC1 is important for physiological activation of I κ B kinase- β (IKK β) complex and sustained activation of NF- κ B pathway in response to TNF- α activation [72]. In response to TNF- α , MUC1-CT also interacts with caspase-8 and Fas-associated death domain following death receptor stimulation and blocks the activation of apoptosis process [73]. In clear renal cell carcinoma (cRCC), overexpression of MUC1 prevents anoikis through anti-apoptotic NF- κ B nuclear localization, caspase-9 expression and the increase of Bcl_XL/Bax ratio [74].

Normal cellular metabolism is associated with the production of reactive oxygen species (ROS) including superoxide ion (O₂⁻), hydrogen peroxide (H₂O₂), hydroxyl radicals, and nitric oxide. Increase in ROS levels can cause damage to DNA, proteins, and lipids [75,76]. MUC1 reduces apoptosis following oxidative stress by regulating endogenous and H₂O₂-induced intracellular levels of ROS. Reciprocally, MUC1 expression is up-regulated by oxidative stress at the transcriptional level [77]. MUC1-CT also induces FOXO3a activation and attenuates the oxidative stress in colon and breast cancer cells [78]. Chronic ROS exposure leads to HIF-1 α stabilization and up-regulation of downstream target genes [79]. Among them, MUC1 is directly regulated by HIF-1 α in kidney cancer cells [80].

5.2. MUC4

An anti-apoptotic role has been described for MUC4 in serum-free conditions [81,82] as well as following drug treatment with thymoquinone, an antioxidant and anti-tumoral compound found in plants [83], or with gemcitabine [40,42]. MUC4 expression induces

apoptosis via multiple intracellular mechanisms (Fig. 3). It reduces mitochondrial cytochrome c release and activation of caspase-9 through phosphorylation of its partner ErbB2 and sequestration of pro-apoptotic Bad in the cytosol [40,41]. Loss of MUC4 oncoprotein is accompanied by a blockage in the G1-early S phases [41,84], an increased expression of the proapoptotic marker Bax and apoptotic mediator p53 as well as a decreased expression of the antiapoptotic Bcl_XL suggesting a higher susceptibility to apoptosis [42]. The overexpression of rMuc4 induces chemoresistance to paclitaxel, doxorubicin and cisplatin. In this case, rMuc4 expression decreases poly ADP ribose polymerase (PARP) and caspase-9 cleavage and modulates apoptosis [44,85]. rMuc4 may also repress apoptosis via an ErbB2-dependent mechanism, in which rMuc4 potentiates the activation of ErbB2 by phosphorylation of the Y1248 leading to activation of the PI3K/Akt pathway, and via an ErbB2-independent mechanism that remains to be determined [86,87]. MUC4 overexpression in ovarian cancer cells also leads to increased CD133⁺ cell population. CD133 is a pentaspan transmembrane glycoprotein associated with chemoresistance, radioresistance and poor prognosis and is commonly used as a stem cell marker (see Section 7). Additionally, CD133⁺ cell population demonstrates significant resistance to drug-, TGF- β - and TNF-Related Apoptosis-Inducing Ligand (TRAIL)-induced apoptosis compared with CD133⁻ cells [88,89] suggesting that CD133 triggers resistance to apoptosis in MUC4-expressing cells via similar mechanisms.

6. Mucins and alteration of drug metabolism

In pancreatic cancer, chemoresistance has been attributed partly to the enhanced expression of multidrug resistance (MDR) genes (Fig. 1C) including ATP-binding cassette transporter genes (ABCC)1, ABCC3, ABCC5 and ABCB1 that encode MDR related proteins (MRP)-1,




Fig. 3. MUC4 overexpression and associated cellular mechanisms of chemoresistance in epithelial cancer cells.

MRP3, MRP5 and MDR1 proteins, respectively [90]. The proteins encoded by this gene family are ATP-dependent drug efflux pumps for xenobiotic compounds with broad substrate specificity [91]. Notably, cleaved MUC1-CT directly associates and activates the ABCC1 promoter and increases (i) ABCC1 protein level in MUC1-high pancreatic cancer cells and (ii) resistance to gemcitabine and etoposide chemotherapeutic drugs [90] (Fig. 2).

One way to explain modifications of cell sensitivity to nucleoside analog such as gemcitabine or 5-FU is an alteration of the actors responsible for their metabolism and more particularly nucleoside transporters. We and others have described a major role for MUC4 in resistance to gemcitabine chemotherapy [40–42] involving alteration of nucleotide metabolism. In that case, MUC4 mucin negatively regulates the human concentrative nucleoside transporter-1 (hCNT1) transporter expression via the NF- κ B pathway, pointing out to MUC4 and hCNT1 as potential new targets to ameliorate the response of pancreatic tumors to gemcitabine treatment [42].

7. Mucins and cancer stem cells

Tumor ability to relapse after therapy is frequently dependent on a small subset of the cell population within the tumor, called cancer stem cells (CSC) or side population (SP) (Fig. 1D), characterized by an extensive self-renewing capacity [92]. Some therapies including chemotherapy may provide strong selection for CSC survival and proliferation [93].

Very recently, MUC1-CT was shown to upregulate expression of breast CSC marker Aldehyde dehydrogenase 1A1 (ALDH1A1) via Erk1 and C/EBP β by forming a transcriptional activating complex on the *ALDH1A1* gene promoter [94]. MUC1 was also shown to be expressed by CD34⁺ CD38[−] acute myeloid leukemia (AML) cells which have been associated with leukemia stem cells (LSC) suggesting that MUC1 represents a potential target on the AML stem cell population [95].

In ovarian cancer, MUC4 overexpression leads to increased CD133-positive CSC [96] (Fig. 3). Stem cell-like marker CD133 is also observed in a small subpopulation of pancreatic epithelial cells in the basal compartment in non-malignant pancreatic tissue specimens. This subpopulation also expresses MUC4 mucin [41]. MUC4 oncoprotein which is normally not expressed in the normal pancreas, is expressed at a high level in both the small CD133⁺ cell progenitor subpopulation as well as their differentiated CD133[−] progenies. Mimeault et al. have suggested that MUC4 down-regulation may partially reverse the resistance of CD133⁺ initiating cells to the gemcitabine treatment [41]. This is notably important since gemcitabine treatment of pancreatic tumor xenografts leads to enrichment of cell subpopulations expressing stem cell markers such as ALDH and CD24 as well as efflux pumps such as ABCB1 and ABCG2 [97,98]. MUC4 oncoprotein may thus represent a promising therapeutic strategy in that matter.

8. Mucins and epithelial–mesenchymal transition

Epithelial–mesenchymal transition (EMT) is a reversible pathophysiological process associated with loss of cell polarity, decreased surface expression of epithelial markers (E-cadherin, cytokeratin-18, ZO-1...) and increased expression of mesenchymal markers (vimentin, N-cadherin, MMP-9, ZEB-1...) (Fig. 1E) [99,100]. Among EMT inducers are found transcription factors, such as Snail, Slug, twist1/2, members of the transforming growth factor (TGF β) pathway or miRNAs [101, 102]. Many reports show that EMT is a major step toward metastatic tumor progression and contributes to drug resistance and acquisition or selection of stem/progenitor-like features and ultimately recurrence [103–106].

MUC1, MUC4 and MUC16 are able to trigger the molecular process of EMT [107–110]. The mucin-induced EMT implies interaction with β -catenin that leads to cell–cell junction disruption and invasiveness. Interestingly, MUC1-CT interacts with β -catenin and translocates to

the nucleus to upregulate EMT initiating genes. Moreover, MUC1-CT occupies and induces the *ZEB1* promoter in a NF- κ B p65-dependent manner. In turn *ZEB1* associates with MUC1-CT and contributes to suppress miR200c expression that promotes EMT [111]. In cRCC, MUC1-CT also occupies the *Snail* promoter, modulates the binding of β -catenin and triggers EMT [112]. The cellular mechanisms underlying EMT that are related to MUC4 or MUC16 are not fully understood but might involve signaling pathway deregulation.

The relation between MUC4 and EMT is controversial since MUC4 was shown to suppress EMT in lung adenocarcinoma [113,114] whereas MUC4 overexpression in ovarian cancer led to decreased expression of epithelial markers and occurrence of mesenchymal markers via an upregulation of Twist1, Twist2 and Snail transcription factors and FAK signaling pathway [110]. Collectively, these results suggest that MUC4 regulation of EMT is cellular dependent. Additional studies need to be carried out in order to elucidate these discrepancies regarding MUC4.

One may hypothesize that targeting EMT using MUC1 or MUC16 mucin as a therapeutic approach may be effective since these mucins are involved in both chemoresistance and EMT that often lead to tumor recurrence. Recently and in favor of that hypothesis, MUC1 siRNA was shown to inhibit both cell proliferation and EMT in urinary bladder cancer [115].

9. Mucins and polymorphisms associated with chemosensitivity?

As genomic sequencing cost declines rapidly, the availability of pangenomic information promotes the era of pharmacogenomics and therefore personalized medicine. The genetic diversity between different populations and ethnic backgrounds might explain a high degree of variability of drug response and adverse drug reactions [116]. Frequency differences of polymorphisms exist across different geographic regions for drug transporters or genes involved in drug metabolism [117,118].

Mucin single nucleotide polymorphisms (SNP), associated with various diseases including cancer, have been described [119,120]. Moreover, the SNP distributions differ between subjects from different origins (Caucasian or Asian) [121]. Mucin genes also exhibit a high degree of polymorphisms in the VNTR domain [122]. The variation of VNTR length leads to quantitative differences of the O-glycan content and alters biophysical properties of the glycoprotein. Among disease-associated mucins, gastric cancer patients harbor a higher proportion of short MUC1 allele compared to a control population [123]. Despite these data, no direct association between mucin polymorphism (SNP or VNTR) and sensitivity to chemotherapeutic drugs has been described so far.

10. Outlook to the future: Using mucins as a therapeutic target to sensitize cancer cells to chemotherapeutic drugs or as biomarkers of chemoresistance?

Since oncomucin expression in cancer cells is linked to a higher susceptibility to apoptosis, an increased expression of multidrug resistance markers, and stem cell characteristics, could represent a valuable therapeutic strategy to decrease tumor resistance. Because of their high molecular weight and their high sugar content, extraction and purification of native mucins for therapy is notoriously difficult [124]. Therefore, native mucins have never been used concomitantly with chemotherapeutic drugs. However, mucin expression or functionality has been targeted to sensitize cancer cells to chemotherapeutic drugs. Peptide inhibitors targeting MUC1-CT were shown to block its interaction with NF- κ B and to activate the induction of late apoptosis/necrosis in myeloma cells [125].

Silencing of MUC1 C-t subunit in ErbB2-overexpressing breast cancer cells or treatment with the penetrating peptide inhibitor, G0-203, reverses chemoresistance to trastuzumab, tamoxifen, taxol and doxorubicin (Fig. 2) [126–128]. Silencing MUC1-CT is associated with

(i) downregulation of ErbB2 phosphorylation and (ii) sensitivity to tamoxifen-induced growth inhibition and loss of clonogenic survival. ErbB2-positive cancer cells that are intrinsically resistant to trastuzumab became sensitive when treated with an antagonist of the cleaved form of the MUC1 protein [36]. In bladder cancer, MUC1 silencing also leads to reduction of the tumor volume *in vivo* and suppresses EMT. However, the impact on chemosensitivity remains to be proven in this case [115].

Currently, phase 1 multi-center clinical trial testing G0-203 dose-escalation, safety, pharmacodynamics and pharmacokinetics in solid tumors and lymphomas is in progress (ClinicalTrials.gov identifier: NCT01279603). G0-203 also results in tumor regression, independently of chemotherapy, in non-small cell lung cancer cells and acute myeloid leukemia [95,129].

Numerous studies are already targeting MUC1 as an immunotherapeutic approach [130–133]. Mucins could also be used in modified vectors sensitizing the tumor to targeted chemotherapy. In OVCAR-3 ovarian cancer cells, a MUC1/let-7i chimera, combines MUC1 aptamer and let-7i miRNA, can specifically reverse chemoresistance to paclitaxel [134]. Similarly a MUC1/miR-29b chimera also reverses chemoresistance to paclitaxel [135]. However, these effects on chemoresistance are independent of MUC1 structure since the MUC1 aptamer alone does not influence the induced apoptosis.

Therapy-predictive markers are factors that prospectively identify response or resistance to a specific treatment. They allow classification of patients with the same histological subtype of malignancy that respond very differently to a specific drug [136]. In CRC, the presence of acellular mucin pools in the resected specimens of patients with a complete response to preoperative chemoradiation is frequently reported as a marker of treatment effect. However, several reports ruled out its prognostic significance [137–139].

An ideal marker implicates non-invasive and rapid procedures of detection. Therefore, clinicians routinely use robust circulating markers in the blood or markers such as tumor-associated antigens. MUC1 (CA15-3) and MUC16 (CA125) tumor-associated antigens are commonly detected in patients' sera by standardized tests such as ELISA or immunohistochemistry [136,140]. MUC1 and MUC16 are overexpressed in ovarian cancer [17,141]. Budiu *et al.*, showed that increased serum MUC1 level has a prognostic value for poor clinical response and reduced overall survival in platinum-resistant or platinum-refractory ovarian cancer whereas MUC16 was not associated with clinical response [140].

In a multivariable analysis, MUC1 protein and mRNA expression were independently predictive in a breast cancer cohort for therapy response and survival after neoadjuvant chemotherapy [50]. CA15.3 level is also correlated with treatment response in patients undergoing chemotherapy for metastatic breast cancer [142]. Detection of (MUC1)-positive circulating tumor cells and MUC1 protein in the peripheral blood of patients with metastatic breast cancer is thus associated with higher progression-free survival (PFS) [143].

11. Conclusion

Over the past two decades, spectacular advances in targeted therapy led to improvement of treatment of a great numbers of cancer patients. These outstanding results are counterbalanced by the increased occurrence of acquired or intrinsic events of chemoresistance that limit long term success. Many pieces of evidence support the involvement of both membrane-bound and secreted mucins in diverse biological mechanisms of resistance (physical barrier, resistance to apoptosis, drug metabolism, cell stemness, EMT). Understanding mechanisms of resistance involving mucins shall contribute to the development of next generation targeted therapy molecules. Alternatively, mucins will help in the prevention because of their potential as tumor biomarkers and orientate the therapeutic choice tree toward the potentially more

successful therapy and more importantly avoid a more aggressive alternate therapy that is promised to failure.

Acknowledgements

We are grateful to Dr M. Perrais and Dr A. Vincent (Inserm UMR837, Lille) for their critical reading of the manuscript. This research was supported by SIRIC ONCOLille, Grant INCa-DGOS-Inserm 6041, by a grant from “Contrat de Plan Etat Région” CPER Cancer 2007–2013, and by a grant from la Ligue Nationale Contre le Cancer (Equipe Labellisée Ligue 2010, IVS). Isabelle Van Seuningen is the recipient of a “Contrat Hospitalier de Recherche Translationnelle”/CHRT 2010, AVIESAN.

References

- [1] P. Roussel, G. Lamblin, M. Lhermitte, N. Houdret, J.J. Lafitte, J.M. Perini, A. Klein, A. Scharfman, The complexity of mucins, *Biochimie* 70 (1988) 1471–1482.
- [2] J. Dekker, J.W. Rossen, H.A. Buller, A.W. Einerhand, The MUC family: an obituary, *Trends Biochem. Sci.* 27 (2002) 126–131.
- [3] J.L. Desseyn, A.P. Spicer, Epithelial mucin genes, *Annu. Rev. Physiol.* 57 (1995) 607–634.
- [4] D.J. Thornton, K. Rousseau, M.A. McGuckin, Structure and function of the polymeric mucins in airways mucus, *Annu. Rev. Physiol.* 70 (2008) 459–486.
- [5] J.L. Desseyn, V. Gouyer, D. Tetaert, Architecture of the gel-forming mucins, in: I. Van Seuningen (Ed.), *The Epithelial Mucins: Structure/Function. Roles in Cancer and Inflammatory Diseases*, Research Signpost, Kerala, India, 2008, pp. 1–16.
- [6] S.E. Baldus, K. Engelmann, F.G. Hanisch, MUC1 and the MUCs: a family of human mucins with impact in cancer biology, *Crit. Rev. Clin. Lab. Sci.* 41 (2004) 189–231.
- [7] D.T. Tran, K.G. Ten Hagen, Mucin-type O-glycosylation during development, *J. Biol. Chem.* 288 (2013) 6921–6929.
- [8] S. Groux-Degroote, A. Harduin-Lepers, P. Delannoy, Biosynthesis of mucin O-glycan chains in normal and pathological states, in: I. Van Seuningen (Ed.), *The Epithelial Mucins: Structure/Function. Roles in Cancer and Inflammatory Diseases*, Research Signpost, Kerala, India, 2008, pp. 39–54.
- [9] J.L. Desseyn, Mucin CYS domains are ancient and highly conserved modules that evolved in concert, *Mol. Phylogenet. Evol.* 52 (2009) 284–292.
- [10] D. Ambort, M.E. Johansson, J.K. Gustafsson, A. Ermund, G.C. Hansson, Perspectives on mucus properties and formation—lessons from the biochemical world, *Cold Spring Harb. Perspect. Med.* 2 (2012) pii: a014159.
- [11] M. Backstrom, D. Ambort, E. Thomsson, M.E. Johansson, G.C. Hansson, Increased understanding of the biochemistry and biosynthesis of MUC2 and other gel-forming mucins through the recombinant expression of their protein domains, *Mol. Biotechnol.* 54 (2013) 250–256.
- [12] C.L. Hattstrup, S.J. Gendler, Structure and function of the cell surface (tethered) mucins, *Annu. Rev. Physiol.* 70 (2008) 431–457.
- [13] N. Jonckheere, N. Skrypek, F. Frenois, I. Van Seuningen, Membrane-bound mucin modular domains: from structure to function, *Biochimie* 95 (2013) 1077–1086.
- [14] N. Jonckheere, I. Van Seuningen, The membrane-bound mucins: how large O-glycoproteins play key roles in epithelial cancers and hold promise as biological tools for gene-based and immunotherapies, *Crit. Rev. Oncog.* 14 (2008) 177–196.
- [15] J.L. Desseyn, D. Tetaert, V. Gouyer, Architecture of the large membrane-bound mucins, *Gene* 410 (2008) 215–222.
- [16] B.H. Perez, I.K. Gipson, Focus on molecules: human mucin MUC16, *Exp. Eye Res.* 87 (2008) 400–401.
- [17] N. Jonckheere, I. Van Seuningen, The membrane-bound mucins: from cell signalling to transcriptional regulation and expression in epithelial cancers, *Biochimie* 92 (2010) 1–11.
- [18] I. Van Seuningen, P. Pigny, M. Perrais, N. Porchet, J.P. Aubert, Transcriptional regulation of the 11p15 mucin genes. Towards new biological tools in human therapy, in inflammatory diseases and cancer? *Front. Biosci.* 6 (2001) D1216–D1234.
- [19] R. Pinto, A.S. Carvalho, T. Conze, A. Magalhaes, G. Picco, J.M. Burchell, J. Taylor-Papadimitriou, C.A. Reis, R. Almeida, U. Mandel, H. Clausen, O. Soderberg, L. David, Identification of new cancer biomarkers based on aberrant mucin glycoforms by in situ proximity ligation, *J. Cell. Mol. Med.* 16 (2012) 1474–1484.
- [20] C. Robbe-Masselot, J.C. Michalski, C. Capon, Tumour-associated antigens of mucin O-glycans. Clinical relevance of glycobiology in cancer and inflammatory diseases of the epithelium, in: I. Van Seuningen (Ed.), *The Epithelial Mucins: Structure/Function. Roles in Cancer and Inflammatory Diseases*, Research Signpost, Kerala, India, 2008, pp. 55–74.
- [21] P. Argueso, I.K. Gipson, Assessing mucin expression and function in human ocular surface epithelia in vivo and in vitro, *Methods Mol. Biol.* 842 (2012) 313–325.
- [22] M.A. Hollingsworth, B.J. Swanson, Mucins in cancer: protection and control of the cell surface, *Nat. Rev. Cancer* 4 (2004) 45–60.
- [23] N. Jonckheere, N. Skrypek, I. Van Seuningen, Mucins and pancreatic cancer, *Cancers* 2 (2010) 1794–1812.
- [24] S. Kaur, S. Kumar, N. Momi, A.R. Sasson, S.K. Batra, Mucins in pancreatic cancer and its microenvironment, *Nat. Rev. Gastroenterol. Hepatol.* 10 (2013) 607–620.
- [25] D.W. Kufe, Mucins in cancer: function, prognosis and therapy, *Nat. Rev. Cancer* 9 (2009) 874–885.
- [26] H. Hoshi, T. Sawada, M. Uchida, H. Saito, H. Iijima, M. Toda-Agetsuma, T. Wada, S. Yamazoe, H. Tanaka, K. Kimura, A. Kakehashi, M. Wei, K. Hirakawa, H. Wanibuchi, Tumor-associated MUC5AC stimulates in vivo tumorigenicity of human pancreatic cancer, *Int. J. Oncol.* 38 (2011) 619–627.
- [27] H. Valque, V. Gouyer, F. Gottrand, J.L. Desseyn, MUC5B leads to aggressive behavior of breast cancer MCF7 cells, *PLoS ONE* 7 (2012) e46699.
- [28] S. Yamazoe, H. Tanaka, T. Sawada, R. Amano, N. Yamada, M. Ohira, K. Hirakawa, RNA interference suppression of mucin 5 AC (MUC5AC) reduces the adhesive and invasive capacity of human pancreatic cancer cells, *J. Exp. Clin. Cancer Res.* 29 (2010) 53.
- [29] S.C. Chauhan, M.C. Ebeling, D.M. Maher, M.D. Koch, A. Watanabe, H. Aburatani, Y. Lio, M. Jaggi, MUC13 mucin augments pancreatic tumorigenesis, *Mol. Cancer Ther.* 11 (2012) 24–33.
- [30] D.M. Maher, B.K. Gupta, S. Nagata, M. Jaggi, S.C. Chauhan, Mucin 13: structure, function, and potential roles in cancer pathogenesis, *Mol. Cancer Res.* 9 (2011) 531–537.
- [31] I. Lakshmanan, M.P. Ponnusamy, S. Das, S. Chakraborty, D. Haridas, P. Mukhopadhyay, S.M. Lele, S.K. Batra, MUC16 induced rapid G2/M transition via interactions with JAK2 for increased proliferation and anti-apoptosis in breast cancer cells, *Oncogene* 31 (2012) 805–817.
- [32] A. Velcich, W. Yang, J. Heyer, A. Fragale, C. Nicholas, S. Viani, R. Kuchlerlapati, M. Lipkin, K. Yang, L. Augenlicht, Colorectal cancer in mice genetically deficient in the mucin Muc2, *Science* 295 (2002) 1726–1729.
- [33] R. Dahiya, T. Lesuffleur, K.S. Kwak, J.C. Byrd, A. Barbat, A. Zweibaum, Y.S. Kim, Expression and characterization of mucins associated with the resistance to methotrexate of human colonic adenocarcinoma cell line HT29, *Cancer Res.* 52 (1992) 4655–4662.
- [34] T. Lesuffleur, N. Porchet, J.P. Aubert, D. Swallow, J.R. Gum, Y.S. Kim, F.X. Real, A. Zweibaum, Differential expression of the human mucin genes MUC1 to MUC5 in relation to growth and differentiation of different mucus-secreting HT-29 cell subpopulations, *J. Cell Sci.* 106 (Pt 3) (1993) 771–783.
- [35] E. Leteurtre, V. Gouyer, K. Rousseau, O. Moreau, A. Barbat, D. Swallow, G. Huet, T. Lesuffleur, Differential mucin expression in colon carcinoma HT-29 clones with variable resistance to 5-fluorouracil and methotrexate, *Biol. Cell.* 96 (2004) 145–151.
- [36] S.P. Fessler, M.T. Wotkowicz, S.K. Mahanta, C. Bamdad, MUC1* is a determinant of trastuzumab (Herceptin) resistance in breast cancer cells, *Breast Cancer Res. Treat.* 118 (2009) 113–124.
- [37] M. Deng, D.D. Jing, X.J. Meng, Effect of MUC1 siRNA on drug resistance of gastric cancer cells to trastuzumab, *Asian Pac. J. Cancer Prev.* 14 (2013) 127–131.
- [38] M. Shi, Z. Yang, M. Hu, D. Liu, Y. Hu, L. Qian, W. Zhang, H. Chen, L. Guo, M. Yu, L. Song, Y. Ma, N. Guo, Catecholamine-induced beta2-adrenergic receptor activation mediates desensitization of gastric cancer cells to trastuzumab by upregulating MUC4 expression, *J. Immunol.* 190 (2013) 5600–5608.
- [39] A.C. Chen, I. Migliaccio, M. Rimawi, S. Lopez-Tarruella, C.J. Creighton, S. Massarweh, C. Huang, Y.C. Wang, S.K. Batra, M.C. Gutierrez, C.K. Osborne, R. Schiuff, Upregulation of mucin4 in ER-positive/HER2-overexpressing breast cancer xenografts with acquired resistance to endocrine and HER2-targeted therapies, *Breast Cancer Res. Treat.* 134 (2012) 583–593.
- [40] S. Bafna, S. Kaur, N. Momi, S.K. Batra, Pancreatic cancer cells resistance to gemcitabine: the role of MUC4 mucin, *Br. J. Cancer* 101 (2009) 1155–1161.
- [41] M. Mimeault, S.L. Johansson, S. Senapati, N. Momi, S. Chakraborty, S.K. Batra, MUC4 down-regulation reverses chemoresistance of pancreatic cancer stem/progenitor cells and their progenies, *Cancer Lett.* 295 (2010) 69–84.
- [42] N. Skrypek, B. Duchene, M. Hebbard, E. Leteurtre, I. Van Seuningen, N. Jonckheere, The MUC4 mucin mediates gemcitabine resistance of human pancreatic cancer cells via the Concentrative Nucleoside Transporter family, *Oncogene* 32 (2013) 1714–1723.
- [43] T.T. Wisniewski, S. Meister, E.G. Hahn, J.R. Kalden, R. Voll, M. Ocker, Mucin production determines sensitivity to bortezomib and gemcitabine in pancreatic cancer cells, *Int. J. Oncol.* 40 (2012) 1581–1589.
- [44] Y.P. Hu, B. Haq, K.L. Carraway, N. Savaraj, T.J. Lampidis, Multidrug resistance correlates with overexpression of Muc4 but inversely with P-glycoprotein and multidrug resistance related protein in transfected human melanoma cells, *Biochem. Pharmacol.* 65 (2003) 1419–1425.
- [45] L.J. Mekenkamp, K.J. Heesterbeek, M. Koopman, J. Tol, S. Teerstra, S. Venderbosch, C.J. Punt, I.D. Nagtegaal, Mucinous adenocarcinomas: poor prognosis in metastatic colorectal cancer, *Eur. J. Cancer* 48 (2012) 501–509.
- [46] T. Nagao, T. Kinoshita, T. Hojo, H. Tsuda, K. Tamura, Y. Fujiwara, The differences in the histological types of breast cancer and the response to neoadjuvant chemotherapy: the relationship between the outcome and the clinicopathological characteristics, *Breast* 21 (2012) 289–295.
- [47] K. Oberholzer, M. Menig, A. Kreft, A. Schneider, T. Junginger, A. Heintz, K.F. Kreitner, A.M. Hotker, T. Hansen, C. Duber, H. Schmidberger, Rectal cancer: mucinous carcinoma on magnetic resonance imaging indicates poor response to neoadjuvant chemoradiation, *Int. J. Radiat. Oncol. Biol. Phys.* 82 (2012) 842–848.
- [48] O. Poujade, P. Morice, R. Rouzier, P. Madelenat, F. Lecuru, J.M. Muray, P. Mathevet, S. Alran, R.J. Salmon, F. Rey, Pathologic response rate after concomitant neoadjuvant radiotherapy and chemotherapy for adenocarcinoma of the uterine cervix: a retrospective multicentric study, *Int. J. Gynecol. Cancer* 20 (2010) 815–820.
- [49] M. Messager, J.H. Lefevre, V. Pichot-Delahaye, A. Souadka, G. Piessen, C. Mariette, The impact of perioperative chemotherapy on survival in patients with gastric signet ring cell adenocarcinoma: a multicenter comparative study, *Ann. Surg.* 254 (2011) 684–693 (discussion 693).
- [50] B.V. Sinn, G. von Minckwitz, C. Denkert, H. Eidtmann, S. Darb-Esfahani, H. Tesch, R. Kronenwett, G. Hoffmann, A. Belau, C. Thommsen, H.J. Holzhausen, S.T. Grasshoff, K. Baumann, K. Mehta, M. Dietel, S. Loibl, Evaluation of Mucin-1 protein and mRNA expression as prognostic and predictive markers after neoadjuvant chemotherapy for breast cancer, *Ann. Oncol.* 24 (2013) 2316–2324.

- [51] S.P. Pitroda, N.N. Khodarev, M.A. Beckett, D.W. Kufe, R.R. Weichselbaum, MUC1-induced alterations in a lipid metabolic gene network predict response of human breast cancers to tamoxifen treatment, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 5837–5841.
- [52] M. Takano, K. Fujii, T. Kita, Y. Kikuchi, K. Uchida, Amplicon profiling reveals cytoplasmic overexpression of MUC1 protein as an indicator of resistance to platinum-based chemotherapy in patients with ovarian cancer, *Oncol. Rep.* 12 (2004) 1177–1182.
- [53] D. Zhang, J. Gao, L. Zhu, Z. Hu, R. Hou, S. Liu, M. Tan, J. Liu, B. Lin, Chemoresistance is associated with MUC1 and Lewis y antigen expression in ovarian epithelial cancers, *Int. J. Mol. Sci.* 14 (2013) 11024–11033.
- [54] P.G. Bhat, D.R. Flanagan, M.D. Donovan, Drug diffusion through cystic fibrotic mucus: steady-state permeation, rheologic properties, and glycoprotein morphology, *J. Pharm. Sci.* 85 (1996) 624–630.
- [55] L.R. Shaw, W.J. Irwin, T.J. Grattan, B.R. Conway, The influence of excipients on the diffusion of ibuprofen and paracetamol in gastric mucus, *Int. J. Pharm.* 290 (2005) 145–154.
- [56] H.H. Sigurdsson, J. Kirch, C.M. Lehr, Mucus as a barrier to lipophilic drugs, *Int. J. Pharm.* 453 (2013) 56–64.
- [57] K. Khanvilkar, M.D. Donovan, D.R. Flanagan, Drug transfer through mucus, *Adv. Drug Deliv. Rev.* 48 (2001) 173–193.
- [58] A.V. Kalra, R.B. Campbell, Mucin impedes cytotoxic effect of 5-FU against growth of human pancreatic cancer cells: overcoming cellular barriers for therapeutic gain, *Br. J. Cancer* 97 (2007) 910–918.
- [59] E. Leteurtre, V. Gouyer, D. Delacour, B. Hemon, A. Pons, C. Richet, J.P. Zanetta, G. Huet, Induction of a storage phenotype and abnormal intracellular localization of apical glycoproteins are two independent responses to GalNAc α -0-bn, *J. Histochem. Cytochem.* 51 (2003) 349–361.
- [60] A.V. Kalra, R.B. Campbell, Mucin overexpression limits the effectiveness of 5-FU by reducing intracellular drug uptake and antineoplastic drug effects in pancreatic tumours, *Eur. J. Cancer* 45 (2009) 164–173.
- [61] G. Lafitte, K. Thuresson, O. Soderman, Diffusion of nutrients molecules and model drug carriers through mucin layer investigated by magnetic resonance imaging with chemical shift resolution, *J. Pharm. Sci.* 96 (2007) 258–263.
- [62] J.A. Gubbels, M. Felder, S. Horibata, J.A. Belisle, A. Kapur, H. Holden, S. Petrie, M. Migneault, C. Rancourt, J.P. Connor, M.S. Patankar, MUC16 provides immune protection by inhibiting synapse formation between NK and ovarian tumor cells, *Mol. Cancer* 9 (2010) 11.
- [63] M. Komatsu, L. Yee, K.L. Carraway, Overexpression of sialomucin complex, a rat homologue of MUC4, inhibits tumor killing by lymphokine-activated killer cells, *Cancer Res.* 59 (1999) 2229–2236.
- [64] E. van de Wiel-van Kemenade, M.J. Ligtenberg, A.J. de Boer, F. Buijs, H.L. Vos, C.J. Melief, J. Hilken, C.G. Figdor, Episialin (MUC1) inhibits cytotoxic lymphocyte-target cell interaction, *J. Immunol.* 151 (1993) 767–776.
- [65] S.A. Price-Schiavi, S. Jepson, P. Li, M. Arango, P.S. Rudland, L. Yee, K.L. Carraway, Rat Muc4 (sialomucin complex) reduces binding of anti-ErbB2 antibodies to tumor cell surfaces, a potential mechanism for herceptin resistance, *Int. J. Cancer* 99 (2002) 783–791.
- [66] P. Nagy, E. Friedlander, M. Tanner, A.I. Kapanen, K.L. Carraway, J. Isola, T.M. Jovin, Decreased accessibility and lack of activation of ErbB2 in JIMT-1, a herceptin-resistant, MUC4-expressing breast cancer cell line, *Cancer Res.* 65 (2005) 473–482.
- [67] J. Ren, N. Agata, D. Chen, Y. Li, W.H. Yu, L. Huang, D. Raina, W. Chen, S. Kharbanda, D. Kufe, Human MUC1 carcinoma-associated protein confers resistance to genotoxic anticancer agents, *Cancer Cell* 5 (2004) 163–175.
- [68] X. Wei, H. Xu, D. Kufe, Human MUC1 oncoprotein regulates p53-responsive gene transcription in the genotoxic stress response, *Cancer Cell* 7 (2005) 167–178.
- [69] R. Ahmad, M. Alam, H. Rajabi, D. Kufe, The MUC1-C oncoprotein binds to the BH3 domain of the pro-apoptotic BAX protein and blocks BAX function, *J. Biol. Chem.* 287 (2012) 20866–20875.
- [70] D. Raina, R. Ahmad, S. Kumar, J. Ren, K. Yoshida, S. Kharbanda, D. Kufe, MUC1 oncoprotein blocks nuclear targeting of c-Abl in the apoptotic response to DNA damage, *EMBO J.* 25 (2006) 3774–3783.
- [71] D. Raina, R. Ahmad, D. Chen, S. Kumar, S. Kharbanda, D. Kufe, MUC1 oncoprotein suppresses activation of the ARF-MDM2-p53 pathway, *Cancer Biol. Ther.* 7 (2008) 1959–1967.
- [72] R. Ahmad, D. Raina, V. Trivedi, J. Ren, H. Rajabi, S. Kharbanda, D. Kufe, MUC1 oncoprotein activates the IkappaB kinase beta complex and constitutive NF-kappaB signalling, *Nat. Cell Biol.* 9 (2007) 1419–1427.
- [73] N. Agata, R. Ahmad, T. Kawano, D. Raina, S. Kharbanda, D. Kufe, MUC1 oncoprotein blocks death receptor-mediated apoptosis by inhibiting recruitment of caspase-8, *Cancer Res.* 68 (2008) 6136–6144.
- [74] A. Bouillez, V. Gnemmi, K. Gaudelot, B. Hemon, B. Ringot, N. Pottier, F. Glowacki, C. Butruille, C. Cauffiez, M. Hamdane, N. Sergeant, I. Van Seuningen, X. Leroy, S. Aubert, M. Perrais, MUC1-C nuclear localization drives invasiveness of renal cancer cells through a sheddase/gamma secretase dependent pathway, *Oncotarget* 5 (2014) 754–763.
- [75] A.T. Lau, Y. Wang, J.F. Chiu, Reactive oxygen species: current knowledge and applications in cancer research and therapeutic, *J. Cell. Biochem.* 104 (2008) 657–667.
- [76] P. Storz, Reactive oxygen species in tumor progression, *Front. Biosci.* 10 (2005) 1881–1896.
- [77] L. Yin, Y. Li, J. Ren, H. Kuwahara, D. Kufe, Human MUC1 carcinoma antigen regulates intracellular oxidant levels and the apoptotic response to oxidative stress, *J. Biol. Chem.* 278 (2003) 35458–35464.
- [78] L. Yin, L. Huang, D. Kufe, MUC1 oncoprotein activates the FOXO3a transcription factor in a survival response to oxidative stress, *J. Biol. Chem.* 279 (2004) 45721–45727.
- [79] J.P. Fruehauf, F.L. Meyskens Jr., Reactive oxygen species: a breath of life or death? *Clin. Cancer Res.* 13 (2007) 789–794.
- [80] S. Aubert, V. Fauquette, B. Hemon, R. Lepoivre, N. Briez, D. Bernard, I. Van Seuningen, X. Leroy, M. Perrais, MUC1, a new hypoxia inducible factor target gene, is an actor in clear renal cell carcinoma tumor progression, *Cancer Res.* 69 (2009) 5707–5715.
- [81] P. Chaturvedi, A.P. Singh, N. Moniaux, S. Senapati, S. Chakraborty, J.L. Meza, S.K. Batra, MUC4 mucin potentiates pancreatic tumor cell proliferation, survival, and invasive properties and interferes with its interaction to extracellular matrix proteins, *Mol. Cancer Res.* 5 (2007) 309–320.
- [82] N. Moniaux, P. Chaturvedi, G.C. Varshney, J.L. Meza, J.F. Rodriguez-Sierra, J.P. Aubert, S.K. Batra, Human MUC4 mucin induces ultra-structural changes and tumorigenicity in pancreatic cancer cells, *Br. J. Cancer* 97 (2007) 345–357.
- [83] M.P. Torres, M.P. Ponnusamy, S. Chakraborty, L.M. Smith, S. Das, H.A. Arafat, S.K. Batra, Effects of thymoquinone in the expression of mucin 4 in pancreatic cancer cells: implications for the development of novel cancer therapies, *Mol. Cancer Ther.* 9 (2010) 1419–1431.
- [84] N. Jonckheere, N. Skrypek, J. Merlin, A.F. Dessein, P. Dumont, E. Leteurtre, A. Harris, J.L. Desseyn, C. Susini, F. Frenois, I. Van Seuningen, The mucin MUC4 and its membrane partner ErbB2 regulate biological properties of human CAPAN-2 pancreatic cancer cells via different signalling pathways, *PLoS ONE* 7 (2012) e32232.
- [85] M. Komatsu, S. Jepson, M.E. Arango, C.A. Carothers Carraway, K.L. Carraway, Muc4/sialomucin complex, an intramembrane modulator of ErbB2/HER2/Neu, potentiates primary tumor growth and suppresses apoptosis in a xenotransplanted tumor, *Oncogene* 20 (2001) 461–470.
- [86] S. Jepson, M. Komatsu, B. Haq, M.E. Arango, D. Huang, C.A. Carraway, K.L. Carraway, Muc4/sialomucin complex, the intramembrane ErbB2 ligand, induces specific phosphorylation of ErbB2 and enhances expression of p27(kip), but does not activate mitogen-activated kinase or protein kinaseB/Akt pathways, *Oncogene* 21 (2002) 7524–7532.
- [87] H.C. Workman, C. Sweeney, K.L. Carraway III, The membrane mucin Muc4 inhibits apoptosis induced by multiple insults via ErbB2-dependent and ErbB2-independent mechanisms, *Cancer Res.* 69 (2009) 2845–2852.
- [88] A. El-Khattouti, D. Selimovic, Y. Haikel, M. Megahed, C.R. Gomez, M. Hassan, Identification and analysis of CD133⁺ melanoma stem-like cells conferring resistance to taxol: an insight into the mechanisms of their resistance and response, *Cancer Lett.* 343 (2014) 123–133.
- [89] J. Li, X.Y. Zhong, Z.Y. Li, J.F. Cai, L. Zou, J.M. Li, T. Yang, W. Liu, CD133 expression in osteosarcoma and derivation of CD133⁺ cells, *Mol. Med. Rep.* 7 (2013) 577–584.
- [90] S. Nath, K. Daneshvar, L.D. Roy, P. Grover, A. Kidiyoor, L. Mosley, M. Sahraei, P. Mukherjee, MUC1 induces drug resistance in pancreatic cancer cells via upregulation of multidrug resistance genes, *Oncogenesis* 2 (2013) e51.
- [91] F.J. Sharom, ABC multidrug transporters: structure, function and role in chemoresistance, *Pharmacogenomics* 9 (2008) 105–127.
- [92] M. Greaves, C.C. Maley, Clonal evolution in cancer, *Nature* 481 (2012) 306–313.
- [93] P.B. Gupta, T.T. Onder, G. Jiang, K. Tao, C. Kuperwasser, R.A. Weinberg, E.S. Lander, Identification of selective inhibitors of cancer stem cells by high-throughput screening, *Cell* 138 (2009) 645–659.
- [94] M. Alam, R. Ahmad, H. Rajabi, A. Kharbanda, D. Kufe, MUC1-C oncoprotein activates ERK → C/EBPbeta signaling and induction of aldehyde dehydrogenase 1A1 in breast cancer cells, *J. Biol. Chem.* 288 (2013) 30892–30903.
- [95] D. Stroopinsky, J. Rosenblatt, K. Ito, H. Mills, L. Yin, H. Rajabi, B. Vasir, T. Kufe, R. Luptakova, J. Arnason, C. Nardella, J.D. Levine, R.M. Joyce, I. Galinsky, Y. Reiter, K. M. Stone, P.P. Pandolfi, D. Kufe, D. Avigan, MUC1 is a potential target for the treatment of acute myeloid leukemia stem cells, *Cancer Res.* 73 (2013) 5569–5579.
- [96] M.P. Ponnusamy, P. Seshacharyulu, A. Vaz, P. Dey, S.K. Batra, MUC4 stabilizes HER2 expression and maintains the cancer stem cell population in ovarian cancer cells, *J. Ovarian Res.* 4 (2011) 7.
- [97] A. Jimeno, G. Feldmann, A. Suarez-Gauthier, Z. Rasheed, A. Solomon, G.M. Zou, B. Rubio-Viqueira, E. Garcia-Garcia, F. Lopez-Rios, W. Matsui, A. Maitra, M. Hidalgo, A direct pancreatic cancer xenograft model as a platform for cancer stem cell therapeutic development, *Mol. Cancer Ther.* 8 (2009) 310–314.
- [98] J. Zhou, C.Y. Wang, T. Liu, B. Wu, F. Zhou, J.X. Xiong, H.S. Wu, J. Tao, G. Zhao, M. Yang, S.M. Gou, Persistence of side population cells with high drug efflux capacity in pancreatic cancer, *World J. Gastroenterol.* 14 (2008) 925–930.
- [99] J.P. Thiery, H. Adloque, R.Y. Huang, M.A. Nieto, Epithelial–mesenchymal transitions in development and disease, *Cell* 139 (2009) 871–890.
- [100] J.P. Thiery, J.P. Sleeman, Complex networks orchestrate epithelial–mesenchymal transitions, *Nat. Rev. Mol. Cell Biol.* 7 (2006) 131–142.
- [101] S. Lamouille, D. Subramanyam, R. Blelloch, R. Derynck, Regulation of epithelial–mesenchymal and mesenchymal–epithelial transitions by microRNAs, *Curr. Opin. Cell Biol.* 25 (2013) 200–207.
- [102] M.K. Wendt, M. Tian, W.P. Schiemann, Deconstructing the mechanisms and consequences of TGF-beta-induced EMT during cancer progression, *Cell Tissue Res.* 347 (2012) 85–101.
- [103] N. Ahmed, K. Abubaker, J. Findlay, M. Quinn, Epithelial mesenchymal transition and cancer stem cell-like phenotypes facilitate chemoresistance in recurrent ovarian cancer, *Curr. Cancer Drug Targets* 10 (2010) 268–278.
- [104] S. Lim, A. Becker, A. Zimmer, J. Lu, R. Buettner, J. Kirfel, SNAI1-mediated epithelial–mesenchymal transition confers chemoresistance and cellular plasticity by regulating genes involved in cell death and stem cell maintenance, *PLoS ONE* 8 (2013) e66558.
- [105] A. Voulgari, A. Pintzas, Epithelial–mesenchymal transition in cancer metastasis: mechanisms, markers and strategies to overcome drug resistance in the clinic, *Biochim. Biophys. Acta* 1796 (2009) 75–90.
- [106] S. Yamada, B.C. Fuchs, T. Fujii, Y. Shimoyama, H. Sugimoto, S. Nomoto, S. Takeda, K. Kanabe, Y. Kodera, A. Nakao, Epithelial-to-mesenchymal transition predicts prognosis of pancreatic cancer, *Surgery* 154 (2013) 946–954.

- [107] M. Comamala, M. Pinard, C. Theriault, I. Matte, A. Albert, M. Boivin, J. Beaudin, A. Piche, C. Rancourt, Downregulation of cell surface CA125/MUC16 induces epithelial-to-mesenchymal transition and restores EGFR signalling in NIH:OVCA3 ovarian carcinoma cells, *Br. J. Cancer* 104 (2011) 989–999.
- [108] G. Horn, A. Gazieli, D.H. Wreschner, N.I. Smorodinsky, M. Ehrlich, ERK and PI3K regulate different aspects of the epithelial to mesenchymal transition of mammary tumor cells induced by truncated MUC1, *Exp. Cell Res.* 315 (2009) 1490–1504.
- [109] L.D. Roy, M. Sahraei, D.B. Subramani, D. Besmer, S. Nath, T.L. Tindler, E. Bajaj, K. Shanmugam, Y.Y. Lee, S.I. Hwang, S.J. Gendler, P. Mukherjee, MUC1 enhances invasiveness of pancreatic cancer cells by inducing epithelial to mesenchymal transition, *Oncogene* 30 (2011) 1449–1459.
- [110] M.P. Ponnusamy, I. Lakshmanan, M. Jain, S. Das, S. Chakraborty, P. Dey, S.K. Batra, MUC4 mucin-induced epithelial to mesenchymal transition: a novel mechanism for metastasis of human ovarian cancer cells, *Oncogene* 29 (2010) 5741–5754.
- [111] H. Rajabi, M. Alam, H. Takahashi, A. Kharbanda, M. Guha, R. Ahmad, D. Kufe, MUC1-C oncoprotein activates the ZEB1/miR-1330 regulatory loop and epithelial-mesenchymal transition, *Oncogene* 33 (2014) 1680–1689.
- [112] V. Gnemmi, A. Bouillez, K. Gaudelot, B. Hemon, B. Ringot, N. Pottier, F. Glowacki, A. Villers, D. Vindrieux, C. Cauffiez, I.V. Seuningen, D. Bernard, X. Leroy, S. Aubert, M. Perrais, MUC1 drives epithelial-mesenchymal transition in renal carcinoma through Wnt/beta-catenin pathway and interaction with SNAIL promoter, *Cancer Lett.* 346 (2014) 225–236.
- [113] L. Gao, J. Liu, B. Zhang, H. Zhang, D. Wang, T. Zhang, Y. Liu, C. Wang, Functional MUC4 suppress epithelial-mesenchymal transition in lung adenocarcinoma metastasis, *Tumour Biol.* 35 (2014) 1335–1341.
- [114] N. Jonckheere, I. Van Seuningen, Comment on: Functional MUC4 suppress epithelial-mesenchymal transition in lung adenocarcinoma metastasis. Gao L, Liu J, Zhang B, Zhang H, Wang D, Zhang T, Liu Y, Wang C. *Tumour Biol.* 2013, in press, *Tumour Biol.* 35 (2014) 3941–3942.
- [115] T. Fujii, K. Shimada, S. Anai, K. Fujimoto, N. Konishi, ALKBH2, a novel AlkB homologue, contributes to human bladder cancer progression by regulating MUC1 expression, *Cancer Sci.* 104 (2013) 321–327.
- [116] J.W. Lee, F. Amineng, A.P. Bhavsar, K. Shaw, B.C. Carleton, M.R. Hayden, C.J. Ross, The emerging era of pharmacogenomics: current successes, future potential, and challenges, *Clin. Genet.* (2014), <http://dx.doi.org/10.1111/cge.12392> (in press).
- [117] C. Ono, H. Kikkawa, A. Suzuki, M. Suzuki, Y. Yamamoto, K. Ichikawa, M. Fukae, I. Ieiri, Clinical impact of genetic variants of drug transporters in different ethnic groups within and across regions, *Pharmacogenomics* 14 (2013) 1745–1764.
- [118] K. Sai, Y. Saito, Ethnic differences in the metabolism, toxicology and efficacy of three anticancer drugs, *Expert Opin. Drug Metab. Toxicol.* 7 (2011) 967–988.
- [119] C.Y. Chang, H.W. Chang, C.M. Chen, C.Y. Lin, C.P. Chen, C.H. Lai, W.Y. Lin, H.P. Liu, J.J. Sheu, F.J. Tsai, MUC4 gene polymorphisms associate with endometriosis development and endometriosis-related infertility, *BMC Med.* 9 (2011) 19.
- [120] N. Saeki, A. Saito, I.J. Choi, K. Matsuo, S. Ohnami, H. Totsuka, S. Chiku, A. Kuchiba, Y.S. Lee, K.A. Yoon, M.C. Kook, S.R. Park, Y.W. Kim, H. Tanaka, K. Tajima, H. Hirose, F. Tanioka, Y. Matsuno, H. Sugimura, S. Kato, T. Nakamura, T. Nishina, W. Yasui, K. Aoyagi, H. Sasaki, K. Yanagihara, H. Katai, T. Shimoda, T. Yoshida, Y. Nakamura, S. Hirohashi, H. Sakamoto, A functional single nucleotide polymorphism in mucin 1, at chromosome 1q22, determines susceptibility to diffuse-type gastric cancer, *Gastroenterology* 140 (2011) 892–902.
- [121] Y. Horimasu, N. Hattori, N. Ishikawa, S. Kawase, S. Tanaka, K. Yoshioka, A. Yokoyama, N. Kohno, F. Bonella, J. Guzman, S. Ohshimo, U. Costabel, Different MUC1 gene polymorphisms in German and Japanese ethnicities affect serum KL-6 levels, *Respir. Med.* 106 (2012) 1756–1764.
- [122] J. Fowler, L. Vinall, D. Swallow, Polymorphism of the human muc genes, *Front. Biosci.* 6 (2001) D1207–D1215.
- [123] F. Carvalho, R. Seruca, L. David, A. Amorim, M. Seixas, E. Bennett, H. Clausen, M. Sobrinho-Simoes, MUC1 gene polymorphism and gastric cancer—an epidemiological study, *Glycoconj. J.* 14 (1997) 107–111.
- [124] J.R. Davies, C. Wickstrom, D.J. Thornton, Gel-forming and cell-associated mucins: preparation for structural and functional studies, *Methods Mol. Biol.* 842 (2012) 27–47.
- [125] L. Yin, R. Ahmad, M. Kosugi, T. Kufe, B. Vasir, D. Avigan, S. Kharbanda, D. Kufe, Survival of human multiple myeloma cells is dependent on MUC1 C-terminal transmembrane subunit oncoprotein function, *Mol. Pharmacol.* 78 (2010) 166–174.
- [126] A. Kharbanda, H. Rajabi, C. Jin, D. Raina, D. Kufe, Oncogenic MUC1-C promotes tamoxifen resistance in human breast cancer, *Mol. Cancer Res.* 11 (2013) 714–723.
- [127] D. Raina, Y. Uchida, A. Kharbanda, H. Rajabi, G. Panchamoorthy, C. Jin, S. Kharbanda, M. Scaltriti, J. Baselga, D. Kufe, Targeting the MUC1-C oncoprotein downregulates HER2 activation and abrogates trastuzumab resistance in breast cancer cells, *Oncogene* (2014), <http://dx.doi.org/10.1038/ncr.2013.308> (in press).
- [128] Y. Uchida, D. Raina, S. Kharbanda, D. Kufe, Inhibition of the MUC1-C oncoprotein is synergistic with cytotoxic agents in the treatment of breast cancer cells, *Cancer Biol. Ther.* 14 (2013) 127–134.
- [129] D. Raina, M. Kosugi, R. Ahmad, G. Panchamoorthy, H. Rajabi, M. Alam, T. Shimamura, G.I. Shapiro, J. Supko, S. Kharbanda, D. Kufe, Dependence on the MUC1-C oncoprotein in non-small cell lung cancer cells, *Mol. Cancer Ther.* 10 (2011) 806–816.
- [130] C.K. Tang, V. Apostolopoulos, Strategies used for MUC1 immunotherapy: preclinical studies, *Expert Rev. Vaccines* 7 (2008) 951–962.
- [131] C.K. Tang, M. Katsara, V. Apostolopoulos, Strategies used for MUC1 immunotherapy: human clinical studies, *Expert Rev. Vaccines* 7 (2008) 963–975.
- [132] V. Lakshminarayanan, P. Thompson, M.A. Wolfert, T. Buskas, J.M. Bradley, L.B. Pathangey, C.S. Madsen, P.A. Cohen, S.J. Gendler, G.J. Boons, Immune recognition of tumor-associated mucin MUC1 is achieved by a fully synthetic aberrantly glycosylated MUC1 tripartite vaccine, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2011) 261–266.
- [133] R.E. Beatson, J. Taylor-Papadimitriou, J.M. Burchell, MUC1 immunotherapy, *Immunotherapy* 2 (2010) 305–327.
- [134] N. Liu, C. Zhou, J. Zhao, Y. Chen, Reversal of paclitaxel resistance in epithelial ovarian carcinoma cells by a MUC1 aptamer-let-7i chimera, *Cancer Investig.* 30 (2012) 577–582.
- [135] F. Dai, Y. Zhang, X. Zhu, N. Shan, Y. Chen, The anti-chemoresistant effect and mechanism of MUC1 aptamer-miR-29b chimera in ovarian cancer, *Gynecol. Oncol.* 131 (2013) 451–459.
- [136] M.J. Duffy, Tumor markers in clinical practice: a review focusing on common solid cancers, *Med. Princ. Pract.* 22 (2013) 4–11.
- [137] L.F. de Campos-Lobato, D.W. Dietz, L. Stocchi, J.D. Vogel, I.C. Lavery, J.R. Goldblum, M. Skacel, R.J. Pelley, M.F. Kalady, Clinical implications of acellular mucin pools in resected rectal cancer with pathological complete response to neoadjuvant chemoradiation, *Colorectal Dis.* 14 (2012) 62–67.
- [138] K.D. Smith, D. Tan, P. Das, G.J. Chang, K. Kattepogu, B.W. Feig, J.M. Skibber, M.A. Rodriguez-Bigas, Clinical significance of acellular mucin in rectal adenocarcinoma patients with a pathologic complete response to preoperative chemoradiation, *Ann. Surg.* 251 (2010) 261–264.
- [139] J. Shia, M. McManus, J.G. Guillem, T. Leibold, Q. Zhou, L.H. Tang, E.R. Riedel, M.R. Weiser, P.B. Paty, L.K. Temple, G. Nash, K. Kolosov, B.D. Minsky, W.D. Wong, D.S. Klimstra, Significance of acellular mucin pools in rectal carcinoma after neoadjuvant chemoradiotherapy, *Am. J. Surg. Pathol.* 35 (2011) 127–134.
- [140] R.A. Budiu, G. Mantia-Smaldone, E. Elishaev, T. Chu, J. Thaller, K. McCabe, D. Lenzner, R.P. Edwards, A.M. Vlad, Soluble MUC1 and serum MUC1-specific antibodies are potential prognostic biomarkers for platinum-resistant ovarian cancer, *Cancer Immunol. Immunother.* 60 (2011) 975–984.
- [141] S. Aubert, I. Van Seuningen, X. Leroy, Mucins in the uro-genital tract. Potential for therapeutic approaches using mucins, in: I. Van Seuningen (Ed.), *The Epithelial Mucins: Structure/Function. Roles in Cancer and Inflammatory Diseases*, Research Signpost, Kerala, India, 2008, pp. 249–272.
- [142] D. Di Gioia, V. Heinemann, D. Nagel, M. Untch, S. Kahlert, I. Bauerfeind, T. Koehnke, P. Stieber, Kinetics of CEA and CA15-3 correlate with treatment response in patients undergoing chemotherapy for metastatic breast cancer (MBC), *Tumour Biol.* 32 (2011) 777–785.
- [143] J.P. Cheng, Y. Yan, X.Y. Wang, Y.L. Lu, Y.H. Yuan, J. Jia, J. Ren, MUC1-positive circulating tumor cells and MUC1 protein predict chemotherapeutic efficacy in the treatment of metastatic breast cancer, *Chin. J. Cancer* 30 (2011) 54–61.