

Enhanced CH₄ emissions from a wetland soil exposed to Elevated CO₂

J. PATRICK MEGONIGAL* & W.H. SCHLESINGER

Department of Botany, Duke University, Durham, NC 27708, U.S.A. (* Current address:
Department of Biology, George Mason University, Fairfax, VA 22030-4444)

Accepted 11 September 1996

Key words: climate change, elevated carbon dioxide, hydrology, methane, transpiration, wetlands

Abstract. Methane emissions from wetland soils are generally a positive function of plant size and primary productivity, and may be expected to increase due to enhanced rates of plant growth in a future atmosphere of elevated CO₂. We performed two experiments with *Orontium aquaticum*, a common emergent aquatic macrophyte in temperate and sub-tropical wetlands, to determine if enhanced rates of photosynthesis in elevated CO₂ atmospheres would increase CH₄ emissions from wetland soils. *O. aquaticum* was grown from seed in soil cores under ambient and elevated (ca. 2-times ambient) concentrations of CO₂ in an initial glasshouse study lasting 3 months and then a growth chamber study lasting 6 months. Photosynthetic rates were 54 to 71% higher under elevated CO₂ than ambient CO₂, but plant biomass was not significantly different at the end of the experiment. In each case, CH₄ emissions were higher under elevated than ambient CO₂ levels after 2 to 4 months of treatment, suggesting a close coupling between photosynthesis and methanogenesis in our plant-soil system. Methane emissions in the growth chamber study increased by 136%. We observed a significant decrease in transpiration rates under elevated CO₂ in the growth chamber study, and speculate that elevated CO₂ may also stimulate CH₄ emissions by increasing the extent and duration of flooding in some wetland ecosystems. Elevated CO₂ may dramatically increase CH₄ emissions from wetlands, a source that currently accounts for 40% of global emissions.

Introduction

Vascular plants enhance CH₄ emissions from wetlands. Carbon assimilation by plants is the ultimate source of organic substrates for methanogenic bacteria (Oremland 1988) and plant vascular tissue is a conduit for CH₄ diffusion from sediments to the atmosphere (Chanton & Dacey 1991). The positive correlation that exists between CH₄ emission rates and net ecosystem production in North American wetlands (Whiting & Chanton 1993) is evidence that plant productivity is a key process regulating CH₄ emission from wetlands at a regional scale. The central role of plant production is also supported by observations that amendments of rice straw stimulate methane production in paddy soils (Sass et al. 1991; Cicerone et al. 1992; Watanabe et al. 1995), and additions of H₂ or labile carbon compounds often increase CH₄ produc-

tion in laboratory soil incubations (Yavitt et al. 1987; Valentine et al. 1994). Environmental factors that influence photosynthesis or plant production in wetlands may indirectly regulate methane emissions.

Carbon dioxide concentrations have increased dramatically during this century, and are expected to double again by the end of the next century (IPCC 1995). Although it is well established that most plants respond to elevated CO_2 with increased rates of photosynthesis (Curtis 1996; Sage 1994; Gunderson & Wullschleger 1994), there are few studies of the impact of increased photosynthesis on other ecosystem processes (Field et al. 1992). Plant detritus provides organic carbon compounds that fuel most biogeochemical transformations in soils and sediments, and an increase in their supply may be expected to impact all heterotrophic soil processes, including methanogenesis.

There are other mechanisms by which elevated CO_2 could stimulate methane emissions. An increase in plant biomass may promote methane diffusion from soils by increasing the volume of porous vascular tissue. Lower transpiration rates – a common response of plants to elevated CO_2 – may raise water tables or lengthen periods of inundation. Each of these effects would stimulate CH_4 emissions from wetland ecosystems, which presently account for 40% of all global sources of CH_4 (Cicerone & Oremland 1988), and constitute important biological feedbacks on climate change.

Dacey et al. (1994) reported that elevated CO_2 increased CH_4 emissions from a brackish tidal marsh by 80% after 4 years of treatment. They suggested that this effect was due, in part, to an increase in leaf and root detritus in the high CO_2 treatment, although it was unclear how rapidly such an effect could occur. Hutchin et al. (1995) observed an increase in CH_4 emissions of 145% after exposing intact cores from an ombrotrophic mire to elevated CO_2 for just 6 weeks. Based on evidence that methanogenesis is limited by the availability of labile organic carbon compounds, we also proposed that elevated CO_2 would increase CH_4 emission rates over relatively short periods of time. A rapid response would indicate a tight temporal coupling between photosynthesis and methanogenesis in wetland ecosystems, although other mechanisms that could explain such an observation must be considered. The objective of our study was to determine if short-term exposures to elevated CO_2 would increase CH_4 emissions from a freshwater wetland plant-soil system.

Methods

We collected seeds of *Orontium aquaticum* from a tidal freshwater swamp on the North Carolina coast in spring of 1993 and 1994. *Orontium aquaticum* is

an emergent aquatic macrophyte that occurs throughout the southeastern US coastal plain (Odum et al. 1984). *O. aquaticum* is similar in morphology and appearance to emergent aquatic macrophytes that have been used in previous studies of CH₄ emissions [i.e., *Peltandra virginica* (Chanton et al. 1992) and *Sagittaria lancifolia* (Schipper & Reddy 1996)]. We chose this species because it is common at our field site and bears fruit in the spring when our studies began. Soils collected from the study site had a silty clay loam texture with 22% organic carbon content. Plants were raised from seed under ambient and elevated concentrations of CO₂ in trays of field soil (4 cm deep) that were continuously flooded with one-half strength Hoagland's solution.

We performed experiments in the summers of 1993 and 1994 to determine if elevated CO₂ would influence CH₄ emissions from a wetland plant-soil system over a relatively short period of time. The 1993 experiment was performed in a glasshouse using relatively small volume containers filled with a mixture of 75% sand and 25% field soil. The design of the study was intended to minimize the contribution of the native soil organic carbon to CH₄ production, so that we could more easily observe changes in the contribution of recent photosynthates. Based on the results of the 1993 study, we undertook a similar experiment in 1994 using controlled environmental chambers and unamended soil (without sand addition) from our field site.

Glasshouse experiment

Seedlings were transferred after 8 weeks (30 July 1993) into polyvinyl chloride (PVC) pots (7.5 cm-i.d. by 25 cm deep) filled 20 cm deep with a sand-soil mixture. Glasshouses at the Duke University Phytotron Facility were automatically controlled (Hellmers & Giles 1979) to maintain CO₂ partial pressures of 35 or 70 Pa. Plants were exposed to natural light intensity and photoperiod; photon flux densities in August were about 1500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at mid-day. Glasshouse temperatures were maintained on a day/night schedule of 27/22 C, with the thermoperiod adjusted to follow the photoperiod. Soils were continuously flooded with one-half strength Hoagland's solution and occasionally flushed with fresh nutrient solution to prevent salt accumulation. Evaporation of flood-water and algal growth were minimized by placing a 2 cm-thick styrofoam disk on the surface of each pot in both experiments.

Methane emissions were determined after 9 and 14 weeks of treatment using a static chamber technique. Both plants and soils were enclosed with a 7.6 cm-i.d. PVC chamber capped with a plastic end-cap that was sealed with silicon. The flux chambers were joined to the pots with a PVC coupling modified with a single internal O-ring at each end to ensure an air-tight seal. A rubber septum (Vacutainer stopper) in the end-cap was used to sample the

headspace 4 times over a period of 3 hours. Methane concentrations were immediately determined on a Varian 3700 gas chromatograph with a flame ionization detector, using a Porapak Q 80/100 mesh column at 50 C and He at 30 ml min⁻¹ as the carrier gas.

Photosynthesis, conductance, relative humidity and photon flux density were measured with a model 6200 Licor Infrared Gas Analyzer (IRGA)(Licor Inc., Lincoln, NE) inside the glasshouses at treatment CO₂ concentrations. We used the second youngest shoot on each plant to control for leaf age effects. The plants were harvested at the end of the experiment and roots were recovered on a 2 mm-mesh sieve by a combination of floating and washing. Mass was determined after drying at 70 C to constant weight.

Controlled chamber experiment

Soils were collected, sieved free of woody detritus and roots, and used to fill PVC pots (7.6 cm-i.d. by 40 cm deep) to a depth of 30 cm. After 4 weeks (14 July 1994), 32 similar-size seedlings were transferred into randomly assigned pots in each treatment. Six additional pots per treatment remained unplanted for a no-plant control. The plants were raised in growth chambers controlled at CO₂ partial pressures of either 35 or 72 Pa for a period of 6 months. Plants and CO₂ treatments were rotated at monthly intervals to mitigate possible chamber effects. Photon flux density was 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and temperature was 30C day/25 C night on a 12-hour cycle. Photosynthesis was measured under growth CO₂ concentrations at a photon flux density of 1300 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Gas exchange measurements were made as described for the glasshouse experiment. The plants were continuously flooded with one-half strength Hoagland's solution. Conductivity measurements in mid-September showed lower salt concentrations in the soil solution (mean = 0.787 m Ω) than in the pure Hoagland's solution (1.264 m Ω), with no significant difference between treatments ($P = 0.61$). Platinum redox electrodes were inserted horizontally into 4 pots per treatment at depths of 5 cm and 28 cm below the soil surface and allowed to equilibrate for 1 d before reading with a pH meter.

Eight plants from each treatment were harvested for biomass at the end of the experiment. The soils from 8 additional pots were sectioned at 4-cm intervals and dried at 110 C to constant weight to determine gravimetric water content. Fresh subsamples from these pots were centrifuged at 3500 rpm for 5 min and the supernatant was analyzed for SO₄ on a Dionex 2101i ion chromatograph.

Statistics

Methane fluxes were based on a linear-regression between CH₄ concentration inside static chambers and time (SAS Institute 1987). Because fluxes in the growth chamber experiment were high and easily measured (Figure 1), slopes with regression coefficients (r^2) < 0.90 were discarded (1 observation) and the chambers were compared with a t-test on untransformed data. Methane emissions in this experiment were normally distributed as determined by the Shapiro-Wilk Statistic using the SAS univariate procedure. In the glasshouse experiment, low regression coefficients (r^2 < 0.90) were not discarded because these data were assumed to indicate below-detection-limit emission rates, and this information was relevant to our hypothesis. Low emission rates were probably caused by a combination of sandy, low organic C soils and small pots. For the two-month sample, we used a parametric t-test on transformed data because the distributions were log-normal despite having 3 values (2 ambient, 1 elevated) below the detection limit. For the 3-month sample, the non-parametric Kuskal-Wallis test was used to compare treatments (SAS procedure nlin) because 10 values were below the detection limit (7 ambient, 3 elevated).

We considered tests of the median to be appropriate in this study because our pots were self-contained units that were wholly sampled for CH₄ emissions (Parkin & Robinson 1992). Statistical analyses were used to judge differences between experimental groups, but because glasshouses and chambers were not replicated, the presence or absence of a difference cannot be interpreted as unequivocal support for treatment effects.

Results and discussion

Methane emission rates were higher in elevated CO₂ than in ambient CO₂ atmospheres after two to four months of treatment. In the initial glasshouse experiment, the two groups were nearly significantly different after two months ($P = 0.06$), and after three months this difference was particularly dramatic ($P = 0.04$, Figure 1). We obtained a similar result when the study was repeated in growth chambers. Although there were no differences between the chambers during the first two months of the study, CH₄ emissions were higher in the elevated CO₂ chamber than in the ambient CO₂ chamber after four months ($P \leq 0.01$ in months 5 and 6). The magnitude of this increase (136%) suggests that future CH₄ emissions from wetlands may increase dramatically from a source that presently accounts for 40% of the CH₄ emissions globally. The response of our pots to elevated CO₂ was similar to the 145% increase reported by Hutchin et al. (1995) for cores from an ombrotrophic

Table 1. Gas exchange characteristics of *Orontium aquaticum* grown under ambient and elevated CO₂. Values are means $\pm$ 1 standard error, with significant differences indicated by asterisks on the larger mean. Relative humidity inside the glasshouses was similar to cuvette values; however, the relative humidity of the growth chambers (ca. 70%) was higher than cuvette levels.

Date	MPa [CO ₂]	Sample Size	Photosynthesis ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	Transpiration ($\text{mol m}^{-2} \text{s}^{-1}$)	Cuvette Rel. Humidity (%)
Glasshouse					
2 months	35	8	13.6 $\pm$ 0.42	7.57 $\pm$ 0.36	54 $\pm$ 4
	70	9	21.4 $\pm$ 0.99**	8.16 $\pm$ 0.28	65 $\pm$ 1
3 months	35	8	19.2 $\pm$ 0.80	7.43 $\pm$ 0.29*	59 $\pm$ 4
	70	9	32.9 $\pm$ 3.16*	5.66 $\pm$ 0.42	68 $\pm$ 2
Growth Chamber					
6 months	35	9	18.1 $\pm$ 0.66	14.11 $\pm$ 0.99	31 $\pm$ 1
	72	10	27.8 $\pm$ 1.62**	9.30 $\pm$ 0.95**	31 $\pm$ 1

* $P \leq 0.01$.

** $P \leq 0.001$.

mire, and higher than the 80% increase reported by Dacey et al. (1994) for soils in a brackish tidal marsh studied in situ. It appears that large increases in methane emissions in response to elevated CO₂ can occur in a wide variety of wetland ecosystems.

Net photosynthetic rates were 54–71% higher under elevated CO₂ (Table 1), a result consistent with studies of *Scirpus olneyi* (Arp & Drake 1991), *Eriophorum vaginatum* (Hutchin et al. 1995), and a large number of upland plant species (Curtis 1996; Sage 1994; Gunderson & Wullschlegel 1994). In a previous study of *Eriophorum vaginatum*, higher photosynthetic rates under elevated CO₂ conditions were short-lived due to down-regulation (Tissue & Oechel 1987). Our photosynthesis measurements were made on relatively young leaves only, and the actual increase in CO₂ assimilation on a whole-plant basis is likely to have been smaller than measured due to leaf aging effects (Field & Mooney 1983). Nonetheless, the data suggest that methane emissions increased because of higher photosynthetic activity and release of additional organic carbon to soils.

Plant biomass tended to be higher in elevated CO₂ than in ambient CO₂ atmospheres (Table 2), but there were no significant differences between the two groups in either experiment ($P > 0.10$). Coarse and fine root biomass comparisons in the growth chamber study had P-values of 0.11. Hutchin et al. (1995) also reported that high photosynthetic rates in elevated CO₂ environments did not increase biomass accumulation in an ombrotrophic mire community, and suggested that the excess organic carbon may have been




Figure 1. Methane emissions under ambient and elevated levels of atmospheric CO₂ in two independent experiments. Box plots for the glasshouse experiment encompass the 25th to 75th percentiles in emissions; horizontal caps represent the 10th and 90th percentiles; circles are the 5th and 95th percentiles. The median is indicated with a horizontal line inside the box. Values in the environmental chamber experiment are means $\pm$ 1 standard error.

released by root exudates. More rapid rates of leaf and root turnover under elevated than under ambient CO₂ conditions is another possible explanation for this result.

Table 2. Dry mass (g) of *Orontium aquaticum* plants at the end of two elevated CO₂ experiments. Values are means $\pm$ 1 standard error based on 9 samples in the glasshouse study and 8 samples in the growth chamber study. Change in biomass due to elevated CO₂ (Δ) was calculated as (elevated-ambient) $\div$ ambient. There were no significant differences in between-treatment comparisons ($P \geq 0.10$).

	Glasshouse			Growth Chamber		
	35 Pa	70 Pa	% Δ	38 pa	72 Pa	% Δ
Shoot	1.48 $\pm$ 0.14	1.91 $\pm$ 0.26	29	5.37 $\pm$ 0.40	5.95 $\pm$ 0.77	11
Fine Root ^a	nd	nd	nd	2.36 $\pm$ 0.18	2.81 $\pm$ 0.22	19
Course Root	nd	nd	nd	12.96 $\pm$ 0.79	15.20 $\pm$ 1.07	17
Total Root	2.51 $\pm$ 0.36	2.77 $\pm$ 0.32	10	15.32 $\pm$ 0.93	18.01 $\pm$ 1.27	18
Total Plant	3.99 $\pm$ 0.43	4.68 $\pm$ 0.56	17	20.69 $\pm$ 1.22	23.96 $\pm$ 1.97	16

^a nd = not determined

Water table depth and redox potential exert strong control over methane emissions from wetlands (Moore & Dalva 1993; Wang et al. 1993; Funk et al. 1994). We maintained anaerobic conditions in the soil profile by continuously flooding the surface (5 cm deep) and sealing the bottom of our pots with a rubber cap. Redox potentials indicated that the soils were anaerobic with no significant differences between treatments in depth-wise comparisons ($P > 0.27$, Table 3). Nonetheless, we found evidence that differences in transpiration rates (Table 1) affected the water balance in our pots. The gravimetric water content declined with depth in both treatments, indicating that transpiration rates were greater than water percolation rates in this fine-textured soil (Figure 2). The water content of soils with plants grown at ambient and elevated CO₂ were significantly different at 28 cm ($P = 0.05$). A higher gravimetric water content in elevated CO₂ pots, compared to ambient CO₂ pots, indicates a long-term shift toward a more positive water balance under elevated CO₂. Drake (1992) reported lower rates of whole-stand transpiration from elevated CO₂ than ambient CO₂ chambers in a brackish tidal marsh despite an increase in leaf biomass. Leaf biomass was similar between treatments in our study (Table 2).

Most wetlands experience periods when evapotranspiration and runoff exceed precipitation, causing the water table to fall and O₂ infusion into the soil (Meronigal et al. 1993; Faulkner & Patrick 1992). On sites where transpiration is an important component of the water budget, elevated CO₂ may extend the portion of the growing season when soils are saturated and anaerobic. Transpiration accounted for up to 74% of evapotranspiration losses from a tidal freshwater marsh in Virginia (Hussey & Odum 1992), and was shown to be a driving force for O₂ infusion into salt-marsh sediments (Dacey & Howes 1984). More studies that quantify the importance of transpiration in wetland




Figure 2. Gravimetric water content (%) at 4-cm intervals for soils in the environmental growth chamber experiment. Values are means $\pm$ 1 standard error.

Table 3. Redox potentials at two depths for pots with *Orontium aquaticum* plants exposed to ambient (38 Pa) or elevated (72 Pa) levels of CO₂, and for pots without plants. Values are means $\pm$ 1 standard error ($n = 4$ for planted pots, $n = 2$ for unplanted pots).

Treatment	5 cm Depth	28 cm Depth
	Redox Potential (mV)	
No plants	11 $\pm$ 41	58 $\pm$ 11
Ambient CO ₂	45 $\pm$ 17	-14 $\pm$ 61
Elevated CO ₂	71 $\pm$ 12	37 $\pm$ 16

water budgets are needed. Pulliam & Meyer (1992) developed an empirically-based simulation model of CH₄ emissions from a temperate wetland forest and concluded that emissions are likely to be more sensitive to future perturbations of the water budget than to changes in temperature. A transpiration feedback on CH₄ emissions would augment increased methanogenesis due to increased primary production.

Freeman et al. (1994) proposed that an experimental lowering of the groundwater table in a peatland increased pore-water SO₄ concentrations, through hydrogen sulfide oxidation, and caused competitive inhibition of methanogenesis by sulfate-reducing bacteria. We investigated the possibility that relatively high rates of transpiration had caused SO₄ levels to increase in the ambient CO₂ pots of the growth chamber study, but found that SO₄ levels were not significantly different in the two groups of pots (ambient mean $\pm$ se = 1.34 ± 0.12 mM, elevated = 0.97 ± 0.24 mM). Thus, the observed differences in CH₄ emissions apparently were not due to a transpiration-induced increase in SO₄-reduction in the ambient pots.

The porous aerenchyma tissue of vascular wetland plants provides a diffusion pathway for gas exchange between sediments and the atmosphere and greatly enhances CH₄ emissions (Chanton & Dacey 1991). This process may contribute to the positive correlation between CH₄ emissions and net ecosystem production observed for North American wetlands (Whiting & Chanton 1993). Although there was a tendency for greater plant biomass in the elevated CO₂ than in the ambient CO₂ treatments in our study (Table 2), the differences were not significant, and roots reached to the bottom of the pots in all treatments. It is unlikely that improved transport of CH₄ through plant tissues can explain the large differences in emissions that we observed.

In conclusion, we have shown that elevated CO₂ can dramatically increase CH₄ emissions from wetland soils and sediments. We believe the effect was due to increased rates of carbon input through root exudation or root turnover. We also speculate that elevated CO₂ may increase CH₄ emissions in some wetland ecosystems by decreasing rates of water loss by transpiration. The significance of this process will depend greatly on the importance of transpiration in the water budgets of particular wetland sites, and the extent to which decreases in transpiration on a leaf-area basis are negated by increases in leaf area.

Acknowledgements

This work was supported by NSF Dissertation Improvement Grant DEB 93-11143 and a NASA Global Change Fellowship to J.P.M., and by NSF grants BSR-87-06429 and DEB-94-15541 to the Duke University Phytotron

Facility. We thank Dwight Billings, Norm Christensen, Chis Martens, Curt Richardson, Boyd Strain and David Tissue for reviewing an early draft of the manuscript and providing advice on the research.

References

- Arp WJ & Drake BG (1991) Increased photosynthetic capacity of *Scirpus olneyi* after four years of exposure to elevated CO₂. *Plant Cell Environ.* 14: 1003–1006
- Chanton JP & Dacey JWH (1991) Effects of vegetation on methane flux, reservoirs, and carbon isotopic composition. In: Sharkey TD, Holland EA & Mooney HA (Eds) *Trace Gas Emissions by Plants* (pp 65–92). Academic Press, San Diego
- Chanton JP, Whiting GJ, Showers WJ & Crill PM (1992) Methane flux from *Peltandra virginica*: stable isotope tracing and chamber effects. *Global Biogeochem. Cycles* 6: 15–31
- Cicerone RJ, Delwiche CC, Tyler SC & Zimmerman PR (1992) Methane emissions from California rice paddies with varied treatments. *Global Biogeochem. Cycles* 6: 233–248
- Cicerone RJ & Oremland RS (1988) Biogeochemical aspects of atmospheric methane. *Global Biogeochem. Cycles* 2: 299–327
- Curtis PS (1996) A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant Cell Environ.* 19: 127–137
- Dacey JWH, Drake BG & Klug MJ (1994) Stimulation of methane emission by carbon dioxide enrichment of marsh vegetation. *Nature* 370: 47–49
- Dacey JWH & Howes BL (1984) Water uptake by roots controls water table movement and sediment oxidation in short *Spartina* marsh. *Science* 224: 487–489
- Drake BG (1992) A field study of the effects of elevated CO₂ on ecosystem processes in a Chesapeake Bay Wetland. *Aust. J. Bot.* 40: 579–595
- Faulkner SP & Patrick WH Jr (1992) Redox processes and diagnostic wetland soil indicators in bottomland hardwood forests. *Soil Sci. Soc. Am. J.* 56: 856–865
- Field CB, Chapin FS, Matson PA & Mooney HA (1992) Responses of terrestrial ecosystems to the changing atmosphere: A resource-based approach. *Ann. Rev. Ecol. and Syst.* 23: 201–235
- Field C & Mooney HA (1983) Leaf age and seasonal effects on light, water, and nitrogen use efficiency in a California shrub. *Oecologia* 56: 348–355
- Freeman C, Hudson J, Lock MA, Reynolds B & Swanson C (1994) A possible role of sulphate in the suppression of wetland methane fluxes following drought. *Soil Biol. Biochem.* 26: 1439–1442
- Funk DW, Pullman ER, Peterson KM, Crill PM & Billings WD (1994) Influence of water table on carbon dioxide, carbon monoxide, and methane fluxes from taiga bog microcosms. *Global Biogeochem. Cycles* 8: 271–278
- Gunderson CA & Wullschlegel SD (1994) Photosynthetic acclimation in trees to rising atmospheric CO₂: A broader perspective. *Photosynth. Res.* 39: 369–388
- Hellmers H & Giles LJ (1979) Carbon dioxide: critique I. In: Tibbitts TW & Kozlowski TT (Eds) *Controlled Environmental Guidelines for Plant Research* (pp 229–234). Academic Press, New York, NY
- Hussey BH & Odum WE (1992) Evapotranspiration in tidal marshes. *Estuaries* 15: 59–67
- Hutchinson PR, Press MC, Lee JA & Ashenden TW (1995) Elevated concentrations of CO₂ may double methane emissions from mires. *Global Change Biol.* 1: 125–128
- IPCC (1995) *Climate change 1995: The science of climate change*. Cambridge University Press, Cambridge, England
- Megonigal JP, Patrick WH Jr & Faulkner SP (1993) Wetland identification in seasonally flooded forest soils: Soil morphology and redox dynamics. *Soil Sci. Soc. Am. J.* 57: 140–149

- Moore TR & Dalva M (1993) The influence of temperature and water table position on carbon dioxide and methane emissions from laboratory columns of peatland soils. *J. Soil Science* 44: 651–664
- Odum WE, Smith TJ, Hoover JK & McIvor CC (1984) *The Ecology of Tidal Freshwater Marshes of the United States East Coast: A Community Profile*. United States Fish and Wildlife Service, FWS/OBS-83/17, Washington, DC
- Oremland RS (1988) Biogeochemistry of methanogenic bacteria. In: Zehnder AJB (Ed) *Biology of Anaerobic Microorganisms* (pp 641–705). John Wiley and Sons, New York, NY
- Parkin TB & Robinson JA (1992) Analysis of lognormal data. *Adv. Soil Sci.* 20: 193–235
- Pulliam WM & Meyer JL (1992) Methane emissions from floodplain swamps of the Ogeechee River: Long-term patterns and effects of climate change. *Biogeochem.* 15: 151–174
- Sage RF (1994) Acclimation of photosynthesis to increasing atmospheric CO₂: The gas exchange perspective. *Photosynth. Res.* 39: 351–368
- SAS Institute (1987) *SAS/STAT Guide for Personal Computers*, Version 6 edition. SAS Institute, Cary, NC
- Sass RL, Fisher FM, Harcombe PA & Turner FT (1991) Mitigation of methane emissions from rice fields: Possible adverse effects of incorporated rice straw. *Global Biogeochem. Cycles* 5: 275–287
- Schipper LA & Reddy KR (1996) Determination of methane oxidation in the rhizosphere of *Sagittaria lancifolia* using methyl fluoride. *Soil Sci. Soc. Am. J.* 60: 611–616
- Tissue DT & Oechel WC (1987) Response of *Eriophorum vaginatum* to elevated CO₂ and temperature in the Alaskan tussock tundra. *Ecology* 68: 401–410
- Valentine DW, Holland EA & Schimel DS (1994) Ecosystem and physiological controls over methane production in northern wetlands. *J. Geophys. Res.* 99: 1563–1571
- Wang ZP, DeLaune RD, Masscheleyn PH & Patrick WH, Jr (1993) Soil redox and pH effects on methane production in a flooded rice soil. *Soil Sci. Soc. Am. J.* 57: 382–385
- Watanabe A, Satoh Y & Kimura M (1995) Estimation of the increase in CH₄ emission from paddy soils by rice straw application. *Plant Soil* 173: 225–231
- Whiting GJ & Chanton JP (1993) Primary production control of methane emission from wetlands. *Nature* 364: 794–795
- Yavitt JB, Lang GE & Wieder RK (1987) Control of carbon mineralization to CH₄ and CO₂ in anaerobic, *Sphagnum*-derived peat from Big Run Bog, West Virginia. *Biogeochemistry* 4: 141–157