

Non-laminar cerebral cortex in teleost fishes?

Hironobu Ito and Naoyuki Yamamoto

Biol. Lett. 2009 **5**, 117-121
doi: 10.1098/rsbl.2008.0397

References

[This article cites 32 articles, 2 of which can be accessed free](#)
<http://rsbl.royalsocietypublishing.org/content/5/1/117.full.html#ref-list-1>

Article cited in:
<http://rsbl.royalsocietypublishing.org/content/5/1/117.full.html#related-urls>

Subject collections

Articles on similar topics can be found in the following collections

[neuroscience](#) (114 articles)
[evolution](#) (541 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Opinion piece

Non-laminar cerebral cortex in teleost fishes?

A large skull is disadvantageous to animals that move quickly in three-dimensional space, such as fishes and birds in water or air. A cerebral neocortex with a six-layered sheet has not evolved, most likely due to the limited cranial space. Instead of the laminar cortex, telencephalic nuclear masses seem to have evolved as the pallium in teleost fishes. We consider that the nuclear masses contain rather simple neural circuits sharing a skeleton of simple circuits in the mammalian cortex, which have been elaborated by additional circuits in mammals. Such basic similarities at the connective level shared by nuclear and cortical pallium might underlie similar or equivalent functions.

Keywords: cortex; pallium; evolution; actinopterygian; teleost

1. NUCLEAR VERSUS LAMINAR TELECEPHALIC PALLIUM

A distinctive feature of the mammalian telencephalon is the presence of a six-layered pallium or neocortex, which is considered to be responsible for 'higher' functions. Despite several 'intelligent' behaviours, however, the pallium of birds is composed mostly of nuclear masses with only a small laminar structure, the Wulst (Shimizu 2007). Laminar pallium, therefore, does not appear to be a requisite for such functions. Karten (1991) suggested that visual information processing in the bird telencephalon is comparable with that in the mammalian neocortex.

The pallium of teleosts also lacks six-layered cortex (or even layered structures), although behavioural studies show some higher cortical-like functions in teleosts including transitive inference (Demski 1983; Grosenick *et al.* 2007; Ito *et al.* 2007). In the present paper, we survey afferent, efferent and intrinsic connections of the teleost telencephalon in search of pallial circuitries comparable with mammalian neocortex.

2. OVERVIEW OF TELEOST TELECEPHALON

The developmental process of the telencephalon is unique in ray-finned fishes. The developing telencephalon does not show evagination being different from other vertebrates, but instead shows eversion. Accordingly, the telencephalon does not form paired tubular structures that contain the lateral ventricles, but forms two solid hemispheres composed of several nuclear masses (figure 1). The hemispheres are separated by a T-shaped common ventricle and covered with a thin tela chorioidea (figure 1). This unique developmental process might explain the formation of the

non-laminar telencephalon in ray-finned fishes. Eversion significantly reduces the pial surface area on the lateral wall of the pallium (figure 1a), leaving little room for cortex formation. Recently, Striedter & Northcutt (2006) have proposed that the eversion evolved because embryos of ray-finned fishes are so small that the telencephalon cannot evaginate, but instead squeeze into a rostral narrow space of the cranium.

The areas dorsalis telencephali (D) and ventralis telencephali (V) named by Nieuwenhuys (1963) are considered to be homologous approximately to the pallium and subpallium of other vertebrates, respectively (Alunni *et al.* 2004; Kage *et al.* 2004; figure 1). The D can be subdivided into several parts: pars centralis (Dc); pars dorsalis (Dd); pars medialis (Dm); pars lateralis (Dl); and pars posterior (Dp). The D also includes the nucleus taenia (nT). These parts may be subdivided further. For example, the Dl is composed at least of two components: dorsal (dDl) and ventral (vDl) regions (Yamane *et al.* 1996). Similarly, the V is composed of pars dorsalis (Vd), pars ventralis (Vv), pars lateralis (Vl), pars supracommissuralis (Vs), pars post-commissuralis (Vp) and so on. In the present survey, we mainly focus on the area dorsalis, the presumed pallial homologue, which we hypothesize to contain functional equivalents to the mammalian neocortex.

3. SENSORY PATHWAYS TO THE TELECEPHALON

Although the telencephalon of fishes has been regarded purely as an olfactory centre for a long time (e.g. Herrick 1924), non-olfactory sensory pathways to the telencephalon have been reported in some teleosts (Ebbesson 1980; Finger 1980; Echterler & Saidel 1981). Based upon more detailed experiments and observations, it has been clarified that different sensory modalities reach different pallial targets (figure 2a; for review see, Yamamoto *et al.* 2007). Secondary olfactory fibres reach the Dp and nT (Yamamoto & Ito 2000). Visual information reaches mainly the dDl mediated by the nucleus prethalamicus (Ito & Vanegas 1983) or a presumed prethalamus homologue within the preglomerular complex, a collection of nuclei in the ventrolateral diencephalon (Northcutt 2006; Yamamoto & Ito 2008). Octavolateral as well as somatosensory information reaches the Dm, Dc and Dl mediated by the ventromedial thalamic nucleus (Ito *et al.* 1986) and the preglomerular complex (Murakami *et al.* 1986a,b; Striedter 1991; Yamamoto & Ito 2005a,b, 2008; Northcutt 2006). Gustatory information reaches mainly the Dm relayed by a portion of the preglomerular complex or its homologue (Kanwal *et al.* 1988; Yoshimoto *et al.* 1998), reaching a region presumably distinct from octavolateral terminal zones within the Dm (Yamamoto & Ito 2005b). Ascending pathways mediated by the preglomerular complex enumerated above exhibit a considerable degree of modality-specific organization (Yamamoto *et al.* 2007; Yamamoto & Ito 2008), similarly to mammalian thalamo-cortical pathways.

It is a critical issue, whether or not the preglomerular complex corresponds to the dorsal thalamus. Northcutt and his co-workers regard the complex as a migrated part of the posterior tuberculum, a non-thalamic diencephalic basal plate derivative, based




Figure 1. (a) Schematic of telencephalic development of actinopterygians (teleost as an example) in comparison with other vertebrates and (b) (i) a frontal section of the telencephalon of tilapia *Oreochromis niloticus* stained by cresyl violet and (ii) line drawing indicating telencephalic structures. Figure 1a adapted from Yamamoto *et al.* (2007). Thick broken lines in (a) and large dots in (b) indicate the boundary between the areas dorsalis (D: presumed pallial homologue) and ventralis (V: presumed subpallial homologue). ca, commissura anterior; dDc, dorsal region of Dc; dDm, dorsal region of Dm; dDml, lateral portion of dDm; dDmm, medial portion of dDm; DPa, dorsal pallium; LPa, lateral pallium; MPa, medial pallium; POA, area preoptica; Se, septum; St, striatum; vDc, ventral region of Dc; vDm, ventral region of Dm; VPa, ventral pallium. See text for other abbreviations. Scale bar, 1 mm.

on developmental analyses with non-experimental materials (Braford & Northcutt 1983; Northcutt 2008). However, analyses of *Pax6* expression during ontogeny suggest that preglomerular neurons are derived from diencephalic alar plate (Kage *et al.* 2004; Ishikawa *et al.* 2007). We therefore consider it likely that preglomerulo-pallial pathways correspond to dorsal thalamo-neocortical pathways in mammals.

4. DESCENDING PATHWAYS FROM THE TELENCEPHALON

Descending pathways originate mainly from the Dc (figure 2b), which are mostly directed to the optic tectum, torus semicircularis, nucleus lateralis valvulae, nucleus paracommissuralis and inferior lobes of the hypothalamus (Murakami *et al.* 1983; Imura *et al.* 2003; Yang *et al.* 2004, 2007). The Dl




Figure 2. Line drawings illustrating (a) ascending sensory pathways to and (b) descending pathways originating from the area dorsalis telencephali in teleosts, and an example of the input–output pathways involved in telencephalic visual processing that can be deduced from tract-tracing studies in (d) acanthopterygian teleosts (Ito *et al.* 1980; Ito & Vanegas 1983; Murakami *et al.* 1983; Yamane *et al.* 1996; Imura *et al.* 2003), in comparison with (c) the mammalian cortical circuit. Ve, common ventricle; tc, tela chorioidea. Other abbreviations are as in the text and those of figure 1.

and Dm also give rise to descending fibres (figure 2b), which are targeted mainly to the inferior lobes of the hypothalamus and preglomerular complex (Folgueira *et al.* 2004; Yamamoto & Ito 2005b, 2008; Northcutt 2006; Yang *et al.* 2007). Thus, the Dc seems to give rise to longer descending pathways that reach down to the meso-rhombencephalic level than those originating from the Dl and Dm that mostly end in the diencephalon. In addition to neurons that project fibres directly out of the telencephalon, the Dl and Dm appear to contain different populations of neurons that send intra-telencephalic fibres to the Dc (Murakami *et al.* 1983; Yamane *et al.* 1996; Yamamoto & Ito 2005b, 2008; Northcutt 2006).

5. NON-LAMINAR CORTIX

Neurons in the Dc resemble those in layer V of the mammalian neocortex, which give rise to long descending fibres. The Dl and Dm appear to contain neurons similar to those in layer VI in terms of originating rather short descending fibres to the diencephalon. The Dl and Dm also contain neuronal populations that participate in telencephalic intrinsic connections as neurons located in other cortical layers in mammals.

Although more detailed analysis of the fibre connections is required to test such a comparison, it appears quite possible that neuronal populations functionally equivalent to neocortical layers in mammals do not form laminar structures but aggregate to form nuclear masses in teleosts, similar to the situation proposed by Karten (1991) for birds. It requires further analyses to see whether the similar neuronal circuitries and their component neuronal populations are homologous.

All experimental procedures were performed under the official Japanese regulations for research on animals and those of the Animal Care Committee of our institutions.

Dr Yuji Ishikawa of National Institute of Radiological Sciences provided important data and helpful suggestions. This study was supported in part by a grant from Fujiwara Natural History Foundation to N.Y.

Hironobu Ito^{1,*} and Naoyuki Yamamoto²

¹Department of Anatomy and Neurobiology, Nippon Medical School, Tokyo 113-8602, Japan

²Laboratory of Fish Biology, Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8601, Japan

*(hiroito@amber.plala.or.jp)

- Alunni, A., Blin, M., Deschet, K., Bourrat, F., Vernier, P. & Rétaux, S. 2004 Cloning and developmental expression patterns of *Dlx2*, *Lhx7* and *Lhx9* in the medaka fish (*Oryzias latipes*). *Mech. Dev.* **121**, 977–983. (doi:10.1016/j.mod.2004.03.023)
- Braford Jr, M. R. & Northcutt, R. G. 1983 Organization of the diencephalon and pretectum of the ray-finned fishes. In *Fish neurobiology*, vol. 2 (eds R. E. Davis & R. G. Northcutt), pp. 117–163. Ann Arbor, MI: University of Michigan Press.
- Demski, L. 1983 Behavioral effects of electrical stimulation of the brain. In *Fish neurobiology*, vol. 2 (R. E. Davis & R. G. Northcutt), pp. 317–359. Ann Arbor, MI: University of Michigan Press
- Ebbesson, S. O. E. 1980 A visual thalamo-telencephalic pathway in a teleost fish (*Holocentrus rufus*). *Cell Tissue Res.* **213**, 505–508. (doi:10.1007/BF00237895)
- Echteler, S. M. & Saidel, W. M. 1981 Forebrain connections in the goldfish support telencephalic homologies with land vertebrates. *Science* **212**, 683–685. (doi:10.1126/science.6971493)
- Finger, T. E. 1980 Nonolfactory sensory pathway to the telencephalon in a teleost fish. *Science* **210**, 671–673. (doi:10.1126/science.7192013)
- Folgueira, M., Anadón, R. & Yáñez, J. 2004 Experimental study of the connections of the telencephalon in the rainbow trout (*Oncorhynchus mykiss*). II: dorsal area and preoptic region. *J. Comp. Neurol.* **480**, 204–233. (doi:10.1002/cne.20341)
- Grosenick, L., Clement, T. S. & Fernald, R. D. 2007 Fish can infer social rank by observation alone. *Nature* **445**, 429–432. (doi:10.1038/nature05511)
- Herrick, C. J. 1924 *Neurological foundations of animal behavior*. New York, NY: Hafner Publishing Company. (Reprinted)
- Imura, K., Yamamoto, N., Sawai, N., Yoshimoto, M., Yang, C.-Y., Xue, H.-G. & Ito, H. 2003 Topographical organization of an indirect telencephalo-cerebellar pathway through the nucleus paracommissuralis in a teleost, *Oreochromis niloticus*. *Brain Behav. Evol.* **61**, 70–90. (doi:10.1159/000069353)
- Ishikawa, Y., Yamamoto, N., Yoshimoto, M. & Ito, H. 2007 Developmental origin of diencephalic sensory relay nuclei in teleosts. *Brain Behav. Evol.* **69**, 87–95. (doi:10.1159/000095197)
- Ito, H. & Vanegas, H. 1983 Cytoarchitecture and ultrastructure of nucleus prethalamicus, with special reference to degenerating afferents from optic tectum and telencephalon, in a teleost (*Holocentrus ascensionis*). *J. Comp. Neurol.* **221**, 401–415. (doi:10.1002/cne.902210404)
- Ito, H., Morita, Y., Sakamoto, N. & Ueda, S. 1980 Possibility of telencephalic visual projections in a teleost, *Holocentrus rufus*. *Brain Res.* **197**, 219–222. (doi:10.1016/0006-8993(80)90448-5)
- Ito, H., Murakami, T., Fukuoka, T. & Kishida, R. 1986 Thalamic fiber connections in a teleost (*Sebastiscus marmoratus*): visual, somatosensory, octavolateral, and cerebellar relay region to the telencephalon. *J. Comp. Neurol.* **250**, 215–227. (doi:10.1002/cne.902500208)
- Ito, H., Ishikawa, Y., Yoshimoto, M. & Yamamoto, N. 2007 Diversity of brain morphology in teleosts: brain and ecological niche. *Brain Behav. Evol.* **69**, 76–86. (doi:10.1159/000095196)
- Kage, T. *et al.* 2004 Morphogenesis and regionalization of the medaka embryonic brain. *J. Comp. Neurol.* **476**, 219–239. (doi:10.1002/cne.20219)
- Kanwal, J. S., Finger, T. E. & Caprio, J. 1988 Forebrain connections of the gustatory system in ictalurid catfishes. *J. Comp. Neurol.* **278**, 353–376. (doi:10.1002/cne.902780306)
- Karten, H. J. 1991 Homology and evolutionary origins of the 'neocortex'. *Brain Behav. Evol.* **38**, 264–272. (doi:10.1159/000114393)
- Murakami, T., Morita, Y. & Ito, H. 1983 Extrinsic and intrinsic fiber connections of the telencephalon in a teleost, *Sebastiscus marmoratus*. *J. Comp. Neurol.* **216**, 115–131. (doi:10.1002/cne.902160202)
- Murakami, T., Fukuoka, T. & Ito, H. 1986a Telencephalic ascending acousticolateral system in a teleost (*Sebastiscus marmoratus*), with special reference to fiber connections of the nucleus preglomerulosus. *J. Comp. Neurol.* **247**, 383–397. (doi:10.1002/cne.902470308)
- Murakami, T., Ito, H. & Morita, Y. 1986b Telencephalic afferent nuclei in the carp diencephalon, with special reference to fiber connections of the nucleus preglomerulosus pars lateralis. *Brain Res.* **382**, 97–103. (doi:10.1016/0006-8993(86)90115-0)
- Nieuwenhuys, R. 1963 The comparative anatomy of the actinopterygian forebrain. *J. Hirnforsch.* **6**, 171–192.
- Northcutt, R. G. 2006 Connections of the lateral and medial divisions of the goldfish telencephalic pallium. *J. Comp. Neurol.* **494**, 903–943. (doi:10.1002/cne.20853)
- Northcutt, R. G. 2008 Forebrain evolution in bony fishes. *Brain Res. Bull.* **75**, 191–205. (doi:10.1016/j.brainresbull.2007.10.058)
- Shimizu, T. 2007 The avian brain revisited: anatomy and evolution of the telencephalon. In *Integration of comparative neuroanatomy and cognition* (eds S. Watanabe & M. H. Hofman), pp. 55–73. Tokyo, Japan: Keio University Publications.
- Striedter, G. F. 1991 Auditory, electrosensory, and mechanosensory lateral line pathways through the forebrain in channel catfishes. *J. Comp. Neurol.* **312**, 311–331. (doi:10.1002/cne.903120213)
- Striedter, G. F. & Northcutt, R. G. 2006 Head size constrains forebrain development and evolution in ray-finned fishes. *Evol. Dev.* **8**, 215–222. (doi:10.1111/j.1525-142X.2006.00091.x)
- Yamamoto, N. & Ito, H. 2000 Afferent sources to the ganglion of the terminal nerve in teleosts. *J. Comp. Neurol.* **428**, 355–375. (doi:10.1002/1096-9861(20001211)428:2<355::AID-CNE12>3.0.CO;2-W)
- Yamamoto, N. & Ito, H. 2005a Fiber connections of the central nucleus of semicircular torus in cyprinids. *J. Comp. Neurol.* **491**, 186–211. (doi:10.1002/cne.20683)
- Yamamoto, N. & Ito, H. 2005b Fiber connections of the anterior preglomerular nucleus in cyprinids with notes on telencephalic connections of the preglomerular complex. *J. Comp. Neurol.* **491**, 212–233. (doi:10.1002/cne.20681)
- Yamamoto, N. & Ito, H. 2008 Visual, lateral line, and auditory ascending pathways to the dorsal telencephalic area through the rostralateral region of lateral preglomerular nucleus in cyprinids. *J. Comp. Neurol.* **508**, 615–647. (doi:10.1002/cne.21717)
- Yamamoto, N., Ishikawa, Y., Yoshimoto, M., Xue, H.-G., Bahaxar, N., Sawai, N., Yang, C.-Y., Ozawa, H. & Ito, H. 2007 A new interpretation on the homology of the teleostean telencephalon based on hodology and a new eversion model. *Brain Behav. Evol.* **69**, 96–104. (doi:10.1159/000095198)
- Yamane, Y., Yoshimoto, M. & Ito, H. 1996 Area dorsalis pars lateralis of the telencephalon in a teleost (*Sebastiscus marmoratus*) can be divided into dorsal and ventral regions. *Brain Behav. Evol.* **48**, 338–349. (doi:10.1159/000113212)

- Yang, C.-Y., Yoshimoto, M., Xue, H.-G., Yamamoto, N., Imura, K., Sawai, N., Ishikawa, Y. & Ito, H. 2004 Fiber connections of the lateral valvular nucleus in a percomorph teleost, tilapia (*Oreochromis niloticus*). *J. Comp. Neurol.* **474**, 209–226. (doi:10.1002/cne.20150)
- Yang, C.-Y., Xue, H.-G., Yoshimoto, M., Ito, H., Yamamoto, N. & Ozawa, H. 2007 Fiber connections of the corpus glomerulosum pars rotunda, with special reference to efferent projection pattern to the inferior lobe in a percomorph teleost, tilapia (*Oreochromis niloticus*). *J. Comp. Neurol.* **501**, 582–607. (doi:10.1002/cne.21261)
- Yoshimoto, M., Albert, J. S., Sawai, N., Shimizu, M., Yamamoto, N. & Ito, H. 1998 Telencephalic ascending gustatory system in a cichlid fish, *Oreochromis (Tilapia) niloticus*. *J. Comp. Neurol.* **392**, 209–226. (doi:10.1002/(SICI)1096-9861(19980309)392:2<209::AID-CNE5>3.0.CO;2-6)