

Potential impacts of climate change on the winter distribution of Afro-Palaearctic migrant passerines

Morgane Barbet-Massin, Bruno A Walther, Wilfried Thuiller, Carsten Rahbek and Frédéric Jiguet

Biol. Lett. 2009 **5**, 248-251 first published online 25 February 2009
doi: 10.1098/rsbl.2008.0715

Supplementary data

["Data Supplement"](#)

<http://rsbl.royalsocietypublishing.org/content/suppl/2009/02/20/rsbl.2008.0715.DC1.html>

References

[This article cites 15 articles, 2 of which can be accessed free](#)

<http://rsbl.royalsocietypublishing.org/content/5/2/248.full.html#ref-list-1>

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (466 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Potential impacts of climate change on the winter distribution of Afro-Palaeartic migrant passerines

Morgane Barbet-Massin^{1,*}, Bruno A. Walther^{2,3,4}, Wilfried Thuiller⁵, Carsten Rahbek³ and Frédéric Jiguet¹

¹CRBPO, UMR 5173 MNHN-CNRS-UPMC, CP51, 55 Rue Buffon, 75005 Paris, France

²DIVERSITAS, 57 rue Cuvier, CP 41, 75231 Paris Cedex 05, France

³Zoological Museum, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark

⁴Centre of Excellence for Invasion Biology (CIB), University of Stellenbosch, Private Bag X1, Matieland, South Africa

⁵Laboratoire d'Ecologie Alpine, UMR CNRS 5553, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 9, France

*Author for correspondence (barbet@mnhn.fr).

We modelled the present and future sub-Saharan winter distributions of 64 trans-Saharan migrant passerines to predict the potential impacts of climate change. These predictions used the recent ensemble modelling developments and the latest IPCC climatic simulations to account for possible methodological uncertainties. Results suggest that 37 species would face a range reduction by 2100 (16 of these by more than 50%); however, the median range size variation is –13 per cent (from –97 to +980%) under a full dispersal hypothesis. Range centroids were predicted to shift by 500 ± 373 km. Predicted changes in range size and location were spatially structured, with species that winter in southern and eastern Africa facing larger range contractions and shifts. Predicted changes in regional species richness for these long-distance migrants are increases just south of the Sahara and on the Arabian Peninsula and major decreases in southern and eastern Africa.

Keywords: Africa; birds; climate change; ensemble forecast; climate suitability model; species' range shift

1. INTRODUCTION

Climate suitability models have recently received a great deal of attention as the methods of choice to estimate the potential risks that biodiversity faces due to global warming (Pearson & Dawson 2003). Although these models are based on a general conceptual framework, the numerous modelling techniques used to predict climatic scenarios, climate suitability and their resultant uncertainties represent major sources of prediction variability. Nevertheless, more robust forecasts can be achieved if ensemble forecasts (use of multiple models) are produced and analysed appropriately (Araújo *et al.* 2005).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2008.0715> or via <http://rsbl.royalsocietypublishing.org>.

Breeding bird populations have been declining in Europe during the past few decades, with long-distance migrants being especially affected (Sanderson *et al.* 2006). Few studies have investigated the impact on migrant species due to changing climate conditions along migration routes on stopover sites and wintering grounds (Tøttrup *et al.* 2008). The potential impacts of climate change on the breeding ranges of European birds have been estimated recently (Huntley *et al.* 2008), and changes in migration phenology have been reported in numerous species (Jonzén *et al.* 2006). However, we still need to study the potential impacts on winter ranges. La Sorte & Thompson (2007) reported recent poleward shifts in the winter ranges of North American birds. It is thus likely that a changing climate in Africa will affect the winter distribution of trans-Saharan migrants, potentially affecting their conservation status.

Here, we estimated the potential impacts of climate change on the winter distributions of 64 species of Afro-Palaeartic trans-Saharan migrant passerines. We used various climatic scenarios (IPCC 2007) in an ensemble forecasting framework (Araújo & New 2007) to account for climate prediction and modelling uncertainties. We first explored the potential changes in range size and position, considering either full or no dispersal ability of species between present and future ranges. We then synthesized potential impacts by mapping predicted changes in species richness of these migrant species.

2. MATERIAL AND METHODS

(a) Bird and climate data

For each migrant species, we used recorded localities obtained from museum specimens, ringing records and field observations (see the project website www.macroecology.ku.dk/africamigrants). We considered records obtained south of 24° N latitude (south of the Sahara) from 1st November to 29th February, to be sure they concerned wintering not migrating birds. A total of 16 638 primary records were used, reduced to 9797 records when combined at a $0.5^\circ \times 0.5^\circ$ grid cell resolution, leaving 6–1201 grid cell records per species. Pseudo-absences were selected randomly, outside the suitable area of the surface range envelope model (appendix 2, electronic supplementary material).

Future climate projections were derived from five general circulation models (GCMs; BCM2, ECHAM5, HADCM3, MIR-OHIC3_2-HI and MK3) and three recent special reports on emission scenarios (SRES) (A1B, B1 and A2 when available), using monthly mean predictions taken from IPCC (2007), for time intervals 1961–1990, 2011–2030, 2046–2065 and 2080–2099. GCMs represent physical processes in the atmosphere, ocean, cryosphere and land surface. SRES reflect the potential impacts of different assumptions about demographic, socio-economic and technological development on the release of greenhouse gases. We interpolated these values onto a $0.5^\circ \times 0.5^\circ$ grid over Africa and the Arabian Peninsula for a total of 11 275 grid cells. For distribution modelling, we used eight climatic variables: (i) annual mean temperature, (ii) mean temperature of the warmest month, (iii) mean temperature of the coldest month, (iv) temperature seasonality, (v) annual precipitation, (vi) precipitation of the wettest month, (vii) precipitation of the driest month, and (viii) precipitation seasonality. The use of climatic variables only assumes that current range limits are mainly driven by climate, which is a reasonable hypothesis at a continental scale (Pearson & Dawson 2003). However, in some cases, climate variables could simply be proxies for the major habitats and these habitats may not be able to track the climate change.

(b) Ensemble forecast

We modelled present and possible future distributions using nine climate suitability models (Thuiller 2003; Phillips *et al.* 2006; details are given in appendix 1 of the electronic supplementary material). Each model was first calibrated using a 70 per cent random subset of the data, and then evaluated with the remaining 30 per cent using a threshold-independent method, the area under

the relative operating characteristic curve (AUC). Data splitting was performed five times, from which we calculated an average AUC. The final runs of the models for obtaining the range predictions used 100 per cent of the data.

For each species and time period, we obtained 45 (9 models \times 5 GCMs) present and 108 (9 models \times 12 (GCMs \times SRES)) future modelled distributions. The consensus distribution was obtained with an ensemble forecast approach, by selecting the outputs of the five modelling techniques with the best AUC scores, and by further calculating the unweighted mean distributions (Marmion *et al.* 2009; appendix 1, electronic supplementary material) for the corresponding 25 (5 models \times 5 GCMs) present or 60 (5 models \times 12 (GCMs \times SRES)) future distributions. For every species, the mean AUC of all models ranged from 0.757 to 0.987 (mean = 0.930 ± 0.04). The mean AUC of the models used for consensus distributions ranged from 0.840 to 0.996 (mean = 0.962 ± 0.03 ; appendix 3, electronic supplementary material). To transform the probabilistic consensus distribution to a presence/absence distribution, we preserved the occurrence probabilities for pixels above the sensitivity–specificity sum maximization threshold, and set to zero the occurrence probability for pixels under the threshold.

(c) Winter range extents and shifts

The range was calculated for each species' consensus distribution, by summing the probabilities of the filtered distribution, weighting each pixel by its area and accounting for the latitudinal variations. The latitude and longitude of the range centroids were also estimated from the filtered distribution, using probabilities as weights. For each species, the distance between present and future range centroids was calculated as a measure of range shift. We further calculated the area ratio between future and present ranges as a measure of the potential impact on range size; this measure assumes that each species is capable of dispersing completely from the present to the future range. We finally calculated the ratio between the overlap of present and future ranges and the present range; this measure assumes that no species is capable of any dispersal into new areas predicted to be climatically suitable in the future. We used a generalized least-squares phylogenetic comparative method (appendix 6, electronic supplementary material) to test for a correlation between the ratio of range sizes, the range overlap or the range shift with the present range size and the latitude and longitude of range centroid. To map the potential evolution of species richness of these migrant birds, we considered the extreme two cases presented above where dispersal is either full or null. Given that all the investigated bird species are migratory, the full dispersal ability is certainly closer to the reality. Species richness per grid cell was obtained by summing the number of species predicted as present.

3. RESULTS

Under the total dispersal hypothesis, we found that 35, 38 and 37 species out of a total of 64 should undergo a reduction in their winter range by 2030, 2065 and 2100, respectively (appendices 4 and 7, electronic supplementary material). There is an increase in the variation of winter range size over time: only one species is predicted to experience a more than 50 per cent reduction by 2030, whereas 11 and 16 species should face such a reduction by 2065 and 2100, respectively. By 2100, there should be a median variation (difference between future and present ranges divided by the present range) of -13 per cent in range sizes, with a maximum increase of 980 per cent for *Locustella naevia* and a maximum decrease of 97 per cent for *Ficedula albicollis* and *Ficedula semitorquata* (appendix 8, electronic supplementary material). Under the null dispersal hypothesis, the median variation in winter range size should be a 28 per cent decrease by 2100.

By 2100, the range centroids are predicted to have shifted by 500 ± 373 km (range 57–1410; figure 1). Most species whose winter range is predicted to decrease are predicted to shift latitudinally, whereas most species whose winter range is predicted to




Figure 1. Predicted range shifts in winter ranges of 64 long-distance Afro-Palaeartic passerine birds under future climatic conditions, predicted for 2100 by an ensemble forecast framework. The distance and direction of each species' range centroid displacement is illustrated with an arrow. We contrasted species whose range is predicted to increase (black) or decrease (grey) in size.

increase are predicted to shift longitudinally. Thus, these predicted shifts are strongly spatially structured, and they depend upon whether the winter range is predicted to decrease or increase (see appendix 6 in the electronic supplementary material and table 1). In generalized least-squares models, we found a negative effect of longitude on the ratio of range size, on range overlap and on range shift, and a positive effect of latitude on range overlap. With spatial effects accounted for, we found a positive effect of present range size on range overlap and a negative effect on range ratio and range shift.

4. DISCUSSION

By 2100, 37 out of 64 long-distance migrant passerines are predicted to suffer a winter range reduction, even assuming that species are capable of full dispersal. Under such conditions, 16 species would lose more than half of their range size, two of them losing almost all suitable wintering conditions. The predicted variations in distributions are highly spatially structured; we predict larger distributional reductions for species wintering in eastern or southern Africa, while those wintering in western Africa are predicted to extend their ranges eastwards. In agreement with previous studies (La Sorte & Thompson 2007; Huntley *et al.* 2008), we predict mostly poleward distributional shifts for species presently wintering in the Southern Hemisphere; however, species presently wintering in West Africa would shift eastward since they are limited northward by the Sahara. Large potential impacts of climate change are predicted for range-restricted species with limited dispersal ability, as they face the smallest overlap between present and future predicted ranges and the largest shifts.

When looking at overall species richness, predicted changes by 2100 (figure 2) show some similarities to those predicted for mammals (Thuiller *et al.* 2006),

Table 1. Results of generalized least-squares models conducted on the predicted ratio of range sizes, the predicted range overlap (both log transformed) and the shift, explained by latitude, longitude and size of the predicted present range (effects of predictors are adjusted to each other).

predictor	ratio of range sizes		range overlap		range shift	
	t_{60}	p -value	t_{60}	p -value	t_{60}	p -value
latitude	1.21	0.23	6.07	<0.001	−1.29	0.201
longitude	−4.19	<0.001	−4.23	<0.001	−2.84	0.006
present range size	−3.36	0.001	3.66	<0.001	−3.09	0.003




Figure 2. Predicted species richness of migrant passerines in Africa under future climatic scenarios (by 2100): (a) future predicted species richness under the hypothesis of full dispersal ability between predicted present and future ranges; (b) present predicted species richness (1990); (c) future predicted species richness under the hypothesis of null dispersal ability; (d, e) present variations in species richness corresponding to (a) and (c), respectively.

especially a large decrease in the western parts of southern Africa, although they are clearly different from those obtained for African breeding birds (Jetz *et al.* 2007). Presently, Afro-Palaearctic migrant passerines winter preferentially in a belt stretching from Senegal to Ethiopia and from Sudan southwards through Kenya and Tanzania (Wisz *et al.* 2007). This distribution corresponds largely with their preferred dry Acacia and moist broadleaved savannah habitats (Salewski & Jones 2006). Consequently, low species-rich areas are found in rainforest habitats at equatorial latitudes and in desert-like habitats, e.g. most of

Namibia. Assuming full dispersal ability, a reasonable assumption for migratory birds, our distribution models predict large decreases in species richness in the initially species-rich East African region. The presently species-poor western part of southern Africa is predicted to lose even more of its species. Under the hypothesis of null dispersal, the charts did not show any major relative differences in patterns. Finally, our simulations do not predict present sub-Saharan wintering climatic conditions to appear north of the Sahara, making it unlikely that sub-Saharan migrant species could substitute habitat loss south of the

Sahara with expansions of distributions north of the Sahara. Further studies of long-distance migrants should also consider the effects of predicted changes in land cover on future distributions.

- Araújo, M. B. & New, M. 2007 Ensemble forecasting of species distributions. *Trends Ecol. Evol.* **22**, 42–47. (doi:10.1016/j.tree.2006.09.010)
- Araújo, M. B., Whittaker, R. J., Ladle, R. J. & Erhard, M. 2005 Reducing uncertainty in projections of extinction risk from climate change. *Glob. Ecol. Biogeogr.* **14**, 529–538. (doi:10.1111/j.1466-822X.2005.00182.x)
- Huntley, B., Collingham, Y. C., Willis, S. G. & Green, R. E. 2008 Potential impacts of climatic change on European breeding birds. *PLoS One* **3**, e1439. (doi:10.1371/journal.pone.0001439)
- IPCC 2007 Climate change 2007: IPCC fourth assessment report. Cambridge University Press, Cambridge, UK.
- Jetz, W., Wilcove, D. S. & Dobson, A. P. 2007 Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biol.* **5**, e157. (doi:10.1371/journal.pbio.0050157)
- Jonzén, N. *et al.* 2006 Rapid advance of spring arrival dates in long-distance migratory birds. *Science* **312**, 1959–1961. (doi:10.1126/science.1126119)
- La Sorte, F. A. & Thompson, F. R. 2007 Poleward shifts in winter ranges of North American birds. *Ecology* **88**, 1803–1812. (doi:10.1890/06-1072.1)
- Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R. K. & Thuiller, W. 2009 Evaluation of consensus methods in predictive species distribution modelling. *Divers. Distrib.* **15**, 59–69. (doi:10.1111/j.1472-4642.2008.00491.x)
- Pearson, R. G. & Dawson, T. P. 2003 Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Glob. Ecol. Biogeogr.* **12**, 361–371. (doi:10.1046/j.1466-822X.2003.00042.x)
- Phillips, S. J., Anderson, R. P. & Schapire, R. E. 2006 Maximum entropy modeling of species geographic distributions. *Ecol. Model.* **190**, 231–259. (doi:10.1016/j.ecolmodel.2005.03.026)
- Salewski, V. & Jones, P. 2006 Palearctic passerines in Afrotropical environments: a review. *J. Ornithol.* **147**, 192–201. (doi:10.1007/s10336-006-0057-5)
- Sanderson, F. J., Donald, P. F., Pain, D. J., Burfield, I. J. & van Bommel, F. P. J. 2006 Long-term population declines in Afro-Palearctic migrant birds. *Biol. Conserv.* **131**, 93–105. (doi:10.1016/j.biocon.2006.02.008)
- Thuiller, W. 2003 BIOMOD—optimizing predictions of species distributions and projecting potential future shifts under global change. *Glob. Chang. Biol.* **9**, 1353–1362. (doi:10.1046/j.1365-2486.2003.00666.x)
- Thuiller, W., Broennimann, O., Hughes, G. O., Alkemade, J. R. M., Midgley, G. F. & Corsi, F. 2006 Vulnerability of African mammals to anthropogenic climate change under conservative land transformation assumptions. *Glob. Chang. Biol.* **12**, 424–440. (doi:10.1111/j.1365-2486.2006.01115.x)
- Tøttrup, A. P., Thorup, K., Rainio, K., Yosef, R., Lehtikoinen, E. & Rahbek, C. 2008 Avian migrants adjust migration in response to environmental conditions en route. *Biol. Lett.* **4**, 685–688. (doi:10.1098/rsbl.2008.0290)
- Wisn, M. S., Walther, B. A. & Rahbek, C. 2007 Using potential distributions to explore determinants of western Palearctic migratory songbird species richness in sub-Saharan Africa. *J. Biogeogr.* **34**, 828–841. (doi:10.1111/j.1365-2699.2006.01661.x)