

Relationships between population size and pollen fates in a moth-pollinated orchid

Steven D Johnson, Erica Torninger and Jon Ågren

Biol. Lett. 2009 **5**, 282-285 first published online 25 February 2009
doi: 10.1098/rsbl.2008.0702

Supplementary data

["Data Supplement"](#)

<http://rsbl.royalsocietypublishing.org/content/suppl/2009/02/20/rsbl.2008.0702.DC1.html>

References

[This article cites 20 articles](#)

<http://rsbl.royalsocietypublishing.org/content/5/2/282.full.html#ref-list-1>

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (466 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Relationships between population size and pollen fates in a moth-pollinated orchid

Steven D. Johnson^{1,*}, Erica Torninger^{1,2} and Jon Ågren²

¹School of Biological and Conservation Sciences, University of KwaZulu-Natal, Scottsville, Pietermaritzburg 3209, South Africa

²Department of Ecology and Evolution, Evolutionary Biology Centre, Uppsala University, 752 36 Uppsala, Sweden

*Author for correspondence (johnsonsd@ukzn.ac.za).

Management of small plant populations requires an understanding of their reproductive ecology, particularly in terms of sensitivity to Allee effects. To address this issue, we explored how components of pollen transfer and pollination success of individual plants varied among 36 populations of the self-compatible moth-pollinated orchid *Satyrium longicauda* in South Africa. Mean fruit set, seed production, proportion of flowers with pollen deposited or removed and proportion of removed pollen that reached stigmas (approx. 8% in this species) were not significantly related to population size (range: 1–450 flowering individuals), density or isolation. Plants in small populations did, however, have significantly higher levels of pollinator-mediated self-pollination (determined using colour-labelled pollen) than those in larger populations. Our results suggest that small populations of this orchid species are resilient to Allee effects in terms of overall pollination success, although the higher levels of pollinator-mediated self-pollination in small populations may lead to inbreeding depression and long-term erosion of genetic diversity.

Keywords: Allee effects; pollen transfer efficiency; pollination; population ecology; *Satyrium*; self-pollination

1. INTRODUCTION

Plants in small or sparse populations often experience lower reproductive output than their conspecifics in large populations (e.g. Lamont *et al.* 1993; Ågren 1996; Kéry *et al.* 2000; Ågren *et al.* 2008). This phenomenon, known as the Allee effect, is of great concern for conservationists as reduction in population size is a common consequence of habitat fragmentation (Eriksson & Ehrlén 2001). However, the actual extent of Allee effects in plants may be obscured by a bias towards publication of studies that find these effects. In addition, the underlying causes of Allee effects when they are present in plant populations remain poorly understood (Hobbs & Yates 2003; Ghazoul 2005).

Positive relationships between population size and seed production in plants may be due to ecological or genetic factors or a combination of both. Some studies have shown that lower seed set of plants in small populations is due to pollen limitation (Ågren 1996; Ward & Johnson 2005). This is usually an ecological consequence of inadequate pollinator visitation affecting pollen quantity, but there is also evidence that the quality of pollen (e.g. compatibility type) received by stigmas can be lower in small populations (Busch & Schoen 2008). Other evidence for a role for genetic factors in Allee effects includes correlations between inbreeding coefficients and low seed production in small populations (reviewed by Leimu *et al.* 2006). Increased homozygosity in small populations can be due to loss of alleles through drift, founder effects, biparental inbreeding or higher rates of self-fertilization (cf. Klinkhamer & de Jong 1990; Ellstrand & Elam 1993; Young *et al.* 1996).

We investigated the relationships between population size and components of pollination success in *Satyrium longicauda*, a widespread South African terrestrial orchid pollinated by hawkmoths (Harder & Johnson 2005; Jersakova & Johnson 2007). As study organisms, orchids have the advantage of possessing flowers in which both pollen removal and pollen deposition can be quantified precisely in the field (Nilsson 1992). In addition, pollen can easily be stained and tracked in order to determine the rates of pollinator-mediated self-pollination and other pollen fates (cf. Peakall 1989; Johnson *et al.* 2004, 2005). This is the first study in which the relationship between pollen fates and plant population size has been explored in any detail.

The aims of this study were to establish whether the size of *S. longicauda* populations influences pollination success, pollen transfer efficiency (PTE), rate of self-pollination and seed production of individual plants.

2. MATERIAL AND METHODS

We studied 36 discrete natural populations of *S. longicauda* in January and February 2003. These were distributed across six grassland subregions (five to seven populations each) close to Pietermaritzburg, South Africa (see the electronic supplementary material). The orchids were the only nectar plants available for hawkmoths at these sites, thus making it unlikely that interspecific interactions would affect pollination. Population size varied from 1 to 450 flowering individuals (median = 11). For each population, we recorded population density, calculated as the number of plants per square metre (range: 0.0001–1.0, median = 0.013) and degree of isolation, calculated as the distance in metres to the nearest population (range: 50–5000, median = 300).

A preliminary breeding system experiment at the 'Wahroonga' site revealed that *S. longicauda* is self-compatible and largely incapable of autonomous self-pollination. Eight randomly chosen plants were bagged to prevent pollinators from visiting them. After anthesis, four flowers per plant were randomly selected to be self-pollinated, four to be cross-pollinated and four to be unmanipulated controls. The median percentage fruit set per plant for self-pollinated flowers (100%, range 50–100%) did not differ significantly from that for cross-pollinated flowers (100%, range 66.6–100%, $Z = 0.92$, $p = 0.35$, Wilcoxon signed-rank test for paired samples). However, the mean (\pm s.e.) angular-transformed percentage of seeds with viable embryos per fruit per plant was much higher in the cross-pollination treatment (back-transformed values, cross: $31.9 \pm 5.5\%$, self: $8.0 \pm 2.9\%$, $t_7 = 4.1$, $p = 0.004$, paired t -test). Just 21 per cent of flowers in the unmanipulated treatment set fruit and these contained a minimal mean percentage (2.4%) of seeds with embryos.

Natural pollination success was scored for one recently wilted flower selected arbitrarily from each of approximately 30 plants per population. In populations containing fewer than 10 individuals,

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2008.0702> or via <http://rsbl.royalsocietypublishing.org>.

Received 26 November 2008

Accepted 13 January 2009




Figure 1. (a,b) Relationships between population size and measures of pollination success, (c) PTE, (d) proportion of self-pollination ($R^2_{\text{adj}} = 0.38$; subregion NS; log population size, $p = 0.004$) and (e,f) fruit and seed production in *S. longicauda*. Symbols indicate population clusters belonging to a particular subregion of the study area (treated as a block factor in the ANCOVA; table 1).

two to four flowers per plant were collected. Collected flowers were examined under a microscope to establish the number of pollinaria (pollen packages) removed and the number of individual pollen massulae (subunits of pollinaria) deposited on stigmas.

For each population, the PTE, a measure of the proportion of removed pollen that reaches stigmas (cf. Johnson *et al.* 2004), was calculated as $M_s/(N_r \times P)$, where M_s is the mean number of pollen massulae on stigmas; N_r is the mean number of pollinaria removed from flowers; and P is the mean number of pollen massulae per pollinarium. The latter was based on counts for 20–30 pollinaria from each of seven populations.

Pollinaria in 5–15 plants in each of 21 populations were stained with histochemicals (Peakall 1989) to assess the rate of self-pollination and its relationship to population size. Up to three focal groups of five plants each were used in each population and each plant in a focal group was assigned to be stained with one of the following stains: gentian violet; fast green; brilliant blue; rhodamine; and neutral red. These stains do not affect transfer probabilities (Johnson *et al.* 2004). Staining was accomplished by injecting the histochemicals into the anthers of up to five flowers per plant with a 5 μ l Hamilton microsyringe. After 3 days, stained plants were collected and the number of stained pollinaria removed and stained massulae deposited were recorded. The rate of self-pollination was calculated as P_s/PTE , where P_s is the proportion of removed stained pollen that was deposited on stigmas of the source plant and PTE is the overall proportion of removed pollen that was deposited on stigmas in the population (see above).

At fruit maturation, the percentage of flowers that set fruit was determined for plants in 17 populations. We scored fruit set in all the plants in these populations, except for a larger population in

which it was scored for a subsample of 20 individuals. We also collected a single fruit from each of 1–18 plants (mean = 6.9) in 24 populations and then, using a dissecting microscope, established the percentage of ovules that had developed into seeds with embryos in each of these fruits.

We examined the relationship between population size and population means for various measures of plant reproductive success using two sets of analysis of covariance (ANCOVA) models. The first set simultaneously included population size and grassland subregion as independent variables, while the second set in addition included population density and isolation as independent variables.

3. RESULTS

We did not detect significant relationships between population size and measures of male and female pollination success in *S. longicauda* (figure 1; table 1). While PTE was apparently unaffected by population size, self-pollination was negatively related to population size in ANCOVA models both that did ($F_{1,9} = 8.61$, $p = 0.017$; table 1) and did not take the effects of population density and isolation into account ($F_{1,12} = 12.6$, $p = 0.004$; figure 1; statistically significant also after sequential Bonferroni correction). For both sets of models, neither the proportion of flowers setting fruit or the percentage of seeds with

Table 1. ANCOVA models of the relationships between four predictor variables—geographical subregion, population size, population density and population isolation—and population means for measures of pollination success, PTE and seed production in populations of *S. longicauda*. Values given are *F* ratios, except for sample size (*n*) that refers to the number of populations included in the model. Population size, density and isolation were log transformed prior to analysis. Response variables were angular-transformed proportions. (**p* < 0.05, ***p* < 0.01, ****p* < 0.001.)

| response variable | <i>n</i> | subregion | pop size | pop density | pop isolation |
|-----------------------------|----------|-----------|----------|-------------|---------------|
| pollinated flowers | 29 | 8.38*** | 0.50 | 0.07 | 0.41 |
| flowers with pollen removed | 28 | 3.32* | 0.24 | 0.12 | 0.07 |
| pollen transfer efficiency | 29 | 5.15** | 1.80 | 0.81 | 1.18 |
| self-pollination | 18 | 1.51 | 8.61* | 3.20 | 0.79 |
| fruit production | 17 | 0.46 | 2.15 | 0.13 | 0.88 |
| seeds with embryos | 23 | 2.88 | 2.90 | 0.29 | 0.32 |

viable embryos showed a significant relationship with population size. No measure of pollination success or fecundity in *S. longicauda* was significantly related to population density or isolation (table 1).

The variance in the examined measures of reproductive success tended to decrease with increasing population size, but except for PTE (regression of absolute residuals from ANCOVA in table 1 on log population size, $b = -0.025$, $p = 0.028$, $R^2_{\text{adj}} = 0.14$), this trend was not statistically significant ($p > 0.33$).

4. DISCUSSION

The orchid *S. longicauda* appears to have an overall resilience to Allee effects in terms of both population size and density. Both small and large populations experience high levels of pollination success, with approximately 70 per cent of flowers having pollen deposited and removed. In terms of PTE, approximately 8 per cent of the pollen removed from flowers reached stigmas, irrespective of population size. Pollen transfer is a notoriously stochastic process (cf. Harder & Johnson 2005) and this could confound the detection of population effects on PTE. However, the proportion of deposited pollen that arose from pollinator-mediated self-pollination was higher in small populations (figure 1; table 1). This is consistent with the theoretical expectation that pollinators should visit more flowers per plant in small or sparse populations, which, in turn, would increase the level of geitonogamous self-pollination (cf. Klinkhamer & de Jong 1990; Goulson 1999).

Although higher levels of self-pollination could potentially result in lower seed production (owing to significant inbreeding depression at the seed development stage, see §2), an effect of population size on fruit and seed production of *S. longicauda* was not evident in this study (figure 1; table 1). A possible explanation for this is that, despite the increased level of self-pollination in small populations, plants in these populations received sufficient cross-pollen for high levels of seed production. Orchids such as *S. longicauda* that have sectile pollinia usually receive mixtures of self- and cross-pollen on their stigmas (Johnson *et al.* 2005). Waites & Ågren (2004) found that variation in seed production of the self-incompatible *Lythrum salicaria* largely reflected variation in deposition of compatible pollen and that seed production was apparently unaffected by the

deposition of large quantities of self-incompatible pollen grains.

Many of the studies, in which a positive effect of population size on plant fecundity has been found, have involved bee-pollinated species (cf. Ågren 1996; Kéry *et al.* 2000). Bees often forage in a local range and are expected to avoid small unprofitable patches (Goulson 1999). It may be that the resilience of *S. longicauda* to Allee effects is partially related to its hawkmoth pollinators, which are migratory animals and largely opportunistic foragers. However, any firm conclusions about the link between Allee effects and particular pollinators must await a wider range of studies.

In conclusion, our study shows that seed production and most pollen fates in *S. longicauda* are not affected by the size, density or isolation of populations. However, the increased level of geitonogamous self-pollination in small populations is an intriguing phenomenon that deserves more attention, both with respect to the underlying process and its possible implications for the genetic structure of populations.

This study was supported by a joint NRF-SIDA grant to S.D.J. and J.Å.

- Ågren, J. 1996 Population size, pollinator limitation, and seed set in the self-incompatible herb *Lythrum salicaria*. *Ecology* **77**, 1779–1790. (doi:10.2307/2265783)
- Ågren, J., Ehrlén, J. & Solbreck, C. 2008 Spatio-temporal variation in fruit production and seed predation in a perennial herb influenced by habitat quality and population size. *J. Ecol.* **96**, 334–345. (doi:10.1111/j.1365-2745.2007.01334.x)
- Busch, J. W. & Schoen, D. J. 2008 The evolution of self-incompatibility when mates are limiting. *Trends Plant Sci.* **13**, 128–136. (doi:10.1016/j.tplants.2008.01.002)
- Ellstrand, N. C. & Elam, D. R. 1993 Population genetic consequences of small population size: implications for plant conservation. *Annu. Rev. Ecol. Syst.* **24**, 217–242. (doi:10.1146/annurev.es.24.110193.001245)
- Eriksson, O. & Ehrlén, J. 2001 Landscape fragmentation and the viability of plant populations. In *Integrating ecology and evolution in a spatial context* (eds J. Silvertown & J. Antonovics), pp. 1–28. Oxford, UK: Blackwell Science.
- Ghazoul, J. 2005 Pollen and seed dispersal among dispersed plants. *Biol. Rev.* **80**, 413–443. (doi:10.1017/S1464793105006731)

- Goulson, D. 1999 Foraging strategies of insects for gathering nectar and pollen, and implications for plant ecology and evolution. *Persp. Plant Ecol. Evol. Syst.* **2**, 185–209. (doi:10.1078/1433-8319-00070)
- Harder, L. D. & Johnson, S. D. 2005 Adaptive plasticity of floral display size in animal-pollinated plants. *Proc. R. Soc. B* **272**, 2651–2657. (doi:10.1098/rspb.2005.3268)
- Hobbs, R. J. & Yates, C. J. 2003 Impacts of ecosystem fragmentation on plant populations: generalising the idiosyncratic. *Aust. J. Bot.* **51**, 471–488. (doi:10.1071/BT03037)
- Jersakova, J. & Johnson, S. D. 2007 Protandry promotes male pollination success in a moth-pollinated orchid. *Funct. Ecol.* **21**, 496–504. (doi:10.1111/j.1365-2435.2007.01256.x)
- Johnson, S. D., Peter, C. I. & Ågren, J. 2004 The effects of nectar addition on pollen removal and geitonogamy in the non-rewarding orchid *Anacamptis morio*. *Proc. R. Soc. B* **271**, 803–809. (doi:10.1098/rspb.2003.2659)
- Johnson, S. D., Neal, P. R. & Harder, L. D. 2005 Pollen fates and the limits on male reproductive success in an orchid population. *Biol. J. Linn. Soc.* **86**, 175–190. (doi:10.1111/j.1095-8312.2005.00541.x)
- Kéry, M., Matthies, D. & Spillmann, H.-H. 2000 Reduced fecundity and offspring performance in small populations of the declining grassland plants *Primula veris* and *Gentiana lutea*. *J. Ecol.* **88**, 17–30. (doi:10.1046/j.1365-2745.2000.00422.x)
- Klinkhamer, P. G. L. & de Jong, T. J. 1990 Effects of plant size, plant density and sex differential nectar reward on pollinator visitation in the protandrous *Echium vulgare* (Boraginaceae). *Oikos* **57**, 399–405. (doi:10.2307/3565970)
- Lamont, B. B., Klinkhamer, P. G. L. & Witkowski, E. T. F. 1993 Population fragmentation may reduce fertility to zero in *Banksia goodii*: a demonstration of the Allee effect. *Oecologia* **94**, 446–450. (doi:10.1007/BF00317122)
- Leimu, R., Mutikainen, P., Koricheva, J. & Fischer, M. 2006 How general are positive relationships between plant population size, fitness and genetic variation? *J. Ecol.* **94**, 942–952. (doi:10.1111/j.1365-2745.2006.01150.x)
- Nilsson, L. A. 1992 Orchid pollination biology. *Trends Ecol. Evol.* **7**, 255–259. (doi:10.1016/0169-5347(92)90170-G)
- Peakall, R. 1989 A new technique for monitoring pollen flow in orchids. *Oecologia* **79**, 361–365. (doi:10.1007/BF00384315)
- Waites, A. R. & Ågren, J. 2004 Pollinator visitation, stigmatic pollen loads and among-population variation in seed set in *Lythrum salicaria*. *J. Ecol.* **92**, 512–526. (doi:10.1111/j.0022-0477.2004.00893.x)
- Ward, M. & Johnson, S. D. 2005 Pollen limitation and demographic structure in small fragmented populations of *Brunsvigia radulosa* (Amaryllidaceae). *Oikos* **108**, 253–262. (doi:10.1111/j.0030-1299.2005.13468.x)
- Young, A., Boyle, T. & Brown, T. 1996 The population genetic consequences of habitat fragmentation for plants. *Trends Ecol. Evol.* **11**, 413–418. (doi:10.1016/0169-5347(96)10045-8)