

## ON THE BINDING SITE OF QUINOLONE ANTIBACTERIALS. AN ATTEMPT TO PROBE THE SHEN MODEL

Stephen Hanessian,\* Raffaele Saladino, and Jose Cid Nunez

Department of Chemistry, Université de Montréal  
C.P. 6128, Centre-Ville, Montréal, P.Q., CANADA, H3C 3J7

**Abstract:** Quinolone-nucleic acid base hybrids were synthesized in an effort to probe a mechanistic model and a proposed mode of antibacterial action where stacked pairs of quinolones interact with DNA through H-bonding.

Copyright © 1996 Elsevier Science Ltd

The quinolones represent a class of potent antibacterial agents that have acquired widespread importance clinically.<sup>1</sup> In spite of their comparatively simple structures *vis-a-vis* other more complex antibiotics,<sup>2</sup> their mode of action at the molecular level remains a subject of debate and some controversy. There is a general consensus that the functional target of the quinolones is the bacterial enzyme DNA gyrase, which is required for the initiation and propagation of DNA synthesis.<sup>3</sup> Since DNA is involved, it is also agreed that the mode of action of quinolones comprises a three component complex, the molecular nature of which is currently of great interest.<sup>4</sup> In 1989, Shen, Pernet, Mitscher and their coworkers<sup>5</sup> proposed a cooperative quinolone-DNA binding model in which DNA-gyrase is intimately involved in a drug-enzyme recognition process. Through extensive SAR and the synthesis of





Figure 1. A. Shen model of quinolone-DNA gyrase-DNA interaction. B. Quinolone-nucleic acid base hybrid interactions with DNA gyrase and DNA.

**Figure 2.** Hypothetical interactions of proposed quinolone-adenine/cytosine hybrids with complementary bases in DNA.


Although we had no quantitative measure that such quinolone-nucleic acid base hybrids would indeed mimic the drug-drug associated model, we were encouraged by the results of binding specificities of quinolones to a single as compared to double stranded DNA.<sup>11</sup> It has been demonstrated that the binding of norfloxacin to poly(dG), poly(dA), poly(dT), and poly(dC) decreases in that order, with a distinct preference for the poly(dG). Binding to double stranded poly(dA)-poly(dT) was virtually undetectable. These and other results have led Shen and coworkers<sup>3-5</sup> to suggest that quinolones exhibit preferential binding to unpaired guanine bases in a single stranded DNA region, possibly through hydrogen bonds involving the 4-keto/3-carboxyl group. We therefore proceeded with the synthesis and biological evaluation of several quinolone-base hybrids, based on the predictions of the Shen model. The synthesis of four quinolone-purine/pyrimidine hybrid molecules **1-4** is shown in Scheme 1.

Scheme 1



(a) i: Ethyl-2,4-dichloro-5-fluoro benzoylacetate (1.2 equiv.),  $\text{CH}(\text{OC}_2\text{H}_5)_3$ ,  $\text{Ac}_2\text{O}$ , 130 °C, 3 h. ii: add **5a**, **6a**, **7a**, **8a**  $\text{CH}_2\text{Cl}_2/\text{EtOH}$  (1:1 v/v), 25 °C, 12 h.; (b) i: NaH (1.5 equiv.), toluene, reflux, 12 h.; ii: NaOH, THF/ $\text{H}_2\text{O}$ , 80 °C, 16 h. (c) Piperazine, 1-methyl-pyrrolidone, 80 °C, 12-24 h., for **1**, **2**, **3**, **4**; 76%, 90%, 75%, 54% respectively.

The required 4-amino-1-substituted bases were prepared by reacting 1-bromo-4-chlorobutane with the respective bases, followed by displacement of the chloro group with sodium azide ( $\text{NaN}_3$ ,  $\text{Bu}_4\text{NBr}$ ,  $25^\circ\text{C}$ ,  $\text{MeCN}$  reflux), then hydrogenation with  $\text{Ra-Ni}$ . The 4-aminobutyl heterocycles **5a-8a** were then condensed with the readily available<sup>12</sup> aromatic precursor (Scheme 1) to yield the enamino derivatives **9a-d** in good to excellent yields. Solubility problems intervened to provide a more modest yield for the guanine analog **9d**. Cyclization to the quinolones **10a-d** were carried out in a routine manner, and subsequent introduction of the piperazine unit at C-7 led to the intended hybrid analogs **1-4**, obtained as amorphous solids. Although these were sparingly soluble in common organic solvents, they were all adequately characterized by NMR and other analytical techniques.<sup>13</sup>



A. Complementary DNA Interaction Domain of compound **4**.

B. Complementary DNA and Gyrase interaction domain of compound **5**.

Figure 3. A. Proposed interaction of the quinolone-guanine hybrid **4** with a GGCC nucleotide sequence. B. Proposed interaction of a quinolone-8-morpholinoadenine hybrid with a TTAT nucleotide sequence.

Gyrase-dependent DNA cleavage assays by the standard agarose gel electrophoresis method<sup>14</sup> at  $50\text{ }\mu\text{g/mL}$  or less of compounds **1-3** in 5% DMSO as solvent, exhibited no drug-stimulated DNA cleavage. In an effort to probe the influence of the nature of the C-7 substituent, we prepared the N-methylpiperazine, morpholino, and pyrrolidino analogs related to the quinolone-thymine hybrid **1**. Also, the adenine analog **2** was prepared with a three-carbon linker chain. Again, biological activity was disappointingly absent, although the C-7 morpholino derivative of **1** gave values approaching Nalidixic acid ( $70\text{ }\mu\text{g/mL}$  vs  $58\text{ }\mu\text{g/mL}$  respectively). The base-pairing design in compounds **1-3** was inspired in part from the fact that the nucleotide sequence in *Micrococcus Luteus*

DNA is TTAT,<sup>15</sup> hence the attachment of T and A units in hybrids **1** and **2**. However, the cleavage site on pBR322 DNA (a derivative of ColE1 DNA) has the nucleotide sequence CGCC between the staggered cuts with *E. coli* DNA gyrase.<sup>16</sup> We therefore prepared the quinolone-guanine hybrid **4** in anticipation of a complementary interaction as shown in Figure 3A. Again, no gyrase-dependent DNA cleavage was observed in the appropriate test system involving these preparations with compound **4**. We reasoned that the lack of a second enzyme-binding domain in the inactive analogs **1-4** was in part the reason for their inactivity. Accordingly, we prepared the 8-morpholinoadeninyl hybrid molecule **5**<sup>17</sup> depicted in Figure 3B. In such a derivative the stacked quinolone and adeninyl portions would still satisfy H-bonded interactions with the DNA, with the added feature that the piperazino and morpholino groups might effectively bind to the enzyme as in the original proposal especially if they are properly orientated in space. Unfortunately, this "rationally designed" hybrid also lacked activity in the gyrase-dependent DNA cleavage assays.

In conclusion, we have attempted to probe the Shen model<sup>5</sup> for the interaction of quinolone antibacterials with DNA and DNA gyrase from bacterial sources. The hybrid molecules, consisting of a quinolone linked to purine and pyrimidine bases through three and four methylene bridges were designed and predicated upon the presumed existence of H-bonding and enzyme binding domains as in the proposed three-component model.<sup>5</sup> The absence of enhancement of gyrase-dependent DNA cleavage in these analogs may be due to unfavorable three-dimensional features, and the deployment of critical functionality that deviate substantially from the elusive yet prevailing situation with the quinolones themselves. Thus, while the results of this study do not support the Shen model,<sup>5</sup> we cannot exclude its validity at this time. Clearly, additional studies are needed to shed some light on the fascinating roles that the quinolones play at the molecular level in conjunction with DNA gyrase and DNA itself. We are currently reevaluating our options for the design and synthesis of other quinolone prototypes in order to gain a better understanding of the relevance of the currently proposed models<sup>5-7</sup> with regard to their antibacterial mode of action.

**Acknowledgments.** We thank NSERCC for generous financial assistance through the Medicinal Chemistry Chair program. We are very grateful to Dr. Linus Shen of Abbott Laboratories for the biological tests and for stimulating discussions. R. Saladino and J. C. Nunez acknowledge fellowships from MURST (Italy) and Xunta de Galicia (Spain), respectively.

## References and Notes

1. For a selection of monographs, see Siporin, C.; Heifetz, C. L.; Domagala, J.-M. In *The New Generation of Quinolones*; M. Dekker: New York, 1990; Crumplin, G. C. In *The 4-Quinolones: Antibacterial Agents in Vitro*; Springer Verlag: New York, 1990; Andriole, V. T. In *The Quinolones*; Academic: New York, 1989; Wolfson, J. S.; Hooper, D. C. In *Quinolone Antimicrobial Agents*; American Society for Microbiology, Washington, D.C 1989; Fernandez, P. B. Ed.; Int. Telesymposium of Quinolones, J. R. Prous, Barcelona, Spain; for recent reviews, see Domagala, J. M. *J. Antimicrob. Chemother* **1994**, *33*, 685; Hammond, M. L. *Ann. Rep. Med. Chem.* **1993**, *28*, 119.

2. See for examples, such classes as macrolides, tetracyclines,  $\beta$ -lactams, In *Emerging Targets in Antibacterial and Antifungal Chemotherapy*; Sutcliffe, J.; Georgopapadakou, N. Eds.; Chapman and Hall: New York, 1992; *Antibiotics and Antiviral Compounds. Chemical Synthesis and Modifications*; Krohn, K.; Kirst, H. A.; Maag, H.; Eds; VCH: New York, 1993.
3. For a recent review, see Shen, L. L. *Adv. Pharmacol.* **1994**, 29A, 285; and references cited therein; Reece, R. J.; Maxwell, A. *CRC-Critical Rev. Biochem. Mol. Biol.* **1991**, 26, 335.
4. For recent discussions, see Shen, L. L. In *Molecular Biology of DNA Topoisomerases and its Application to Chemotherapy*, Andoh, T.; Ikeda, H.; Oguro, M. Eds.; CRC: Ann Arbor, 1993; p 177; Shen, L. L. In *Quinolone Antimicrobial Agents*, Hooper, D. C.; Wolfson, J. S., Eds; American Society of Microbiology, Washington, D.C., 1993; p 11.
5. Shen, L. L.; Mitscher, L. A.; Sharma, P. N.; O'Donnell, T. J.; Chu, D. T. W.; Cooper, C. S.; Rosen, T.; Pernet, A. G. *Biochem.* **1989**, 28, 3886.
6. Palumbo, M.; Gatto, B.; Zagotto, G.; Palu, G. *Trends in Microbiol.* **1993**, 1, 232; Palu, G.; Valisena, S.; Ciarrocchi, G.; Gatto, B.; Palumbo, M. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, 89, 9671.
7. Fam, J.; Sun, D.; Kerwin, S. M.; Hurley, L. H. *J. Med. Chem.* **1995**, 38, 408.
8. Maxwell, A. *J. Antimicrob. Chemother.* **1992**, 30, 409.
9. Llorente, B.; Leclerc, F.; Cedergren, R. *Bioorg. Med. Chem.* **1996**, 4, 61.
10. For some examples of drugs that interact with DNA, see Perun, T. J.; Propst, C. L. In *Nucleic Acid Targeted Drug Design*; Propst, C. L.; Perun, T. J. Eds., M. Dekker: New York, 1992, p 1; and other chapters.
11. Shen, L. L.; Pernet, A. G. *Proc. Natl. Acad. Sci. U.S.A.* **1985**, 82, 307.
12. Koga, H.; Itoh, A.; Murayama, S.; Suzue, S.; Irikura, T. *J. Med. Chem.* **1980**, 23, 1358.
13. All new compounds were characterized by  $^1\text{H}$ ,  $^{13}\text{C}$  NMR spectroscopy, by HRMS and microanalytical data when possible.
14. Shen, L. L.; Baranowski, J.; Fostel, J.; Montgomery, D. A.; Larty, P. A. *Antimicrob. Agents Chemother.* **1992**, 36, 2778.
15. Kirkegaard, K.; Wang, J. C. *Cell* **1981**, 23, 721.
16. Fisher, L. M.; Mizuuchi, K.; O'Dea, M.H.; Ohmori, H.; Gellert, M. *Proc. Natl. Acad. Sci. USA* **1981**, 78, 4165; Fisher, L. M.; Barot, H. A.; Cullen, M. E. *EMBO* **1986**, 5, 1411.
17. Prepared from 8-bromoadenine via: (a) morpholine, 140 °C, 4 h, 75%; (b) NaH, THF, then 4-bromo-1-chlorobutane, 25 °C, 2d; (c)  $\text{NaN}_3$ ,  $\text{Me}_4\text{NBr}$ , NaI, MeCN, reflux 24h, 50-50% (2 steps); (d) Pd/C,  $\text{H}_2$ , EtOH, 90%; (e) condensation according to Scheme 1.

(Received in USA 18 July 1996; accepted 6 September 1996)