



NOVEL NON-NUCLEOSIDIC PHOSPHORAMIDITES FOR OLIGONUCLEOTIDE MODIFICATION AND LABELING


Sheng-Hui Su*, Rajkumar S. Iyer, Sunil K. Aggarwal, and Krishan, L. Kalra

BioGenex Laboratories, 4600 Norris Canyon Road, San Ramon, CA 94583, U.S.A.

Abstract: A series of novel labeled phosphoramidites and CPG supports **1a–f** with a new non-nucleosidic backbone based upon cyclohexyl-4-amino-1,1-dimethanol have been synthesized. These reagents have been used to label oligonucleotides with biotin and fluorescein at the 5'-, the 3'-, and internal positions.

© 1997 Elsevier Science Ltd.

Oligodeoxynucleotides bearing suitable reporter groups have proven to be potent tools in molecular biology, finding applications in diagnostics, sequencing, and molecular studies of nucleic acids.^{1,2} Various nucleosidic as well as non-nucleosidic phosphoramidites have been successfully incorporated into oligomers to obtain the desired functionalities. The non-nucleosidic tails used for labeling of oligonucleotides include both branched and linear structures. Branched dendrimeric structures have, for example, been assembled on the 5'-terminus of the oligonucleotide by using bis(4,4'-dimethoxytrityl)alkanetriol phosphoramidite as a building block.^{3–5} Linear tails have also, in turn, been prepared by using non-nucleosidic phosphoramidites derived from



Compound	X	Label	R
1a	-	Biotinyl	-O-P-OCH ₂ CH ₂ CN(N- ¹ Pr ₂)
1b	-CO(CH ₂) ₅ NH-	N-(4- ¹ Bu-Bz)-Biotinyl	-O-P-OCH ₂ CH ₂ CN(N- ¹ Pr ₂)
1c	-CO-	Fluoresceinyl	-O-P-OCH ₂ CH ₂ CN(N- ¹ Pr ₂)
1d	-CO-	-(CH ₂) ₅ NHCOCF ₃	-O-P-OCH ₂ CH ₂ CN(N- ¹ Pr ₂)
1e	-	Biotinyl	-OCO(CH ₂) ₂ CONH-CPG
1f	-	Fluoresceinyl	-OCO(CH ₂) ₂ CONH-CPG

- i. BzCl , pyridine; ii. $\text{BF}_3\text{Et}_2\text{O}$, NaBH_4 , diglyme, $\text{H}_2\text{NSO}_3\text{H}$; iii. Biotin-NHSu, TEA, CH_2Cl_2 ; iv & v. CH_3ONa , CH_3OH ; vi. Fluorescein-NHSu, TEA, DMF; vii. DMTCL, pyridine; viii. $\text{CNCH}_2\text{CH}_2\text{O-P}(\text{N-}i\text{Pr})_2$, tetrazole-DIPA salt; ix. $\text{CNCH}_2\text{CH}_2\text{O-P}(\text{Cl})(\text{N-}i\text{Pr})_2$, DIPEA.

The reaction of compound **4** with biotin-N-hydroxysuccinimide ester (biotin-NHSu) in the presence of triethylamine afforded the corresponding amide **5** in 90% purified yield. Removal of the benzoyl group from **5** followed by dimethoxytritylation resulted in the formation of compound **9**. It is likely that compound **9** and its derivatives exist as a mixture of geometric isomers. However, they were not resolved chromatographically and no attempt was made to assign their stereochemistry. The phosphitylation of compound **9** under standard conditions gave the biotin phosphoramidite **1a** in 75% yield.²⁰ Phosphoramidite **1a** when used as a 0.1 M solution in acetonitrile using standard 0.2 μ mol scale DNA synthesis protocols, exhibits consistently high coupling efficiency (>98%).²¹ Figure 1A shows the capillary electrophoresis trace of a crude 17mer labeled at 5'-position by using biotin amidite **1a**. The desired full length oligo was 70% of the crude product mixture.²² In order to facilitate the kinetics of biotin binding to streptavidin, we have also synthesized the long chain biotin phosphoramidite **1b**.

For fluorescein amidite, the benzoyl derivative **4** was initially deprotected with sodium methoxide in methanol to give compound **7**. Treatment of 5(6)-carboxyfluorescein with pivaloyl chloride gave the non-fluorescent lactone which was converted to its NHSu ester with dicyclohexylcarbodiimide and N-hydroxysuccinimide.¹² The coupling of NHSu ester with compound **7** in pyridine followed by dimethoxytritylation and phosphitylation gave the phosphoramidite derivative **1c**.²³ The fluorescein phosphoramidite **1c**, was used in the same manner as nucleoside phosphoramidites on an automated DNA synthesizer. However, a coupling time of 10 min was used for fluorescein incorporation which resulted in a coupling efficiency of > 97%.²⁴ This is a rather significant improvement over other non-nucleosidic fluorescein




Figure 1: Microgel capillary electrophoretic traces of crude labeled oligonucleotides synthesized using biotin phosphoramidite (**1a**) and fluorescein phosphoramidite (**1c**). **Panel A:** biotinylated oligonucleotide 5'- B-GTA ATG CGA GCG CCA GT-3', where B = biotin. **Panel B:** fluoresceinated oligonucleotide 5'-F-TGG TCC GCT AAG ACA TTA GT-3', where F = fluorescein.

phosphoramidites that have been reported.¹² Figure 1B shows the capillary electrophoresis trace of a 20mer labeled at 5'-position with **1c**. The full length oligo was 76% of the crude mixture.

For functionalization of oligonucleotides with primary amino group, we have also synthesized the amino phosphoramidite **1d**. This phosphoramidite is extremely stable and under normal conditions of automated synthesis, it consistently gave a coupling efficiency >98%. This compound can be used for the post-labeling of oligonucleotides with other reporter groups such as acridine and enzymes in addition to biotin and fluorescein.²⁵

In order to label oligonucleotides with biotin and fluorescein at the 3' end, biotin-CPG (500 Å) and fluorescein-CPG (500 Å) were synthesized by reaction of the corresponding succinate with LCCA-CPG.²⁶ Both biotin-CPG (loading 31.3 µmol/g) and fluorescein-CPG (loading 9.3 µmol/g) gave excellent coupling efficiency results (>99%) when 20 mers were synthesized using standard synthesis cycles.

Biotin phosphoramidite **1b** was tested as follows: the 30mer 5'-p-TTT TTT TTT TCG BAA TCA BGG TCA BAG CTG-3' (B = biotin, p = phosphate) was synthesized with either B = **1b** or B = commercially available nucleosidic biotin-dC phosphoramidite. Both oligos were covalently bound to polystyrene (Covalink-NH₂) microtiter plates via 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide mediated coupling of the primary amino and phosphate groups. After washing off the unbound oligonucleotides, the biotin residues were detected by binding of streptavidin-alkaline phosphatase conjugate. Para-nitrophenyl phosphate was used as the substrate for the enzyme and the development of color was monitored at 405 nm. At concentrations of 5 fmol/µL and 10 fmol/µL, the observed signal intensity (OD 405 nm) was essentially identical for both the oligos tested. This indicates that the non-nucleosidic biotin phosphoramidite **1b** performs as well as nucleosidic biotin-dC phosphoramidite in functional assays. The ability of oligonucleotides synthesized using phosphoramidite **1b** to participate in hybridization assays was then tested and the results compared to those obtained by using commercially available nucleosidic biotin-dC phosphoramidite. The 56mer, Alu-011 with one internal biotin residue, is complementary to the template Alu-011A and has the sequence 5'-GGC CGG GCG CGG TGG CTC ACG CGT TAA BTT AAC GCT GTA ATC CCA GCA CTT TGG GA-3'. The template Alu-011A was immobilized on CovaLink polystyrene microtiter plates as described above. The biotinylated oligonucleotides (25 fmol/µL) were then hybridized with the template for 5 h at 42 °C. After washing off the excess probes, the bound biotinylated probes were detected by incubating with streptavidin-alkaline phosphatase conjugate. Under identical conditions, the **1b** labeled oligomers generated approximately a fivefold stronger signal than oligomers labeled with biotin-dC.

Preliminary results on the testing of fluorescein phosphoramidite **1c**, indicate that probes synthesized using **1c** allowed for the more sensitive detection of kappa immunoglobulin light chains on paraffin embedded tissue sections in in situ hybridization assays. Fluoresceinated probes synthesized using **1c** gave much stronger signal and less background relative to similar probes that had the fluorescein moiety introduced post-synthetically.²⁴

To conclude, we report novel biotin, fluorescein and amino labeled phosphoramidites and CPG supports based upon the cyclohexyl backbone. These compounds can be used advantageously for the introduction of multiple reporter groups onto oligonucleotides in a cost effective and efficient manner. The natural 3-carbon atom internucleotide phosphate distance in DNA/RNA is retained in the above structures which will not affect the hybridization and annealing properties of the duplex. The reporter group is kept outside of the nucleotide chain by the rigid cyclohexane ring resulting in high coupling efficiency and oligonucleotide yields.

Acknowledgment: Technical assistance received from Deguang Yang and Zhong-Yi Lu is gratefully acknowledged. The authors wish to thank Kenneth A. Sikes for critical evaluation of the manuscript.

References and Notes

1. Uhlman, E.; Peyman, A. *Chem. Rev.* **1990**, *90*, 543.
2. Beaucage, S. L.; Iyer, R. P. *Tetrahedron* **1993**, *49*, 1925.
3. Bazin, H.; Roget, A.; Teoule, R. *Nucleosides Nucleotides* **1991**, *10*, 363.
4. Teigelkamp, S.; Ebel, S.; Will, D. W.; Brown, T.; Bebbs, J. D. *Nucleic Acids Res.* **1993**, *21*, 4651.
5. De. Vos, M. J.; Van Elsen, A.; Bollen, A. *Nucleosides Nucleotides* **1994**, *13*, 2245.
6. Nelson, P. S.; Kent, M.; Muthini, S. *Nucleic Acids Res.* **1992**, *20*, 6253.
7. Korshun, V. A.; Pestov, N. B.; Birikh, K. R.; Berlin, Y. A. *Bioconjugate Chem.* **1992**, *3*, 559.
8. Burmeister, J.; Azzawi, A.; von Kiedrowski, G. *Tetrahedron Lett.* **1995**, *36*, 3667.
9. Kane, R. R.; Drechsel, K.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1993**, *115*, 8853.
10. Drechsel, K.; Lee, C. S.; Leung, E. W.; Kane, R. R.; Hawthorne, M. F. *Tetrahedron Lett.* **1994**, *35*, 6217.
11. Misiura, K.; Durrant, I.; Evans, M. R.; Gait, M. J. *Nucleic Acids Res.* **1990**, *18*, 4345.
12. Theisen, P.; McCollum, C.; Upadhy, K.; Jacobson, K.; Vu, H.; Andrus, A. *Tetrahedron Lett.* **1992**, *33*, 5033.
13. Neuner, P. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 147.
14. Behrens, C.; Petersen, K. H.; Egholm, M.; Nielsen, J.; Buchardt, O.; Dahl, O. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 1785.
15. Misiura, K.; Durrant, I.; Evans, M. R.; Gait, M. J. *Nucleosides Nucleotides* **1991**, *10*, 671.
16. Guzaev, A.; Salo, H.; Azhaye, A.; L'nnberg, H. *Bioconjugate Chem.* **1996**, *7*, 240.
17. Su, S.; Iyer, R. S.; Aggarwal, S. K.; Kalra, K. L. Application for U.S. patent pending.
18. Brown, H. C.; Heydkamp, W. R.; Breuer, E.; Murphy, W. S. *J. Am. Chem. Soc.* **1964**, *86*, 3565.

19. Compound **4**: ^1H NMR (360 MHz, $\text{DMSO}-d_6$) δ 7.9 (m, 4H, ortho ArH), 7.55 (m, 2H, para ArH), 7.4 (m, 4H, meta ArH), 4.3 (s, 2H, $-\text{OCH}_2$), 4.1 (s, 2H, $-\text{OCH}_2$), 2.92 (bs, 2H, NH_2), 2.6 (m, 1H, $-\text{CH}$), 1.15–1.75 (m, 8H, 4X CH_2).
20. Compound **1a**: ^1H NMR (360 MHz, CDCl_3) δ 7.4–6.65 (m, 13H, ArH), 6.1 (s, 1H, NH), 5.45 (d, 1H, NH), 5.35 (s, 1H, NH), 4.45–4.1 (m, 2H, $-\text{OCH}_2$), 3.75 (s, 6H, 2X OCH_3), 3.6–3.25 (m, 2H, $-\text{OCH}_2$), 2.8 (m, 2H, $-\text{CH}_2\text{S}-$), 2.7 (m, 1H, CH), 2.5 (m, 1H, CH), 2.2–1.4 (m, 24H, ring and chain CH_2), 1.3–0.9 (dd, 12H, 4X $\text{CH}_3\text{-CH}$); ^{31}P NMR (146 MHz, CDCl_3) 147.9, 148.4 ppm, M/S M^+ 888 ($\text{C}_{48}\text{H}_{66}\text{N}_5\text{O}_7\text{PS}$).
21. Oligodeoxynucleotides were synthesized using a Eppendorf D-100 DNA Synthesizer utilizing standard β -cyanoethyl N,N-diisopropyl phosphoramidite monomers (dA^{Bz} , dG^{tBu} , and dC^{Bz}). Oligodeoxynucleotides were cleaved from the support and deprotected by heating at 55 °C overnight in concentrated aqueous ammonia solution. Fluoresceinated oligodeoxynucleotides were deprotected by heating at 60 °C for 5 h in concentrated aqueous ammonia solution.
22. The integrity of the label was confirmed by checking the MALDI-TOF mass spectrum of a 5'-biotin-(T)₁₉-T-3' (biotin = **1a**) oligomer: M^+ calculated for $\text{C}_{218}\text{H}_{291}\text{N}_{43}\text{O}_{144}\text{P}_{20}\text{S}$ = 6466.12; found = 6479.08 utilizing a 2,4,6-trihydroxyacetophenone (THAP) matrix.
23. Compound **1c**: ^1H NMR (360 MHz, CDCl_3) δ 6.8–8.1 (m, 22H, ArH), 3.4–3.85 (m, 12H), 2.55–2.95 (m, 4H), 1.9–2.1 (m, 6H), 1.6–1.85 (m, 5H), 1.15–1.45 (m, 4X CH_3 , 4X CH_2), 1.2 (dd, 12H, 4X $\text{CH}_3\text{-CH}$); ^{31}P NMR (146 MHz, CDCl_3) 147.6, 147.1 ppm.
24. Preliminary results obtained from the use of phosphoramidite **1c** in in situ hybridization assays were presented at the Annual Symposium of National Society for Histotechnology held in Albuquerque, U.S.A., October 19-23, 1996, Abstract P10.
25. Chu, B. F.; Orgel, L. E. In *Protocols for Oligonucleotide conjugates*; Agrawal, S. Ed.; Humana: New Jersey, 1994; Vol. 26, pp 145–165.
26. Pon, R. In *Protocols for oligonucleotides and analogs*; Agrawal, S. Ed.; Humana: New Jersey, 1993; Vol. 20, pp 465–468.

(Received in USA 14 January 1997; accepted 22 May 1997)