

Corrigendum


Corrigendum to "Synthesis and structure–activity relationships of 7-(2-aminoalkyl)morpholinoquinolones as anti-*Helicobacter pylori* agents" [Bioorg. Med. Chem. Lett. 8 (1998) 2185][†]

Nobuhiro Sakurai,* Mitsuharu Sano, Fumihiro Hirayama, Tsuyoshi Kuroda,
Satoru Uemori, Akihiko Moriguchi, Katsuhiro Yamamoto, Yoshifumi Ikeda and
Takeshi Kawakita

Research Laboratories, Yoshitomi Pharmaceutical Industries Ltd., 955 Koiwai, Yoshitomi-cho, Chikugo-gun,
Fukuoka 871-8550, Japan

On page 2188, Table 1, the unit of MIC should read $\mu\text{g/mL}$ as shown below:

Table 1. Antimicrobial activities of 7-(2-aminoalkyl)morpholinoquinolones and other antimicrobial agents against *H. pylori* ATCC 43504



Compd. No.	R	X	mp (°C)	MIC ($\mu\text{g/mL}$)
1 (Y-26611)	NH ₂	F	180–182	0.05
2	NHMe	F	251–254	0.05
(R)-11	NMe ₂	OMe	144–145	0.39
(S)-11 (Y-34867)	NMe ₂	OMe	144–146	0.012
levofloxacin				0.20
tosufloxacin				0.39
sparfloxacin				0.20
amoxicillin				0.025
clarithromycin				0.025

[†]PII of original article: S0960-894X(98)00390-4