

Corrigendum

Corrigendum to “Formation of Glucoraphanin by Chemoselective Oxidation of Natural Glucoerucin: A Chemoenzymatic Route to Sulforaphane” (*Bioorg. Med. Chem. Lett.* 1999, 9 1047–1048)[†]

R. Iori,^a R. Bernardi,^a D. Gueyrard,^b P. Rollin^{b,*} and S. Palmieri^a

^a*Istituto Sperimentale per le Colture Industriali, MiPA, Via di Corticella, 133, I-40129 Bologna, Italy*

^b*Institut de Chimie Organique et Analytique, Université d'Orléans, BP 6759, F-45067 Orleans, Cedex 2, France*

Reference 9 is incorrect and should read as follows:

9. ¹H NMR (D₂O): 1.93 (m, 2 H, H-10), 1.98 (m, 2 H, H-9), 2.77 (m, 3 H, H-12), 2.88 (t, 2 H, $J_{8,9}$ = 6.8 Hz, H-8), 3.02 (m, 2 H, H-11), 3.54 (m, 2 H, H-2 and H-4), 3.65 (m, 2 H, H-3 and H-5), 3.78–3.97 (2 dd, 2 H, $J_{6a,6b}$ = 12.5 Hz, $J_{6a,5}$ = 1.7 Hz, $J_{6b,5}$ = 5.4 Hz, H-6a and H-6b), 5.12 (d, 1 H, $J_{1,2}$ = 9.5 Hz, H-1); ¹³C NMR (D₂O): 21.9 (C-10), 26.0 (C-9), 32.3 (C-8), 37.0 (C-12), 52.8 (C-11), 61.3 (C-6), 69.8 (C-4), 72.6 (C2), 77.7 (C-3), 80.8 (C-5), 82.4 (C-1), 164.4 (C-7).

*Corresponding author. Tel.: +33-2-3841-7370; fax: +33-2-3849-4579; e-mail: patrick.rollin@univ-orleans.fr

[†]PII of original article: S0960-894X(99)00136-5