

Corrigendum


Corrigendum to “Amino Acid Derived Sulfonamide Hydroxamates as Inhibitors of Procollagen C-Proteinase: Solid-Phase Synthesis of Ornithine Analogues” [Bioorg. Med. Chem. Lett. 11 (2001) 2085][†]

Sharon M. Dankwardt,^{a,*} Robert L. Martin,^a Christine S. Chan,^a Harold E. Van Wart,^a Keith A. M. Walker,^a Nancy G. Delaet^b and Leslie A. Robinson^b

^aRoche Bioscience, Inflammatory Diseases Unit, 3401 Hillview Ave, Palo Alto, CA 94304, USA

^bDuPont Pharmaceuticals Company Research Laboratories, formerly CombiChem Inc., 4570 Executive Drive Ste. 400, San Diego, CA 92121, USA

Scheme 1 contained an error. The corrected version is shown below.



Scheme 1. For resin **A**: Conditions for E⁺ = RCOOH, HATU/HOAt/DIPEA/DMF/CH₂Cl₂ E⁺ = ROCOOSu, pyr/CH₂Cl₂. Abbreviations: Dde = 1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl; ADDP = 1,1'-(azodicarbonyl)dipiperidine. For resin **B**: Conditions for E⁺ = ROCOOSu/ROCOCl/RNCO/RNCS: Et₃N/DMAP/THF, for ROCOCl/RSO₂Cl: pyr/CH₂Cl₂. TMAD, azodicarboxylic acid bis(dimethylamide).

*Corresponding author. Tel.: +1-650-354-2512; fax: +1-650-354-2442; e-mail: sharon.dankwardt@roche.com

[†]PII of original article: S0960-894X(01)00384-5