

Corrigendum

Corrigendum to “N-Isonicotinoyl-(L)-4-aminophenylalanine Derivatives as Tight Binding VLA-4 Antagonists” [Bioorg. Med. Chem. Lett. 13 (2003) 1891]☆


George A. Doherty,^{a,*} Ginger X. Yang,^a Edite Borges,^a Sharon Tong,^a
Ermengilda D. McCauley,^b Kelly M. Treonz,^b Gail Van Riper,^b Stephen Pacholok,^b
Qian Si,^b Gloria C. Koo,^b Kashmira Shah,^b Richard A. Mumford^b
and William K. Hagmann^a

^aDepartment of Medicinal Chemistry, Merck Research Laboratories, Rahway, NJ 07065, USA

^bDepartment of Immunology and Rheumatology Research, Merck Research Laboratories, Rahway, NJ 07065, USA

The authors regret that in the above article there was an error within the structures of Tables 2 and 3. The corrected tables are below:

Table 2. Inhibition of VLA-4 by sulfonlated 2(S)-methyl-proline-4-isonicotinoyl-aminophenylalanine derivatives




Compd	2	4	6	VLA-4 (IC ₅₀ , nM)		Off-rate (37 °C) % bound @ 1 h
				Mn ²⁺ , ^a	Ca ²⁺ , Mg ²⁺ , ^b	
5	Cl	—	Cl	0.11	0.11	57
7	H	—	Cl	0.14	0.21	26
8	H	—	H	0.27	0.25	12
9	Cl	—	OMe	0.13	0.08	40
10	Cl	O	Cl	0.10	0.08	61

^a¹²⁵I-VCAM-Ig, *n* = 3.

^b³⁵S-2, *n* = 3.

*Corresponding author. Tel.: +1-732-594-3686; fax: +1-732-594-5966; e-mail: george_doherty@merck.com

☆DOI of original article: 10.1016/S0960-894X(03)00308-1

Table 3. Inhibition of VLA-4 by sulfonlated 2(*S*)-methyl-proline-3-isonicotinoyl-aminophenylalanine derivatives

Compd	2	4	6	VLA-4 (IC ₅₀ , nM)		Off-rate (37°C) % bound @ 1 h
				Mn ²⁺ , ^a	Ca ²⁺ , Mg ²⁺ , ^b	
11	Cl	Me	Cl	0.19	0.10	61
12	H	Me	Cl	0.16	0.31	0
13	H	H	SMe	0.36	1.56	0

^a¹²⁵I-VCAM-Ig, *n* = 3.^b³⁵S-**2**, *n* = 3.