

Corrigendum

Corrigendum to “Structure–activity relationship of benzo[*b*]thiophene-2-sulfonamide derivatives as novel human chymase inhibitors” [Bioorg. Med. Chem. Lett. 13 (2003) 4085]☆


Hidekazu Masaki,^{a,*} Yusuke Mizuno,^a Akira Tatui,^b Akira Murakami,^b Yuuki Koide,^a Shoji Satoh^a and Atsuo Takahashi^a

^a*Drug Research Department, Tokyo Research Laboratories, TOA EIYO Ltd., 2-293-3 Amanuma-cho, Omiya-ku, Saitama-shi, Saitama 330-0834, Japan*

^b*Drug Research Department, Fukushima Research Laboratories, TOA EIYO Ltd., 1 Tanaka, Yuno, Iizaka-cho, Fukushima-shi, Fukushima 960-0280, Japan*

The authors regret that the general structure in Table 1 was not shown in the above article. The full table appears below.

Table 1. Effect of substitution on the benzo[*b*]thiophene ring and *N*-phenyl ring



Compd	X	R ¹	R ²	R ³	IC ₅₀ (nM) ^a Human chymase
1	Cl	H	SO ₂ (CH ₂) ₂ CH ₃	H	10,600
8	Cl	H	S(CH ₂) ₂ CH ₃	H	NI ^b
9	Cl	H	CN	H	NI
10	Cl	H	NO ₂	H	NI
11	Cl	H	CO ₂ Me	H	1647
12	Cl	NO ₂	CO ₂ Me	H	880
13	Cl	OMe	CO ₂ Me	H	663
14	Cl	CO ₂ Me	CO ₂ Me	H	NI
15	Cl	SO ₂ Me	CO ₂ Me	H	203
16	Cl	H	CO ₂ Me	CO ₂ Me	4208
17	Cl	SO ₂ Me	CO ₂ Me	Me	594
18	Me	SO ₂ Me	CO ₂ Me	H	325
19	H	SO ₂ Me	CO ₂ Me	H	305
7	F	SO ₂ Me	CO ₂ Me	H	56

^a Values are means of three experiments.

^b No inhibition at 10 μM.

☆ DOI of original article: 10.1016/j.bmcl.2003.08.040

* Corresponding author. Tel.: +81-48-647-7971; fax: +81-48-648-0078; e-mail: hidekazu.masaki@toaieyo.co.jp