

Corrigendum

**Corrigendum to “Identification of
4-piperazin-1-yl-quinazoline template based aryl and benzyl
thioureas as potent, selective, and orally bioavailable inhibitors
of platelet-derived growth factor (PDGF) receptor”
[Bioorg. Med. Chem. Lett. 14 (2004) 4867]**

Julie A. Heath,* Mukund M. Mehrotra,* Shannon Chi, Jin-Chen Yu,
Athiwat Hutchaleelaha, Stanley J. Hollenbach, Neil A. Giese,
Robert M. Scarborough and Anjali Pandey

*Departments of Medicinal Chemistry, Biology DMPK, and In Vivo Sciences Millennium Pharmaceuticals, Inc.,
256 E. Grand Avenue, South San Francisco, CA 94080, USA*

Available online 26 October 2004

Regrettably, the units of measurement used in Table 8 for $C_{\max}$ and AUC were incorrect. The corrected table is published below.

Table 8. Pharmacokinetic data of selected analogues

	Analogue		
	14	49	25
<i>Rat</i>			
Dose iv/po (mg/kg)	30	10	30
F (%)	13.7	8.82	39.1
$T_{1/2}$ (h)	14.1	17.4	26.7
$C_{\max}$ (ng/mL)	239	34.1	146
AUC_{PO} (ng h mL ⁻¹)	32.2	26.9	69.0
<i>Dog</i>			
Dose iv/po (mg/kg)	8.69	5	8.09
F (%)	38.8	30.2	101.6
$T_{1/2}$ (h)	10.8	7.58	15.7
$C_{\max}$ (ng/mL)	297	38.9	456
AUC_{PO} (ng h mL ⁻¹)	1507	333	3117
<i>Monkey</i>			
Dose iv/po (mg/kg)	10	10	10
F (%)	n.d.	68	n.d.
$T_{1/2}$ (h)	20.9	8.46	9.25
$C_{\max}$ (ng/mL)	235	143	92.5
AUC_{PO} (ng h mL ⁻¹)	832	1510	477

n.d. = not determined.

DOI of original article: 10.1016/j.bmcl.2004.07.041

* Corresponding authors at present address: Portola Pharmaceuticals, Inc., 270 E. Grand Ave., Suite 22, South San Francisco, CA 94080, USA.
Tel.: +1 650 246 7016; fax: +1 650 244 9287 (J.A.H.); e-mail addresses: jheath@portola.com; mukund_mehrotra@hotmail.com