



Corrigendum

**Corrigendum to “Synthesis and in vitro pharmacological studies
of new C(4) modified salvinorin A analogues”
[Bioorg. Med. Chem. Lett. 16 (2006) 5498–5502]**

David Y. W. Lee,^a Minsheng He,^a Lee-Yuan Liu-Chen,^b Yulin Wang,^b Jian-Guo Li,^b
Wei Xu,^b Zhongze Ma,^a William A. Carlezon, Jr.^c and Bruce Cohen^{d,*}

^aBioorganic and Natural Products Laboratory, McLean Hospital, Harvard Medical School, 115 Mill Street, Belmont, MA 02478, USA

^bDepartment of Pharmacology, School of Medicine, Temple University, 3420 N. Broad Street, Philadelphia, PA 19140, USA

^cBehavioral Genetics Laboratory, McLean Hospital, Harvard Medical School, 115 Mill Street, Belmont, MA 02478, USA

^dMolecular Pharmacology Laboratory, McLean Hospital, Harvard Medical School, 115 Mill Street, Belmont, MA 02478, USA

Available online 7 February 2008

The authors regret that this article contains two errors:

(1) On page 5500, left-hand column, in Scheme 6, “Synthesis of the isostere (**23**) of salvinorin acid” should be changed to “Synthesis of the tetrazole (**23**)”.

(2) On page 5502, the NMR data for compound **3** were incorrectly given, and those reported data belong to 8-*epi* **3**. The revised ¹H and ¹³C NMR data of **3** should be revised as follows: ¹H NMR (CDCl₃, 300 MHz) δ 9.91 (1H, d, *J* = 1.2 Hz), 7.43 (1H, d, *J* = 0.9 Hz), 7.41 (1H, t, *J* = 1.8 Hz), 6.38 (1H, dd, *J* = 0.9 and 1.8 Hz), 5.54 (1H, dd, *J* = 5.4 and 11.7 Hz), 5.17 (1H, dd, *J* = 7.2 and 11.7 Hz), 2.73 (1H, dd, *J* = 3.3 and 12.9 Hz), 2.51 (1H, dd, *J* = 5.1 and 13.5 Hz), 2.39 (1H, m), 2.10–2.30 (4H, m), 2.13 (3H, s), 1.55–1.90 (4H, m), 1.46 (3H, s), 1.07 (3H, s); ¹³C NMR (CDCl₃, 75 MHz) δ 202.2, 200.6, 170.9, 170.0, 143.8, 139.4, 125.1, 108.3, 75.2, 72.0, 63.6, 59.7, 51.2, 43.2, 42.5, 38.0, 35.4, 28.3, 20.6, 17.9, 17.8, 15.1.