

## Corrigendum

Corrigendum to "Synthesis and antiplatelet, antiinflammatory, and antiallergic activities of 2-substituted 3-chloro-1,4-naphthoquinone derivatives"

[Bioorg. Med. Chem. 5 (1997) 2111]<sup>1</sup>

J.-C. Liena, L.-J. Huanga, J.-P. Wangb, C.-M. Tengc, K.-H. Leed, S.-C. Kuo\*a

"Graduate Institute of Pharmaceutical Chemistry, China Medical College, Taichung, Taiwan

bDepartment of Medical Research, Taichung Veterans' Hospital, Taichung, Taiwan

Pharmacological Institute, College of Medicine, National Taiwan University, Taiwan

Matural Products Laboratory, Division of Medicinal Chemistry and Natural Products, School of Pharmacy,

University of North Carolina, Chapel Hill, NC 27599, USA

On page 2112, under the title **Antiinflammatory activity**, the third line should read:

As can be seen in Table 2, at 0.03–0.1µg/ml, compounds 1–3 had a significant inhibitory effect on neutrophil degranulation induced by FMLP.

Page 2113, Table 1, line 11. The data for compound 2 at 0.5μg/ml need to be shifted one column to the right. Therefore, the following values should appear under the headings AA, Collagen and PAF, respectively:

 $40.5 \pm 16.6^{**}$ ,  $0.0 \pm 0.0^{***}$  and  $73.7 \pm 7.3^{*}$ .

<sup>1</sup>PII of original article: S0968-0896(97)00133-8

0968-0896/98/\$19.00 © 1998 Elsevier Science Ltd. All rights reserved.

PII: S0968-0896(98)00016-9