



Pergamon

Bioorganic & Medicinal Chemistry 9 (2001) 2213

BIOORGANIC &
MEDICINAL
CHEMISTRY

Corrigendum

Corrigendum to “a Comparative Study on the Inhibition of Human and Bacterial Kynureninase by Novel Bicyclic Kynurenine Analogues”: [Bioorg. Med. Chem. 9 (2001) 983][†]

Deirdre H. Fitzgerald,[†] Karen M. Muirhead and Nigel P. Botting*

School of Chemistry, University of St. Andrews, St Andrews, Fife KY16 9ST, UK

In the paper, the K_m and V_{max} values for human kynureninase were given wrongly. The correct data, determined at 37°C and pH 7.5, were for bacterial kynureninase K_m was found to be $44.2 \pm 1.1 \mu\text{M}$ and

V_{max} $1.45 \pm 0.09 \mu\text{mol/min/mg}$ for L-kynurenine. Values obtained for human kynureninase were K_m $9.39 \pm 1.78 \mu\text{M}$ and V_{max} $8.45 \pm 1.24 \text{ nmol/mg/min}$ for L-3-hydroxykynurenine.

*Corresponding author. Tel.: +44-334-463-856; fax: +44-334-463-808; e-mail: nbp@st-andrews.ac.uk

[†]Current address: School of Dentistry, Trinity College, Lincoln Place, Dublin 2, Ireland.

PII of original article: S0968-0896(00) 00318-7