



Preparation of Glycosylated Amino Acid Derivatives for Glycoprotein Synthesis by In Vitro Translation System

Shino Manabe,^a Kimitoshi Sakamoto,^a Yoshiaki Nakahara,^{a,b} Masahiko Sisido,^c
Takahiro Hoshaka^c and Yukishige Ito^{a,*}

^a*RIKEN (The Institute of Physical and Chemical Research) and CREST, Japan Science and Technology Corporation (JST),
2-1 Hirosawa, Wako-shi, Saitama 351-0198, Japan*

^b*Department of Applied Biochemistry, Tokai University, Kitakaname 1117, Hiratsuka-shi, Kanagawa 259-1292, Japan*

^c*Department of Bioscience and Biotechnology, Faculty of Engineering, Okayama University, 3-1-1 Tsushimanaka,
Okayama 700-8350, Japan*

Received 3 July 2001; accepted 23 August 2001

Abstract—General preparation of glycosylated amino acylated nucleotide for in vitro peptide synthesis was described. Both *O*-glycosylated amino acyl nucleotides and *C*-glycosylated amino acyl nucleotide were synthesized by choosing the appropriate protecting group. © 2002 Elsevier Science Ltd. All rights reserved.

Introduction

Glycosylation is one of the most prominent post-translational modifications that controls protein conformation, functions, stability, and intra/intercellular trafficking. Due to increased understanding of the biological functions of glycan chains as the recognition signals in various biological events, synthetic effort toward glycoprotein has become a major issue in organic synthesis and chemical biology.¹ Thanks to the recent advances in peptide as well as oligosaccharide synthesis technology, chemical synthesis of relatively complex glycopeptide is now a quite feasible task.² However, it is still considered to be a daunting challenge to produce intact glycoproteins by a purely chemical means (Schemes 1–4).


In vitro protein synthesis has proved to be a powerful method to incorporate non-natural or modified amino acids in a site-specific manner.³ For such purpose, the use of amino acylated suppressor *t*RNA, which recognizes amber nonsense codon, has been explored. Based on this approach, Hecht et al. successfully incorporated glucosylated serine into firefly luciferase,⁴ albeit in modest incorporation efficiency.

An alternative in vitro translation system which utilizes four-base codons, CGGG and AGGU, has been developed.^{5,6} Since these quadruplets has been derived from the minor codons of arginine (CGG and AGG), competition between four-base and three-base should be in favor of the former. With a set of orthogonal four-base codons, incorporation of two different nonnatural amino acids was demonstrated to be possible. Considering the success of incorporating various types of modified amino acids into proteins, exploration of the possibility to produce glycoproteins with four-base codon approach seems to be highly promising (Fig. 1).

Prerequisite to the incorporation of a tailored amino acid residue (Xaa) by in vitro protein synthesis is the preparation of *t*RNA amino acylated at the terminal A residue (*t*RNA-Xaa). It has been demonstrated that *t*RNA-Xaa can be prepared by enzymatic ligation of *t*RNA lacking terminal pCpA [*t*RNA(–CA)] with amino acylated dinucleotide pCpA-Xaa⁷ or pdCpA-Xaa.⁸ As an active ester, cyanomethyl ester has proved to be the most effective in terms of the ease of preparation and chemoselective reactivity toward ribose hydroxy groups.⁹

We report here the preparation of three types of glycosylated amino acid cyanomethyl esters **1**, **2**, and **3**, which were designed as key intermediates for the in vitro production of glycoproteins having novel carbohydrate


*Corresponding author. Tel.: +81-48-467-9430; fax: +81-48-432-4680;
e-mail: yukito@postman.riken.go.jp




Results and Discussion

Adequate choice of protecting group for hydroxy groups of sugar part and amino group of amino acid is crucial for preparation of glycosylated aminoacyl pdCpA because of the sensitivity of the ester linkage between pdCpA and amino acid in the presence of nucleophile.¹¹ While Hecht et al. adopted the allyloxy-carbonyl group for hydroxy protection, we focused our attention on the use of acid labile groups. In this context, the ester linkage was shown to be tolerant under *N*-Boc (*tert*-butoxycarbonyl) deprotection conditions (TFA).¹² It was then presumed that either the silyl or acetonide group can be adopted for the protection of carbohydrate components. Simultaneous removal of *N*- and *O*-protecting groups should then be possible without deteriorating the ester linkage.


As the glycosylated amino acid residue to be incorporated by in vitro system, serine linked *N*-acetyl-D-glucosamine (*O*-GlcNAc-Ser), serine linked D-mannose (*O*-Man-Ser), and *C*-linked D-mannosyl tryptophan (*C*-Man-Trp) were selected because of their biological significance.




Scheme 1. Reagent and conditions. (i) TMSOTf, 1,2-dichloroethane, 50 °C, 10 h, 64%; (ii) H₂, 10% Pd/C, EtOAc, Boc₂O, rt, 3 h, 91%; (iii) NaOMe, MeOH, rt, 3 h, TBSCl, imidazole, DMF, then TESCl, imidazole, DMF, 66%; (iv) LiOH, MeOH, H₂O, then bromoacetonitrile, *i*-Pr₂NEt, CH₃CN, rt, overnight, 41%.



Scheme 2. Reagent and conditions: (i) AgOTf, CH₂Cl₂, –10 °C, overnight, 56%; (ii) 10% Pd/C, Boc₂O, MeOH, rt, overnight, 92%; (iii) NaOMe, MeOH, then TESCl, pyridine, rt, overnight, 66%; (iv) LiOH, H₂O, MeOH, rt, 3 h, 84%; (v) bromoacetonitrile, *i*-Pr₂NEt, CH₃CN, rt, overnight, 84%.



Scheme 3. Reagent and conditions: (i) 2,2-dimethoxypropane, PPTS, DMF, rt, 2 days, 74%; (ii) 10% NaOH aq, EtOH, rt-reflux, overnight; (iii) bromoacetonitrile, *i*-Pr₂NEt, CH₃CN, rt, overnight, 80% (two steps).



Scheme 4. Reagent and conditions: (i) DMF, 0 °C, **15a**, 44%, **16a**, 50%, **17a**, 53%; (ii) TFA for **15b**, 42%, and **16b**, quant for **17b**, 4 M HCl dioxane, H₂O, rt, 65%.

For instance, *O*-Linked *N*-acetyl-D-glucosamine is found in a variety of nuclear and cytoplasmic proteins and functions in signal transduction.¹³ *O*-Mannosyl linkage in α -dystroglycan may play a significant role in binding to laminin, which influences development and regeneration of the nervous system.¹⁴ *C*-linked mannosyl tryptophan is a naturally occurring novel subclass of glycoprotein, which is synthesized by our group for the first time.¹⁵ The biological role of mannosylated tryptophan is not clear at the present time, but recent study suggests that *C*-glycosylation is not a rare post-translational modification as previously thought.¹⁶ Not only from human, *C*-Mannosylated tryptophan is also obtained from marine ascidians.¹⁷

Oxazoline **4** was glycosylated in the presence of TMSOTf with serine derivative **5**, which was synthesized from serine methyl ester by the action of TfN₃.¹⁸ After transformation of azide group to Boc in one pot,

the acetyl group was removed. The hydroxy groups were protected as TBS and TES group in order that the deprotection at the final stage can be performed under acidic conditions. After hydrolysis of methyl ester under alkaline conditions, the acid was converted to cyanomethyl ester **1**. Similarly, α -*O*-linked D-mannosyl serine cyanomethyl ester **2** was prepared from bromide **7** in five steps. β -Elimination during the removal of acetate could be minimized (< 3%) by controlling the NaOMe concentration (0.03%).

Mannosyl tryptophan cyanomethyl ester **3** was synthesized as below. Tetraol **10**¹⁵ was protected as its corresponding diacetonide by 2,2-dimethoxypropane in the presence of PPTS without affecting the Boc group. After hydrolysis of methyl ester and sulfone amide at indole nitrogen, carboxylic acid was changed to cyanomethyl ester **3** in 80% yield.

Subsequently, condensation of cyanomethyl ester **1**, **2**, **3** and nucleotide was investigated. The condensation of **1** with AMP tetrabutylammonium salt **11** as a model of pdCpA was performed in 44% yield at 0 °C to reduce the amount of β -elimination of glycosylated amino ester under basic conditions. Similarly, condensation of **2** and **11** was performed at 0 °C in DMF to afford **13a** in 50% yield. When **3** was used as the glycosylated amino acid source, the reaction proceeded relatively slowly and required a high temperature (60 °C), probably because of steric hindrance of the acetonide. Deprotection of silyl group and Boc group of **12a** and **13a** were performed in TFA at the same time in a short period and **12b** and **13b** were obtained. Because TFA treatment of **14a** caused decomposition of mannosyl tryptophan moiety, acidic hydrolysis of acetonide and Boc was performed in 4 M aq HCl in dioxane to afford **14b**.

Conclusion

In conclusion, glycosylated amino acid derivatives for in vitro synthesis of glycoproteins were synthesized. Although incorporation of a glucose residue into firefly luciferase was demonstrated successfully by Hecht et al., the generality of in vitro production of glycoproteins is not clear. Since synthetic routes are now established for *O*-linked- β -D- (**1**), *O*-linked- α -D- (**2**) and *C*-linked- α -D- (**3**) glycosylated amino acids, comparative studies on incorporation efficiencies of these extremely different structures would reveal the scope of the in vitro system for the synthesis of biologically relevant glycoproteins.

Experimental

General procedures

¹H NMR spectra were taken by JEOL EX-400 as solutions in CDCl₃ otherwise mentioned. Chemical shifts are expressed in ppm downfield from the signal for Me₄Si. Optical rotations were measured by Jasco DIP-310 at ambient temperature. Melting points were uncorrected. Kanto silica gel (spherical, neutral, 100–210 μ m) were

used for column chromatography. Merck TLC (silica gel 60F254) was used for TLC analysis. Molecular sieves were activated by heating to 180 °C just before use.

Methyl (2S)-2-azide-3-hydroxypropanoate (5). To a mixture of NaN₃ (66.7 g, 1.0 mol) in H₂O (160 mL) and CH₂Cl₂ (240 mL), Tf₂O (66.7 g, 0.23 mol) was added with vigorous stirring at 0 °C. The mixture was stirred for 4 h at 0 °C. The organic layer was separated, and the aqueous layer was extracted with CH₂Cl₂ (70 mL × 2). The combined organic layers were washed with satd NaHCO₃ (120 mL) and H₂O (120 mL). The organic layer was dried over Na₂SO₄. To a solution of L-serine methyl ester-HCl salt (25.8 g, 186 mmol) and 4-(dimethylamino)pyridine (112 g, 0.92 mol) in MeCN (840 mL), TfN₃ solution (350 mL, as prepared above) was added at 0 °C, and stirred at 0 °C for 2 h. The mixture was diluted with 1,2-dichloroethane (500 mL) and concentrated at 30 °C in vacuo to ca. 300 mL, and filtered. The mother liquid was subjected to silica gel column chromatography (hexane/EtOAc = 6:4) to afford **5** (15.0 g, 62%) as a pale yellow oil: $[\alpha]_D^{25}$ –92.2 (*c* 1.0, CHCl₃); ¹³C NMR (100 MHz, CDCl₃) δ 169.0 (C), 63.4 (CH), 62.7 (CH₂), 52.9 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 4.11 (dd, 1H, *J* = 4.4, 5.4 Hz), 3.94 (d, 1H, *J* = 4.4 Hz), 3.94 (d, 1H, *J* = 5.4 Hz), 3.85 (s, 3H), 2.11 (brs, 1H); HRMS (FAB), *m/z* 146.0567 (*M* + H)⁺, (C₄H₈N₃O₃ requires 146.0566).

3-O-(2-Acetamido-2-deoxy-3,4,6-tri-O-acetyl-β-D-glucopyranosyl)-(2S)-2-azide-3-hydroxypropionic acid methyl ester (6). To a solution of oxazoline **4** (15.2 g, 46.2 mmol), azide **5** (13.4 g, 92.4 mmol) and 1,1,3,3-tetramethylurea (5.5 mL, 46.2 mmol) in 1,2-dichloroethane (100 mL) was added TMSOTf (8.8 mL, 48.5 mmol) dropwise at 0 °C. After addition, the mixture was stirred at 50 °C for 10 h under N₂ atmosphere. After the reaction mixture was cooled to 0 °C, MeOH (10 mL) and Et₃N (6.8 mL, 48.5 mmol) were added. The solvent was evaporated, and the residue was purified by silica gel column chromatography (hexane/AcOEt/MeOH = 8:9:0.7) to afford **6** (14.1 g, 64%) as a colorless solid: mp 135–137 °C; $[\alpha]_D^{25}$ –22.4 (*c* 1.0, CHCl₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.3 (C), 170.3 (C), 169.0 (C), 168.1 (C), 100.1 (CH), 72.0 (CH), 71.7 (CH), 68.6 (CH), 68.5 (CH₂), 62.0 (CH₂), 61.3 (CH), 54.3 (CH), 52.8 (CH₃), 23.2 (CH₃), 20.7 (CH₃), 20.6 (CH₃), 20.6 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 5.52 (d, 1H, *J* = 8.8 Hz), 5.32 (dd, 1H, *J* = 9.3, 10.5 Hz), 5.07 (t, 1H, *J* = 9.3 Hz), 4.83 (d, 1H, *J* = 8.3 Hz), 4.24 (m, 1H), 4.14 (dd, 1H, *J* = 2.4, 12.2 Hz), 4.04 (t, 1H, *J* = 4.0 Hz), 3.90 (dd, 1H, *J* = 3.7, 11.0 Hz), 3.82 (s, 3H), 3.80–3.75 (m, 1H), 3.71 (m, 1H), 2.09 (s, 3H), 2.03 (s, 3H), 2.03 (s, 3H), 1.98 (s, 3H). Anal. calcd for C₁₈H₂₆N₄O₁₁: C, 45.57; H, 5.52; N, 11.81. Found: C, 45.33; H, 5.45; N, 11.60.

N-(tert-Butoxycarbonyl)-3-O-(2-acetamido-2-deoxy-3,4,6-tri-O-acetyl-β-D-glucopyranosyl)-S-serine methyl ester. To a solution of azide **6** (10.6 g, 22.3 mmol) and di-*tert*-butyl dicarbonate (5.9 g, 27 mmol) in EtOAc (200 mL) was added a suspension of 10% Pd/C (1.0 g) in EtOAc (20 mL). The mixture was stirred vigorously under H₂ atmosphere for 3 h. The catalyst was filtered

off through Celite and the solvent was evaporated. The residue was purified by silica gel column chromatography (CHCl₃/MeOH = 98:2) to afford Boc derivative (11.1 g, 91%) as an amorphous: $[\alpha]_D^{25}$ –3.5 (*c* 1.1, CHCl₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.6 (C), 170.4 (C), 170.3 (C), 170.1 (C), 169.1 (C), 155.2 (C), 100.9 (CH), 80.1 (C), 72.1 (CH), 71.9 (CH), 69.2 (CH₂), 68.4 (CH), 62.0 (CH₂), 54.5 (CH), 53.9 (CH), 52.7 (CH₃), 28.4 (CH₃), 23.4 (CH₃), 20.8 (CH₃), 20.8 (CH₃), 20.7 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 5.74 (d, 1H, *J* = 8.1 Hz), 5.43 (d, 1H, *J* = 7.6 Hz), 5.27 (dd, 1H, *J* = 9.3, 10.5 Hz), 5.05 (t, 1H, *J* = 9.6 Hz), 4.71 (d, 1H, *J* = 8.3 Hz), 4.41 (m, 1H), 4.30–4.18 (m, 2H), 4.12 (dd, 1H, *J* = 2.3, 12.3 Hz), 3.88–3.80 (m, 2H), 3.76 (s, 3H), 3.65 (m, 1H), 2.10 (s, 3H), 2.04 (s, 3H), 2.02 (s, 3H), 1.98 (s, 3H), 1.46 (s, 9H). Anal. calcd for C₂₃H₃₆N₂O₁₃: C, 50.36; H, 6.62; N, 5.11. Found: C, 50.23; H, 6.61; N, 5.13.

(N-tert-Butoxycarbonyl)-3-O-[2-acetamido-2-deoxy-3,4-bis-O-triethylsilyl-6-(tert-butyldimethylsilyl)-β-D-glucopyranosyl]-S-serine methyl ester. To a solution of acetate (8.2 g, 14.9 mmol) in dry MeOH (150 mL) was added MeONa (0.1% wt MeOH solution, 30 mL) at room temperature. The mixture was stirred for 3 h at room temperature. The mixture was neutralized with Amberlyst 15-E. The resins were filtered off and the solvent was evaporated. The material was pumped to dryness. The residue was dissolved in DMF (36 mL) and imidazole (6.1 g, 90 mmol) was added. To the solution was added *t*-butyldimethylsilyl chloride (2.0 g, 13.0 mmol) at room temperature. After the mixture was stirred for 2 h, triethylsilyl chloride (4.6 mL, 27.5 mmol) was added at 0 °C and the mixture was stirred for 12 h at room temperature. To the solution was added MeOH (4 mL) and stirred for 30 min to destroy the excess of silyl chloride. The reaction mixture was diluted with EtOAc, and H₂O was added to the solution. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with aq 5% citric acid, satd. NaHCO₃ aq, and brine. The organic layer was dried over Na₂SO₄. The solvent was evaporated and the residue was purified by silica gel column chromatography (10–30% EtOAc in hexane) to afford silyl ether (7.5 g, 66%, 2 steps) as a colorless solid: mp 108–109 °C; $[\alpha]_D^{25}$ –9.1 (*c* 1.0, benzene); ¹³C NMR (100 MHz, C₆D₆) δ 170.8 (C), 168.9 (C), 155.6 (C), 100.8 (CH), 79.5 (C), 78.4 (CH), 73.4 (CH), 70.8 (CH), 69.6 (CH₂), 63.6 (CH₂), 55.0 (CH), 54.7 (CH), 52.1 (CH₃), 28.7 (CH₃), 26.4 (CH₃), 23.7 (CH₃), 18.8 (C), 7.6 (CH₃), 7.5 (CH₃), 5.6 (CH₂), –4.5 (CH₃), –4.9 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 6.63 (d, 1H, *J* = 9.0 Hz), 5.41 (d, 1H, *J* = 8.3 Hz), 4.61 (d, 1H, *J* = 2.4 Hz), 4.40 (m, 1H), 4.20 (dd, 1H, *J* = 3.8, 9.9 Hz), 4.07–3.95 (m, 2H), 3.88 (m, 1H), 3.71 (s, 3H), 3.75–3.67 (m, 3H), 3.62 (dd, 1H, *J* = 3.9, 10.0 Hz), 1.96 (s, 3H), 1.45 (s, 9H), 1.01–0.94 (m, 18H), 0.90 (s, 9H), 0.69–0.59 (m, 12H), 0.08 (s, 3H), 0.07 (s, 3H). Anal. calcd for C₃₅H₇₂N₂O₁₀Si₃: C, 54.95; H, 9.48; N, 3.66. Found: C, 55.00; H, 9.59; N, 3.74.

(N-tert-Butoxycarbonyl)-3-O-[2-acetamido-2-deoxy-3,4-bis-O-triethylsilyl-6-(tert-butyldimethylsilyl)-β-D-glucopyranosyl]-S-serine cyanomethyl ester (1). The methyl ester (647 mg, 846 μmol) was dissolved in THF (8.4 mL)

and the solution was cooled to 0 °C. To the solution was added ice cooled solution of LiOH (24.3 mg, 1.02 μ mol) in water (2.5 mL). The mixture was stirred at 0 °C for 90 min. The reaction was quenched with aq 10% citric acid, and aqueous layer was extracted with EtOAc three times, then the combined organic layers were washed with brine and dried over Na₂SO₄. The solvent was evaporated, and the residue was purified by silica gel column chromatography (0–4% MeOH in CHCl₃) to afford carboxylic acid (311 mg, 414 μ mol). The obtained acid (152 mg, 202 μ mol) and *i*-Pr₂NEt (37 μ L, 212 μ mol) were dissolved in MeCN (1.0 mL), and bromoacetonitrile (28 μ L, 404 μ mol) was added at room temperature. After stirring for 9 h, the solvent was evaporated. To the residue was added water and aqueous layer was extracted with EtOAc three times, then the combined organic layers were washed with brine and dried over Na₂SO₄. The solvent was evaporated, and the residue was purified by silica gel column chromatography (hexane/EtOAc=7:3) to afford cyanomethyl ester **1** (135 mg, 41%, two steps) as a colorless solid: mp 116–118 °C; $[\alpha]_D^{25}$ –18.9 (*c* 1.0, benzene); ¹³C NMR (100 MHz, C₆D₆) δ 169.4 (C), 169.2 (C), 155.6 (C), 114.4 (C), 100.8 (CH), 80.0 (C), 78.6 (CH), 73.9 (CH), 71.0 (CH), 68.7 (CH₂), 63.6 (CH₂), 55.4 (CH), 54.6 (CH), 48.9 (CH₂), 28.6 (CH₃), 26.4 (CH₃), 23.7 (CH₃), 18.9 (C), 7.5 (CH₃), 7.5 (CH₃), 5.7 (CH₂), 5.7 (CH₂), –4.5 (CH₃), –4.8 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 6.54 (d, 1H, *J*=9.3 Hz), 5.38 (d, 1H, *J*=8.3 Hz), 4.78 (s, 2H), 4.63 (d, 1H, *J*=3.2 Hz), 4.48 (m, 1H), 4.19 (dd, 1H, *J*=4.6, 9.8 Hz), 3.94 (m, 2H), 3.85–3.79 (m, 2H), 3.74–3.65 (m, 3H), 1.96 (s, 3H), 1.46 (s, 9H), 1.01–0.94 (m, 18H), 0.91 (s, 9H), 0.69–0.59 (m, 12H), 0.08 (d, 6H, *J*=4.6 Hz). Anal. calcd for C₃₆H₇₁N₃O₁₀Si₃: C, 54.72; H, 9.06; N, 5.32. Found: C, 54.77; H, 9.18; N, 5.23.

***N*-(Benzyloxycarbonyl)-3-*O*-(2,3,4,6-tetra-*O*-acetyl- α -D-mannopyranosyl)-*S*-serine methyl ester (**9**).** To a suspension of L-Z-Ser-OMe **8** (7.76 g, 30.64 mmol), AgOTf (10.34 g, 40.24 mmol) and activated molecular sieves 4A (12.0 g) in CH₂Cl₂ (100 mL) was added a solution of bromide **7** (16.47 g, 40.05 mmol) in CH₂Cl₂ (30 mL) at –20 °C under Ar. The mixture was stirred at –10 °C overnight. The reaction mixture was quenched with triethylamine (8 mL). The mixture was filtered through Celite, and the filter cake was washed with CH₂Cl₂ (30 mL $\times$ 2). After removal of the solvent, the residue was directly subjected to silica gel column chromatography (CHCl₃/EtOAc 4:1–1:1) to afford **9** (9.99 g, 56%) as an amorphous: $[\alpha]_D^{25}$ 48.2 (*c* 1.1, CHCl₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.2 (C), 169.6 (C), 169.5 (C), 169.4 (C), 169.3 (C), 155.5 (C), 135.8 (C), 128.3 (CH), 128.0 (CH), 98.1 (CH), 69.2 (CH₂), 69.1 (CH), 69.0 (CH), 68.6 (CH), 67.1 (CH₂), 65.8 (CH), 62.2 (CH₂), 54.2 (CH), 52.8 (CH₃), 20.8 (CH₃), 20.7 (CH₃), 20.6 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.28 (m, 5H), 5.77 (d, 1H, *J*=7.8 Hz), 5.26–5.22 (m, 2H), 5.18 (m, 1H), 5.14 (s, 2H), 4.79 (d, 1H, *J*=1.5 Hz), 4.57 (m, 1H), 4.23 (dd, 1H, *J*=5.6, 12.2 Hz), 4.09 (dd, 1H, *J*=2.2, 12.2 Hz), 4.04–3.92 (m, 3H), 3.81 (s, 3H), 2.15 (s, 3H), 2.08 (s, 3H), 2.03 (s, 3H), 1.99 (s, 3H). Anal. calcd for C₂₆H₃₃N₃O₁₄: C, 53.51; H, 5.70; N, 2.4. Found: C, 53.65; H, 5.80; N, 2.23.

***N*-(*tert*-Butoxycarbonyl)-3-*O*-(2,3,4,6-tetra-*O*-acetyl- α -D-mannopyranosyl)-*S*-serine methyl ester.** To a solution of **9** (6.55 g, 11.22 mmol) and di-*tert*-butyl dicarbonate (3.19 g, 14.62 mmol) in MeOH (150 mL) was added 10% Pd/C (800 mg). The mixture was stirred vigorously under H₂ atmosphere for 1 h. The mixture was diluted with EtOAc (200 mL), and the catalyst was filtered off through Celite and the catalyst was washed with EtOAc. After the solvent was evaporated, the residue was purified by silica gel column chromatography (hexane/EtOAc=6:4) to afford Boc derivative (5.65 g, 92%) as an amorphous: $[\alpha]_D^{25}$ 48.3 (*c* 1.0, CHCl₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.3 (C), 170.1 (C), 169.6 (C), 169.5 (C), 169.3 (C), 155.0 (C), 98.2 (CH), 80.2 (C), 69.3 (CH), 69.2 (CH₂), 68.9 (CH), 68.8 (CH), 65.9 (CH), 62.2 (CH₂), 53.8 (CH), 52.8 (CH₃), 28.4 (CH₃), 20.9 (CH₃), 20.8 (CH₃), 20.8 (CH₃), 20.7 (CH₃); ¹H NMR (400 MHz, CDCl₃) δ 5.47 (d, 1H, *J*=7.8 Hz), 5.31–5.25 (m, 2H), 5.18 (dd, 1H, *J*=1.7, 2.9 Hz), 4.80 (d, 1H, *J*=1.5 Hz), 4.50 (m, 1H), 4.28 (dd, 1H, *J*=5.2, 12.3 Hz), 4.12 (dd, 1H, *J*=2.3, 12.3 Hz), 4.00–3.92 (m, 3H), 3.80 (s, 3H), 2.15 (s, 3H), 2.12 (s, 3H), 2.05 (s, 3H), 1.99 (s, 3H), 1.47 (s, 9H). Anal. calcd for C₂₃H₃₅N₃O₁₄: C, 50.27; H, 6.42; N, 2.55. Found: C, 50.33; H, 6.47; N, 2.48.

***N*-(*tert*-Butoxycarbonyl)-3-*O*-(2,3,4,6-tetrakis-*O*-triethylsilyl- α -D-mannopyranosyl)-*S*-serine methyl ester.** To a solution of acetate (1.15 g, 2.09 mmol) in dry MeOH (30 mL) was added MeONa (0.1% wt MeOH solution, 2 mL) at room temperature in the presence of phenolphthalein as an indicator. The mixture was stirred for 2 h and then neutralized with Amberlyst 15E. The resins were filtered and the solvent was evaporated. The residue was dissolved in pyridine (30 mL) and chlorotriethylsilane (2.5 mL, 14.63 mmol) was added at 0 °C and the mixture was stirred for 8 h at room temperature. To the solution was added MeOH (3 mL) and stirred for 30 min to destroy the excess of silyl chloride, and the solvent was evaporated. The residue was dissolved into EtOAc and satd NH₄Cl. The aqueous layer was extracted with EtOAc, and the combined organic layers were washed with brine. The organic layer was dried over Na₂SO₄. The solvent was evaporated and the residue was purified by silica gel column chromatography (hexane/EtOAc=9:1) to afford silyl ether (1.48 g, 84%, 2 steps) as a colorless oil: $[\alpha]_D^{25}$ 33.9 (*c* 1.0, benzene); ¹³C NMR (100 MHz, C₆D₆, 70 °C) δ 170.7 (C), 155.3 (C), 101.6 (CH), 79.6 (C), 77.3 (CH), 75.5 (CH), 73.7 (CH), 70.1 (CH), 68.4 (CH₂), 63.1 (CH₂), 55.0 (CH), 51.8 (CH₃), 28.7 (CH₃), 7.6 (CH₃), 7.5 (CH₃), 7.3 (CH₃), 7.3 (CH₃), 6.2 (CH₂), 6.1 (CH₂), 5.9 (CH₂), 5.5 (CH₂); ¹H NMR (400 MHz, C₆D₆, 70 °C) δ 5.44 (brs, 1H), 4.71 (d, 1H, *J*=3.2 Hz), 4.50 (brs, 1H), 4.21 (t, 1H, *J*=7.2 Hz), 4.03–3.91 (m, 5H), 3.81 (m, 1H), 3.61 (brs, 1H), 3.38 (s, 3H), 1.44 (s, 9H), 1.13–1.02 (m, 36H), 0.83–0.68 (m, 24H). Anal. calcd for C₃₉H₈₃N₃O₁₀Si₄: C, 55.87; H, 9.98; N, 1.67. Found: C, 55.79; H, 10.11; N, 1.76.

***N*-(*tert*-Butoxycarbonyl)-3-*O*-(2,3,4,6-tetrakis-*O*-triethylsilyl- α -D-mannopyranosyl)-*S*-serine.** To a solution of methyl ester (2.77 g, 3.31 mmol) in THF (50 mL) was added a solution of LiOH (95 mg, 3.97 mmol) in H₂O

(10 mL) at room temperature. The mixture was stirred for 3 h, and diluted with CH_2Cl_2 . The reaction was quenched with aq 10% citric acid, and the aq layer was extracted with CH_2Cl_2 . Then the combined organic layers were washed with brine and dried over Na_2SO_4 . The solvent was evaporated, and the residue was purified by silica gel column chromatography (hexane/EtOAc 7:1–1:1) to afford the acid (2.29 g, 84%). ^{13}C NMR ($\text{DMSO}-d_6$, 100°C) δ 170.5 (C), 156.9 (C), 99.6 (CH), 77.8 (C), 75.6 (CH), 73.7 (CH), 72.0 (CH), 66.7 (CH_2), 61.5 (CH_2), 60.6 (CH), 53.8 (CH), 27.6 (CH_3), 6.1 (CH_3), 6.0 (CH_3), 6.0 (CH_3), 5.9 (CH_3), 5.9 (CH_3), 5.4 (CH_2), 4.5 (CH_2), 4.4 (CH_2), 4.3 (CH_2), 3.9 (CH_2). ^1H NMR ($\text{DMSO}-d_6$, 100°C) δ 5.68 (1H, bs), 4.09 (1H, d, $J=2.8$ Hz), 3.65 (1H, m), 3.46 (1H, t, $J=7.6$ Hz), 3.3–3.15 (7H, m), 2.93 (1H, m), 0.91 (9H, s), 0.50–0.41 (36H, m), 0.18–0.10 (20H, m), –0.01 (4H, q, $J=8.0$ Hz); $[\alpha]_D^{24} +28.3$ (c 0.58, CHCl_3); HRMS (FAB), m/z 824.5007 ($\text{C}_{38}\text{H}_{82}\text{O}_{10}\text{NSi}_4$ requires 824.5016).

***N*-(tert-Butoxycarbonyl)-3-*O*-(2,3,4,6-tetrakis-*O*-triethylsilyl- α -D-mannopyranosyl)serine cyanomethyl ester (2).** The above prepared acid (1.99 g, 2.41 mmol) and *i*-Pr₂NEt (0.84 mL, 4.82 mmol) were dissolved in 6:1 MeCN/DMF (35 mL), and bromoacetonitrile (0.58 mL, 4.82 mmol) was added at 4°C . After stirring the mixture for 1 day at room temperature, solvent was evaporated. To the residue was added H_2O , and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . The solvent was evaporated, and the residue was purified by silica gel column chromatography (hexane/EtOAc=9:1–4:1) to afford cyanomethyl ester **2** (1.75 g, 84%) as a colorless oil: $[\alpha]_D^{25} 29.7$ (c 1.1, benzene); ^{13}C NMR (100 MHz, C_6D_6 , 70°C) δ 169.2 (C), 155.3 (C), 113.9 (C), 101.8 (CH), 80.1 (C), 77.7 (CH), 75.6 (CH), 73.3 (CH), 70.3 (CH), 68.2 (CH_2), 63.2 (CH_2), 54.9 (CH), 48.6 (CH_2), 28.6 (CH_3), 7.6 (CH_3), 7.4 (CH_3), 7.3 (CH_3), 7.3 (CH_3), 6.2 (CH_2), 6.0 (CH_2), 5.9 (CH_2), 5.5 (CH_2); ^1H NMR (400 MHz, $\text{DMSO}-d_6$, 80°C) δ 6.83 (brs, 1H), 4.95 (s, 2H), 4.77 (d, 1H, $J=3.2$ Hz), 4.59 (d, 1H, $J=3.2$ Hz), 4.32 (m, 1H), 3.95 (t, 1H, $J=7.3$ Hz), 3.86 (dd, 1H, $J=4.8$, 10.6 Hz), 3.79–3.69 (m, 5H), 3.41 (m, 1H), 1.39 (s, 9H), 0.99–0.91 (m, 36H), 0.67–0.55 (m, 24H). Anal. calcd for $\text{C}_{40}\text{H}_{82}\text{N}_2\text{O}_{10}\text{Si}_4$: C, 55.64; H, 9.57; N, 3.24. Found: C, 55.36; H, 9.69; N, 3.21.

***N*-(tert-Butoxycarbonyl)-[2,3,4,6-bis-*O*-(1-methylethylidene)- α -D-mannopyranosyl]-(1-phenylsulfonyl)-tryptophan cyanomethyl ester.** To a solution of tetraol **10** (121.0 mg, 0.195 mmol) and 2,2-dimethoxypropane (0.25 mL) in DMF (0.7 mL), PPTS (19 mg) was added at room temperature. The mixture was stirred at room temperature for 2 days. Triethylamine (0.3 mL) was added to quench PPTS, and stirred for 15 min. After the mixture was diluted with CHCl_3 , purified by silica gel column chromatography directly ($\text{CHCl}_3/\text{MeOH}$ 7:3). The material recrystallized from EtOAc–hexane to afford the diacetone (100.1 mg, 74%): $[\alpha]_D^{24} -11$ (c 0.47, CHCl_3); mp $142\text{--}143^\circ\text{C}$; ^{13}C NMR (67.8 MHz) δ 172.83 (C), 155.40 (C), 133.11 (CH), 128.75 (CH), 122.90 (CH), 125.71 (CH), 123.89 (CH), 120.16 (C), 119.42 (C), 115.50 (CH), 110.63 (C), 99.36 (C), 79.77 (C), 76.50 (CH), 71.96 (CH), 70.60

(CH_2), 64.82 (CH), 62.83 (CH_2), 53.22 (CH), 52.28 (CH_3), 29.05 (CH_3), 28.51 (CH_3), 27.27 (CH_2), 26.54 (CH_3), 24.35 (CH_3), 19.24 (CH_3). Anal. calcd for $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_{11}\text{S}$: C, 59.99; H, 6.33; N, 4.00. Found: C, 59.92; H, 6.33; N, 3.92.

***N*-(tert-Butoxycarbonyl)-[2,3,4,6-bis-*O*-(1-methylethylidene)- α -D-mannopyranosyl]-tryptophan cyanomethyl ester (3).** To the suspension of the above methyl ester (107 mg, 0.153 mmol) in EtOH (10 mL), 10% NaOH aq (1.0 mL) was added at room temperature. The mixture was stirred at room temperature for 3 h until the methyl ester was consumed on TLC. After starting material was consumed, the mixture became homogenous. Then the mixture was refluxed overnight. After cooling the mixture to room temperature, the solvent was evaporated, and the residue was diluted with CHCl_3 . The mixture was washed with aq 10% citric acid. The aq layer was extracted with CHCl_3 . The combined layers were washed with brine. After removal of solvent in vacuo, the residue was pumped. The residue was dissolved in CH_3CN (1 mL), then *i*-Pr₂NEt (53 μL , 0.306 mmol) and bromoacetonitrile (21 μL , 0.306 mmol) were added at room temperature. The mixture was stirred at room temperature over night. The mixture was diluted with CHCl_3 and brine was added. The aq layer was extracted with CHCl_3 , then the combined layers were dried over Na_2SO_4 . After evaporation, the residue was purified by silica gel column chromatography (hexane/EtOAc 3:2). The material was washed with hexane to give the cyanomethyl ester **3** (71.2 mg, 80%): $[\alpha]_D^{24} +103$ (c 1.25, PhH), ^{13}C NMR (C_6D_6 , C_6D_6 as 128 ppm) 172.06 (C), 155.84 (C), 136.23 (C), 133.38 (C), 123.01 (CH), 120.56 (CH), 119.32 (CH), 111.58 (CH), 110.96 (C), 108.18 (C), 99.62 (C), 79.55 (C), 76.86 (CH), 76.41 (CH), 72.13 (CH), 69.18 (CH), 66.14 (CH), 63.63 (CH_2), 54.45 (CH), 48.48 (CH_2), 29.54 (CH_3), 28.55 (CH_3), 27.22 (CH_3), 27.03 (CH_2), 24.54 (CH_3), 19.25 (CH_3). ^1H NMR δ 8.29 (s, 1H), 7.63 (d, 1H, $J=8.0$ Hz), 7.37 (d, 2H, $J=8.0$ Hz), 7.3–7.1 (m, 2H), 6.30 (d, 1H, $J=4.8$ Hz), 4.92–4.86 (m, 2H), 4.74–4.69 (m, 2H), 4.50–4.45 (m, 2H), 4.36–4.34 (m, 1H), 4.02 (dd, 1H, $J=10.8$ Hz, 6.2 Hz), 3.84–3.71 (m, 2H), 3.38 (d, 1H, $J=14.0$ Hz, 3.6 Hz), 3.00 (t, 1H, $J=11.6$ Hz), 1.71 (s, 3H), 1.71 (s, 3H), 1.60 (s, 3H), 1.49 (s, 3H), 1.40 (s, 3H), 1.34 (s, 9H). Anal. calcd for $\text{C}_{30}\text{H}_{39}\text{N}_3\text{O}_9$: C, 61.53; H, 6.71; N, 7.18. Found: C, 61.30; H, 6.80; N, 7.00.

3-*O*-[2-Acetamido-2-deoxy-3,4-bis-*O*-triethylsilyl-6-(tert-butyl-dimethylsilyl)- β -D-glucopyranosyl]-*N*-(tert-butoxycarbonyl)-*S*-serine AMP ester (12a). AMP-tetra-*n*-butylammonium salt **11** (97.6 mg, 138 μmol) was dissolved in DMF (1.5 mL) and cooled to 0°C under Ar atmosphere. To the solution was added pre-cooled (0°C) solution of cyanomethyl ester **1** (432 mg, 547 μmol) in DMF (0.8 mL). The mixture was stirred for 2 h at 0°C , and then 5% (v/v) acetic acid in water (221 μL) was added to neutralize the mixture. The mixture was washed with hexane (8 mL $\times$ 8) to remove the excess of cyanomethyl ester. The DMF solution was diluted with 30% MeCN in water (12 mL), and the suspension was subjected to C_{18} reverse phase column chromatography (Waters Sep-Pak Vac C_{18} 5 g/12 cc, MeCN/ H_2O =3:7 $\rightarrow$ 1:1 $\rightarrow$ 8:2). Fractions containing target molecule were

diluted with three volumes of MeCN. To the solution was added 5% (v/v) acetic acid in water (158 μ L) and the solution was evaporated at 30 °C for avoiding β -elimination and saponification. The residue was pumped to give **12a** (95.5 mg, 44%). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:10. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ^{13}C NMR (100 MHz, DMSO- d_6) δ 169.8 (C), 169.2 (C), 155.8 (C), 155.1 (C), 152.3 (CH), 150.0 (C), 139.1 (CH), 118.3 (C), 100.7 (CH), 85.0 (CH), 82.8 (CH), 78.5 (C), 76.3 (CH), 75.9 (CH), 75.7 (CH), 72.5 (CH), 71.4 (CH), 67.8 (CH₂), 63.6 (CH₂), 61.7 (CH₂), 57.5 (CH), 55.4 (CH), 54.1 (CH), 28.1 (CH₃), 25.7 (CH₃), 23.2 (CH₃), 23.1 (CH₂), 19.2 (CH₂), 18.1 (C), 13.5 (CH₃), 6.9 (CH₃), 6.9 (CH₃), 4.9 (CH₂), 4.7 (CH₂), -4.8 (CH₃), -5.6 (CH₃); ^1H NMR (400 MHz, DMSO- d_6) δ 8.58 (s, 1H), 8.11 (s, 1H), 8.01 (d, 1H, J = 9.3 Hz), 7.22 (br s, 2H), 6.85 (d, 1H, J = 8.5 Hz), 5.98 (d, 1H, J = 7.8 Hz), 5.37 (d, 1H, J = 4.4 Hz), 5.11 (m, 1H), 4.49 (d, 1H, J = 7.8 Hz), 4.32 (m, 1H), 4.06 (br s, 1H), 3.93 (m, 1H), 3.85 (m, 1H), 3.76 (m, 2H), 3.68 (m, 1H), 3.60 (m, 1H), 3.50 (t, 1H, J = 8.5 Hz), 3.40 (m, 1H), 3.15 (m, 9H), 1.83 (s, 3H), 1.55 (m, 8H), 1.40 (s, 9H), 1.29 (m, 8H), 0.95–0.80 (m, 30H), 0.83 (s, 9H), 0.66 (q, 6H, J = 8.0 Hz), 0.57 (q, 6H, J = 7.8 Hz), 0.05 (s, 3H), 0.04 (s, 3H); HRMS (FAB), m/z 1080.4940 ($M + \text{H}$)⁺, (C₄₄H₈₃N₇O₁₆PSi₃ requires 1080.4942).

3-O-(2-Acetamido-2-deoxy- β -D-glucopyranosyl)-S-serine AMP ester (12b). To a test tube containing **12a** (34.7 mg, 22.0 μ mol) was added pre-cooled (0 °C) TFA (3.0 mL), and the solution was stirred for 1 h at 0 °C. TFA was removed by blowing N₂ gas. The pellet remained was washed with ether (4 mL $\times$ 3) and dried in vacuo. The precipitate was dissolved in 3% MeCN in water (2.0 mL), and the solution was subjected to C₁₈ reverse phase column chromatography (Waters Sep-Pak Vac C₁₈ 10 g/35 cc, MeCN/H₂O = 3:97 $\rightarrow$ 1:9). To avoid saponification, column was performed at 4 °C. Fractions containing target molecule were diluted with 10 volumes of MeCN and evaporated at 30 °C, and the residue was pumped to give **12b** (5.9 mg, 42%). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:5. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ^{13}C NMR (100 MHz, DMSO- d_6) δ 169.8 (C), 166.7 (C), 155.7 (C), 152.5 (CH), 149.4 (C), 138.9 (CH), 118.5 (C), 100.8 (CH), 86.2 (CH), 80.9 (CH), 76.9 (CH), 75.3 (CH), 74.0 (CH), 72.2 (CH), 70.4 (CH), 66.7 (CH₂), 64.1 (CH₂), 60.9 (CH₂), 55.1 (CH), 53.1 (CH), 23.1 (CH₃); ^1H NMR (400 MHz, DMSO- d_6) δ 8.46 (s, 1H), 8.15 (s, 1H), 8.02 (d, 1H, J = 8.5 Hz), 7.32 (br s, 2H), 6.00 (d, 1H, J = 6.6 Hz), 5.47 (m, 1H), 4.93 (m, 1H), 4.47 (d, 1H, J = 8.3 Hz), 4.34 (br s, 1H), 4.25 (br s, 1H), 4.13 (m, 1H), 4.05 (br d, 1H, J = 9.0 Hz), 3.93 (m, 2H), 3.70 (d, 2H, J = 10.3 Hz), 3.45 (m, 2H), 3.32 (dd, 1H, J = 8.8, 9.8 Hz), 3.20 (m, 1H), 3.09 (dd, 1H, J = 8.8, 9.0 Hz), 1.85 (s, 3H); HRMS (FAB), m/z 638.1839 ($M + \text{H}$)⁺ (C₂₁H₃₃N₇O₁₄P requires 638.1823).

N-tert-Butoxycarbonyl-3-O-(2,3,4,6-tetrakis-O-triethylsilyl- α -D-mannopyranosyl)-S-serine AMP ester (13a). AMP-tetra-*n*-butylammonium salt **11** (111 mg, 156

μ mol) was dissolved in DMF (1.6 mL) and cooled to 0 °C under Ar atmosphere. To the solution was added pre-cooled (0 °C) solution of cyanomethyl ester **2** (525 mg, 608 μ mol) in DMF (1.0 mL). The mixture was stirred for 2 h at 0 °C, and then 5% (v/v) acetic acid in water (250 μ L) was added to neutralize the mixture. The mixture was washed with hexane (8 mL $\times$ 2) to remove the excess of cyanomethyl ester. The DMF solution was diluted with 90% MeCN in water (10 mL), and the solution was subjected to C₁₈ reverse phase column chromatography (Waters Sep-Pak Vac C₁₈ 5 g/12 cc, MeCN/H₂O = 9:17 $\rightarrow$ 1:1 $\rightarrow$ THF/H₂O 1:1). Fractions containing target molecule were diluted with same volume of THF and four volumes of MeCN. To the solution was added 5% (v/v) acetic acid in water (25 μ L) and the solution was evaporated at 30 °C for avoiding β -elimination and saponification. The residue was pumped to give **13a** (110 mg, 50%). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:10. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ^{13}C NMR (100 MHz, DMSO- d_6) δ 169.1 (C), 155.6 (C), 154.9 (C), 152.1 (CH), 150.0 (C), 139.0 (CH), 118.2 (C), 100.5 (CH), 84.9 (CH), 83.1 (CH), 78.3 (C), 76.6 (CH), 74.4 (CH), 74.1 (CH), 73.0 (CH), 72.7 (CH), 67.2 (CH), 66.9 (CH₂), 63.5 (CH₂), 61.3 (CH₂), 57.5 (CH₂), 54.3 (CH), 28.1 (CH₃), 23.1 (CH₂), 19.2 (CH₂), 13.5 (CH₃), 6.9 (CH₃), 6.9 (CH₃), 6.6 (CH₃), 4.9 (CH₂), 4.6 (CH₂), 4.6 (CH₂), 4.2 (CH₂); ^1H NMR (400 MHz, DMSO- d_6) δ 8.63 (s, 1H), 8.11 (s, 1H), 7.21 (br s, 2H), 7.01 (d, 1H, J = 8.1 Hz), 6.00 (d, 1H, J = 8.1 Hz), 5.32 (d, 1H, J = 4.6 Hz), 5.07 (m, 1H), 4.58 (d, 1H, J = 2.2 Hz), 4.43 (m, 1H), 4.00 (m, 2H), 3.90 (m, 1H), 3.83 (m, 1H), 3.82–3.70 (m, 6H), 3.37 (m, 1H), 3.15 (m, 8H), 1.55 (m, 8H), 1.40 (s, 9H), 1.38–1.25 (m, 8H), 0.95–0.83 (m, 48H), 0.65–0.50 (m, 24H); HRMS (FAB), m/z 1153.5546 ($M + \text{H}$)⁺, (C₄₈H₉₄N₆O₁₆PSi₄ requires 1153.5541).

3-O- α -D-Mannopyranosyl-S-serine AMP ester (13b). To a test tube containing **13a** (28.7 mg, 20.3 μ mol) was added pre-cooled (0 °C) TFA (3.0 mL), and the solution was stirred for 1 h at 0 °C. TFA was removed by blowing N₂ gas. The pellet remained was washed with ether (4 mL $\times$ 3) and dried in vacuo. The precipitate was dissolved in 1.5% MeCN in water (2.0 mL), and the solution was subjected to C₁₈ reverse phase column chromatography (Waters Sep-Pak Vac C₁₈ 10 g/35 cc, MeCN/H₂O = 1.5:98.5 $\rightarrow$ 1:9). To avoid saponification, column was performed at 4 °C, fractions containing target molecule were diluted 10 volumes of MeCN and evaporated at 30 °C, and the residue was pumped to give **13b** (12.1 mg, quant). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:5. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ^{13}C NMR (100 MHz, DMSO- d_6) δ 167.3 (C), 155.7 (C), 152.4 (CH), 149.3 (C), 139.0 (CH), 118.6 (C), 101.0 (CH), 86.5 (CH), 80.9 (CH), 74.8 (CH), 74.0 (CH), 71.8 (CH), 70.4 (CH), 69.5 (CH), 67.0 (CH), 65.8 (CH₂), 63.7 (CH₂), 61.1 (CH₂), 52.8 (CH); ^1H NMR (400 MHz, DMSO- d_6) δ 8.44 (s, 1H), 8.16 (s, 1H), 7.31 (br s, 2H), 5.98 (d, 1H, J = 6.3 Hz), 5.47 (m, 1H), 4.95 (t, 1H, J = 6.1 Hz), 4.69 (s, 1H), 4.36 (br s, 1H), 4.31 (d, 1H, J = 3.2 Hz), 4.00–3.90 (m, 4H), 3.69 (m, 2H), 3.57

(m, 1H), 3.46 (m, 1H), 3.39 (m, 2H); HRMS (FAB), m/z 597.1551 ($M+H$)⁺, ($C_{19}H_{30}N_6O_{14}P$ requires 597.1558).

***N*-(*tert*-Butoxycarbonyl)-[2,3,4,6-bis-*O*-(1-methylethylidene)- α -D-mannopyranosyl]-tryptophan AMP ester (**14a**).**

To a solution of AMP-tetra-*n*-butylammonium salt **11** (9.3 mg, 13.1 μ mol) in DMF (190 μ L) was added cyanomethyl ester **3** (37.2 mg, 63.5 μ mol). The mixture was stirred at 40 °C for 14 h, at 60 °C for 8 h, cooled to room temperature and poured into ether (40 mL). The precipitate was collected by centrifuge, and washed with ether (10 mL) twice. The pellet was dried in vacuo and dissolved in 10% MeCN in H₂O (2.0 mL). The solution was subjected to C₁₈ reverse phase column chromatography (Waters Sep-Pak Vac C₁₈ 2 g/12 cc, MeCN/H₂O = 2/8→3/7). The fractions containing target molecule were diluted with five volumes of MeCN and evaporated at 30 °C for avoiding saponification. Then **14a** was obtained as an amorphous (7.9 mg, 53%). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:5. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ¹³C NMR (100 MHz, DMSO-*d*₆) δ 171.5 (C), 155.6 (C), 154.9 (C), 149.9 (C), 139.1 (CH), 135.6 (C), 133.4 (C), 126.7 (C), 121.4 (CH), 118.5 (CH), 118.2 (CH), 118.2 (C), 111.3 (CH), 109.2 (C), 108.3 (C), 98.6 (C), 85.1 (CH), 82.9 (CH), 78.5 (C), 75.8 (CH), 75.5 (CH), 75.1 (CH), 72.5 (CH), 71.2 (CH), 67.6 (CH), 64.8 (CH₂), 63.6 (CH₂), 62.2 (CH₂), 57.4 (CH₂), 54.7 (CH), 28.9 (CH₃), 28.0 (CH₃), 26.9 (CH₃), 24.6 (CH₃), 23.1 (CH₂), 19.2 (CH₂), 13.5 (CH₃); ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.11 (s, 1H), 8.62 (s, 1H), 8.12 (s, 1H), 7.58 (d, 1H, J = 8.1 Hz), 7.36 (d, 1H, J = 8.1 Hz), 7.22 (br s, 2H), 7.10 (t, 1H, J = 7.6 Hz), 7.02 (t, 1H, J = 7.6 Hz), 6.81 (d, 1H, J = 6.8 Hz), 5.97 (d, 1H, J = 7.8 Hz), 5.30 (d, 1H, J = 4.9 Hz), 5.10–5.00 (m, 2H), 4.61 (dd, 1H, J = 7.6, 8.1 Hz), 4.45 (t, 1H, J = 7.1 Hz), 4.45 (m, 1H), 4.30 (dd, 1H, J = 7.6, 10.3 Hz), 3.84–3.54 (m, 4H), 3.54 (m, 1H), 3.27 (dd, 1H, J = 4.6, 14.2 Hz), 3.14 (m, 8H), 3.05 (dd, 1H, J = 11.2, 14.6 Hz), 1.59–1.47 (m, 14H), 1.32–1.20 (m, 14H), 1.30 (s, 9H), 0.92 (t, 9H, J = 7.3 Hz); HRMS (FAB), m/z 876.3182 ($M+H$)⁺, ($C_{38}H_{51}N_7O_{15}P$ requires 876.3181).

(α -D-Mannopyranosyl)-tryptophan AMP ester (14b**).** To a test tube containing **14a** (7.8 mg, 6.97 μ mol) was added pre-cooled (0 °C) 4 M HCl in H₂O/dioxane = 1:1 (500 μ L), and the solution was stirred for 13 h at 0 °C. The mixture was diluted with pre-cooled (0 °C) water (5.0 mL), and the solution was subjected to C₁₈ reverse phase column chromatography (Waters Sep-Pak Vac C₁₈ 2 g/12 cc, MeCN/H₂O = 1:1). To avoid saponification, column was performed at 4 °C, fractions containing target molecule were diluted with 10 volumes of MeCN and evaporated at 30 °C. The residues was pumped to give **14b** (4.8 mg, 65%). The ratio of 2'-acyl and 3'-acyl AMP was determined by NMR as 2'-acyl/3'-acyl = 1:5. The peaks corresponding to 3'-acyl AMP were picked up for NMR data: ¹³C NMR (125 MHz, DMSO-*d*₆) δ 168.5, 156.0, 152.7, 149.7, 139.0, 135.7, 135.7, 127.2, 121.3, 118.7, 117.7, 111.6, 106.0, 86.1, 80.8, 79.7, 75.2, 72.7, 71.2, 69.1, 67.9, 65.6, 64.2, 59.7, 57.5, 53.2, 25.9, 23.1, 19.2, 13.5; ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.09 (s, 1H), 8.37 (s, 1H), 8.15 (s, 1H), 7.54 (d, 1H, J = 7.8 Hz),

7.38 (d, 1H, J = 7.8 Hz), 7.30 (br s, 1H), 7.08 (t, 1H, J = 7.1 Hz), 7.04 (t, 1H, J = 7.1 Hz), 5.89 (d, 1H, J = 6.6 Hz), 5.39 (d, 1H, J = 4.2 Hz), 4.94 (d, 1H, J = 8.8 Hz), 4.79 (m, 1H), 4.28 (t, 1H, J = 7.3 Hz), 4.16 (m, 1H), 3.95–3.80 (m, 4H), 3.75–3.68 (m, 2H), 3.65 (dd, 1H, J = 4.2, 11.5 Hz), 3.42 (dd, 1H, J = 6.8, 14.6 Hz), 3.34 (dd, 1H, J = 8.05, 14.6 Hz), 3.15 (m, 8H), 1.55 (m, 8H), 1.30 (m, 8H), 0.92 (t, 12H, J = 7.3 Hz); HRMS (FAB), m/z 696.2024 ($M+H$)⁺, ($C_{27}H_{35}N_7O_{13}P$ requires 696.2030).

Acknowledgements

We thank Dr. H. Koshino and Ms. T. Chijimatsu for NMR measurement, and Dr. T. Chihara and his staff for elemental analysis. We thank Ms. A. Takahashi for technical assistance.

References and Notes

- Varki, A. *Glycobiology* **1993**, *3*, 97.
- (a) For recent review of glycopeptide synthesis Seitz, O. *Chem. Biol. Chem.* **2000**, *1*, 214. Herzner, H.; Reipen, T.; Schluz, M.; Kunz, H. *Chem. Rev.* **2000**, *100*, 4495. (b) Solid-phase chemistry Guo, Z-W.; Nakahara, Y.; Nakahara, Y.; Ogawa, T. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1464. Wang, Z-G.; Zhang, X. F.; Live, D.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2000**, *39*, 3652. (c) Use of intein Tolbert, T. J.; Wong, C.-H. *J. Am. Chem. Soc.* **2000**, *122*, 5421. (d) Native chemical ligation method Shin, Y.; Winans, K. A.; Backes, B. J.; Kent, S. B. H.; Ellman, J. A.; Bertozzi, C. R. *J. Am. Chem. Soc.* **1999**, *121*, 11684. (e) Expressed chemical ligation Macmillan, D.; Bertozzi, C. R. *Tetrahedron* **2000**, *56*, 9515. (f) Use of enzyme Mizuno, M.; Haneda, K.; Iguchi, R.; Muramoto, I.; Kawakami, T.; Aimoto, S.; Yamamoto, K.; Inazu, T. *J. Am. Chem. Soc.* **1999**, *121*, 284.
- Gillmore, M. A.; Steward, L. E.; Chamberlin, A. R. *Topics. Curr. Chem.* **1999**, *202*, 77 Recent review of in vitro peptide synthesis. Dougherty, D. A. *Curr. Opin. Chem. Biol.* **2000**, *4*, 645.
- Mamaev, S. V.; Laikhter, A. L.; Arslan, T.; Hecht, S. M. *J. Am. Chem. Soc.* **1996**, *118*, 7243. Arslan, T.; Mamaev, S. V.; Mamaeva, N. V.; Hecht, S. M. *J. Am. Chem. Soc.* **1997**, *119*, 10877.
- Hohsaka, T.; Ashizuka, Y.; Murakami, H.; Sisido, M. *J. Am. Chem. Soc.* **1996**, *118*, 9778. Hohsaka, T.; Ashizuka, Y.; Sasaki, H.; Murakami, H.; Sisido, M. *J. Am. Chem. Soc.* **1999**, *121*, 12194. Magliery, T. J.; Anderson, J. C.; Schultz, P. G. *J. Mol. Biol.* **2001**, *307*, 755.
- Hohsaka, T.; Ashizuka, Y.; Taira, H.; Murakami, H.; Sisido, M. *Biochemistry* **2001**, *40*, 11060.
- Heckler, T. G.; Chang, L.-H.; Zama, Y.; Naka, T.; Chorghade, M. S.; Hecht, S. M. *Biochemistry* **1984**, *23*, 1468.
- Robertson, S. A.; Noren, C. J.; Anthony-Cahill, S. J.; Griffith, M. C.; Schultz, P. G. *Nucleic Acid Res.* **1989**, *17*, 9649.
- Robertson, S. A.; Ellman, J. A.; Schultz, P. G. *J. Am. Chem. Soc.* **1991**, *113*, 2722.
- Vliegenthart, J. F. G.; Casset, F. *Curr. Opin. Str. Biol.* **1998**, *8*, 565.
- Lodder, M.; Golovine, S.; Laikhter, A. L.; Karginov, V. A.; Hecht, S. M. *J. Org. Chem.* **1998**, *63*, 794 Synthetic effort of glycosylated amino acyl-pdCpA has been made. Fahmi, N. E.; Golovine, S.; Wang, B.; Hecht, S. M. *Carbohydr. Res.* **2001**, *330*, 149.
- Niinomi, T.; Sisido, M. *Chem. Lett.* **1993**, 1305.
- Hart, G. W.; Kreppel, L. K.; Corner, F. I.; Arnold, C. S.;

- Snow, D. M.; Ye, Z.; Chemg, X.; DellaManna, D.; Caine, D. S.; Earles, B. J.; Akimoto, Y.; Cole, R. N.; Hayes, B. K. *Glycobiology* **1996**, *6*, 711. Comer, F. I.; Hart, G. W. *J. Biol. Chem.* **2000**, *275*, 29179.
14. Endo, T. *Biochim. Biophys. Acta* **1999**, *1473*, 237.
15. Manabe, S.; Ito, Y. *J. Am. Chem. Soc.* **1999**, *121*, 9754.
16. Hofsteenge, J.; Müller, D. R.; Beer, T.; Löffler, A.; Richter, W. J.; Vliegenthart, J. F. G. *Biochemistry* **1994**, *33*, 13524. Beer, T.; Vliegenthart, J. F. G.; Löffler, A.; Hofsteenge, J. *Biochemistry* **1995**, *34*, 11785. Löffler, A.; Doucey, M.-A.; Jansson, A. M.; Müller, D. R.; Beer, T.; Hess, D.; Meldal, M.; Richter, W. J.; Vliegenthart, J. F. G.; Hofsteenge, J. *Biochemistry* **1996**, *35*, 12005. Krieg, J.; Gläsner, W.; Vicentini, A.; Doucey, M.-A.; Löffler, A.; Hess, D.; Hofsteenge, J. *J. Biol. Chem.* **1997**, *272*, 26687. Doucey, M.-A.; Hess, D.; Cacan, R.; Hofsteenge, J. *Mol. Biol. Cell.* **1998**, *9*, 291. Krieg, J.; Hartmann, S.; Vicentini, A.; Gläsner, W.; Hess, D.; Hofsteenge, J. *Mol. Biol. Cell.* **1998**, *9*, 301. Doucey, M.-A.; Hess, D.; Blommers, M. J. J.; Hofsteenge, J. *Glycobiology* **1999**, *9*, 435. Hofsteenge, J.; Blommers, M.; Hess, D.; Furmanek, A.; Miroshnichenko, O. *J. Biol. Chem.* **1999**, *274*, 32786. Hartmann, S.; Hofsteenge, J. *J. Biol. Chem.* **2000**, *275*, 28569. Hofsteenge, J.; Huwiler, K. G.; Macek, B.; Hess, D.; Lawler, J.; Mosher, D. F.; Peter-Katalinic, J. *J. Biol. Chem.* **2001**, *276*, 6485.
17. Garcia, A.; Lenis, L. A.; Jiménez, C.; Debitus, C.; Quiñoá, E.; Riguera, R. *Org. Lett.* **2000**, *2*, 2765.
18. Vasella, A.; Witzig, C.; Chiara, J.-L.; Martin-Lomas, M. *Helv. Chim. Acta.* **1991**, *74*, 2073.