

Synthesis and anti-inflammatory evaluation of 4-anilino-furo[2,3-*b*]quinoline and 4-phenoxyfuro[2,3-*b*]quinoline derivatives. Part 3

Yeh-Long Chen,^{a,*} I-Li Chen,^a Chih-Ming Lu,^a Cherng-Chyi Tzeng,^a
Lo-Ti Tsao^b and Jih-Pyang Wang^b

^aFaculty of Medicinal and Applied Chemistry, College of Life Science, Kaohsiung Medical University, Kaohsiung City 807, Taiwan

^bDepartment of Education and Research, Taichung Veterans General Hospital, Taichung 407, Taiwan

Received 8 August 2003; accepted 27 October 2003

Abstract—Mast cells, neutrophils and macrophages are important inflammatory cells that have been implicated in the pathogenesis of acute and chronic inflammatory diseases. To explore a novel anti-inflammatory agent, we have synthesized certain 4-anilino-furo[2,3-*b*]quinoline and 4-phenoxyfuro[2,3-*b*]quinoline derivatives and evaluated their anti-inflammatory activities by reaction of 3,4-dichlorofuro[2,3-*b*]quinoline with appropriate Ar-NH₂ or Ar-OH. Compounds **6a** and **15** were proved to be more potent than the reference inhibitor, mepacrine for the inhibition of rat peritoneal mast cell degranulation with IC₅₀ values of 6.5 and 16.4 μM, respectively. Compounds **2b**, **6a**, **10**, and **15** also showed potent inhibitory activity (IC₅₀ = 7.2–29.4 μM) for the secretion of lysosomal enzyme and β-glucuronidase from neutrophils. These results also indicated that oxime derivatives are more potent than the respective ketone precursors (**6a** ≥ **2a**; **7a** ≥ **3**), and the substituent such as Me at the oxime decreased inhibitory activity (**6a** ≥ **6b**; **7a** ≥ **7b**). Among these derivatives, compound **6a** showed the most potent activity with IC₅₀ values of 6.5–11.6 μM for the inhibition of mast cell degranulation and neutrophil degranulation.

© 2003 Elsevier Ltd. All rights reserved.


1. Introduction

9-Aminoacridine has been used clinically as an anti-septic drug. This tricyclic heterocycle may interact with DNA through intercalation, thus disrupting DNA replication.^{1,2} A large number of its derivatives have been prepared and evaluated for biological activities.^{3–5} Two notable examples are mepacrine (quinacrine), the acridine derivative to be clinically used as an anti-malarial drug which also acts as a calmodulin inhibitor to suppress the histamine secretion process in mast cell^{6–9} and amsacrine (m-AMSA), an antileukemic agent.^{10–12} Certain 9-thioacridines have also been synthesized as inhibitors of trypanothione reductase from *Trypanosoma cruzi*, the causative agent of Chagas' disease.¹³ Due to the biological versatility of acridine derivatives, we have synthesized certain 9-anilinoacridine, 9-phenoxyacridine, and 4-phenoxyfuro[2,3-*b*]quinoline deri-


vatives and evaluated their anti-inflammatory activities.^{14,15} Recently, we have also synthesized certain 4-anilino-furo[2,3-*b*]quinoline derivatives from a natural alkaloid, dictamnine, and evaluated their cytotoxicity.^{16,17} We were especially interested in the study of furo[2,3-*b*]quinoline because it constitutes an important group of bioactive natural products such as dictamnine, acrophylline, confusameline, skimmianine, kokusaginine, and haplopine.^{18–23} These alkaloids were found to possess wide-ranging biological properties including anti-allergic,¹⁸ anti-inflammatory,¹⁹ cytotoxic,²⁰ anti-platelet aggregation,^{21,22} and the voltage-gated potassium channel blocking activities.²³ Although the 4-anilino-furo[2,3-*b*]quinoline ring belongs to the isosteric isomer of 9-anilinoacridine, comparison on the biological activities between these two rings have not been explored. In continuation of our studies on 9-anilinoacridine derivatives, we report herein the preparation and anti-inflammatory activities of certain 4-anilino-furo[2,3-*b*]quinoline derivatives. We have also synthesized and evaluate a number of 4-phenoxyfuro[2,3-*b*]quinoline derivatives for the establishment of anti-inflammatory structure–activity relationships.

Keywords: Anti-inflammatory; 4-Anilino-furo[2,3-*b*]quinoline; 4-Phenoxyfuro[2,3-*b*]quinoline; Furo[2,3-*b*]quinoline.


* Corresponding author. Tel.: +886-7-312-1101x2249; fax: +886-7-312-5339; e-mail: eloch@cc.kmu.edu.tw



Scheme 1. (i) Substituted-anilines, EtOH–H₂O (2:1 v/v), HCl, reflux; (ii) R'ONH₂, EtOH, reflux; (iii) *N*-[4-(3-oxobut-1-enyl)phenyl]acetamide, EtOH–H₂O (2:1 v/v), HCl, reflux.



Scheme 2. (i) 3-Hydroxyacetophenone, EtOH–H₂O (2:1 v/v), HCl, reflux; (ii) R'ONH₂, EtOH, reflux; (iii) H₂, Lindlar catalyst, EtOH–CH₂Cl₂ (1:1 v/v).



Scheme 3. Reagents (i) 4-Hydroxybenzaldehyde, K₂CO₃, THF in a sealed bomb; (ii) 4-Hydroxybenzylideneacetone, K₂CO₃, acetone in a sealed bomb; (iii) Lindlar catalyst, H₂, MeOH–CH₂Cl₂ (1:1 v/v).

2. Chemistry

4-Anilino-furo[2,3-*b*]quinolines 2–7 were prepared as described in Scheme 1. Reaction of the known 3,4-dichloro-furo[2,3-*b*]quinoline (**1**)²⁴ with 4-aminobenzoic

acid in EtOH/H₂O (2:1) afforded 4-(3-chlorofuro[2,3-*b*]quinolin-4-ylaminobenzoic acid (**2b**) in 57% yield. Accordingly, 4-[4-(3-chlorofuro[2,3-*b*]quinolin-4-ylamino)phenyl]-but-3-en-2-one (**5**) was prepared from *N*-[4-(3-oxobut-1-enyl)phenyl]acetamide²⁵ under the same reaction condition. Preparation of **2a**, **3**, and **4** had been described in our previously paper.¹⁶ Compounds **2a** and **3** were reacted with hydroxylamine or methoxylamine, respectively, in EtOH to give the corresponding oximes **6a**, **7a** and methyloximes **6b**, **7b** in 62–97% yield.

Preparation of 4-phenoxyfuro[2,3-*b*]quinolines **8–10** is described in Scheme 2. Compound **1** was treated with 3-hydroxyacetophenone to afford 1-[3-(3-chlorofuro[2,3-*b*]quinolin-4-yloxy)phenyl]ethanone (**8**) which was then reduced with H₂ in the presence of Lindlar catalyst to give 1-[3-(furo[2,3-*b*]quinolin-4-yloxy)phenyl]ethanone (**10**) in 69% yield. Reaction of **8** with hydroxylamine or methoxylamine, respectively, in EtOH gave the corresponding oximes **9a** and methyloximes **9b** in 87–90% yield.

4-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)benzaldehyde (**11**) was synthesized from **1** and 4-hydroxybenzaldehyde, which was then reduced with Lindlar catalyst and H₂ to give 4-(furo[2,3-*b*]quinolin-4-yloxy)-benzaldehyde (**13**) and [4-(furo[2,3-*b*]quinolin-4-yloxy)-phenyl]methanol (**14**) (Scheme 3). Accordingly, reaction of **1** and 4-hydroxybenzylideneacetone afforded **12**, which was reduced to give 4-[4-(furo[2,3-*b*]quinolin-4-yloxy)phenyl]butan-2-one (**15**) and 4-[4-(furo[2,3-*b*]quinolin-4-yloxy)phenyl]butan-2-ol (**16**).

3. Biological results and discussion

3.1. Mast cell degranulation

In the present study, assessment of inhibitory efficacy with respect to mast cell degranulation was performed by measuring the content of β-glucuronidase in supernatant. As shown in Table 1, (*E*)-1-[3-(3-chlorofuro[2,3-*b*]quinolin-4-ylamino)phenyl]ethanone oxime (**7a**) demonstrated only a weak inhibitory activity (IC₅₀ value of 35.8 μM) while (*E*)-1-[4-(3-chlorofuro[2,3-*b*]quinolin-4-ylamino)phenyl]ethanone oxime (**6a**) (6.5 μM) and 4-[4-(furo[2,3-*b*]quinolin-4-yloxy)phenyl]butan-2-one (**15**) (16.4 μM) exhibited more potent activity than the reference inhibitor, mepacrine (20.6 μM). These results also indicated that oxime derivatives are more potent than the respective ketone precursors (**6a** ≥ **2a**; **7a** ≥ **3**) and the substituent such as Me at the oxime decreased inhibitory activity (**6a** ≥ **6b**; **7a** ≥ **7b**). The result that compound **15** was more potent than **13** suggested that the distance between furoquinoline and the carbonyl group also play an important role.

3.1.1. Neutrophil degranulation. Activation of neutrophils with 1 μM formyl-methionyl-leucyl-phenylalanine (fMLP) in the presence of cytochalasin B (5 μg/mL) evoked the release of 21.2 and 19.8% of lysozyme and β-glucuronidase, respectively, of the initial cellular content. Compound **6a**, with an IC₅₀ value of 11.6 and

Table 1. IC₅₀ (μM)^a values of furo[2,3-*b*]quinoline derivatives against mast cell and neutrophil degranulation

Compd	Mast cell degranulation β-glucuronidase ^b	Neutrophil degranulation	
		Lysozyme ^c	β-glucuronidase ^c
2a	> 30.0	> 30.0	> 30.0
2b	> 30.0	23.3 ± 6.9	18.9 ± 5.7
3	> 30.0	> 30.0	> 30.0
4	> 30.0	> 30.0	> 30.0
5	> 30.0	> 30.0	> 30.0
6a	6.5 ± 1.1	11.6 ± 1.7	7.2 ± 0.9
6b	> 30.0	> 30.0	> 30.0
7a	35.8 ± 1.1	> 30.0	> 30.0
7b	> 30.0	> 30.0	> 30.0
8	> 30.0	> 30.0	> 30.0
9a	> 30.0	> 30.0	> 30.0
9b	> 30.0	> 30.0	> 30.0
10	> 30.0	29.4 ± 2.0	26.1 ± 5.9
11	> 30.0	> 30.0	> 30.0
12	> 30.0	> 30.0	32.0 ± 2.4
13	> 30.0	> 30.0	13.0 ± 0.5
14	> 30.0	> 30.0	> 30.0
15	16.4 ± 1.8	14.0 ± 1.2	9.3 ± 1.2
Mepacrine	20.6 ± 1.2		
Trifluoperazine		11.9 ± 0.6	10.6 ± 0.9

^a Values are means ± SE of at least three separate experiments.^b Induced by compound 48/80 (10 μg/mL).^c Induced by fMLP (1 μM)/cytochalasin B (5 μg/mL).

7.2 μM against lysozyme and β-glucuronidase release, respectively, was more potent than its positional isomer **7a** and was comparable to the calmodulin inhibitor, trifluoperazine (11.9 and 10.6 μM, respectively) which inhibits the degradation and superoxide anion generation in neutrophils.^{26,27} 4-(3-Chlorofuro[2,3-*b*]quinolin-4-ylaminobenzoic acid (**2b**) with an IC₅₀ value of 23.3 and 18.9 μM against lysozyme and β-glucuronidase release, respectively, was more potent than its acetyl analogue **2a**. In analogy to the inhibition of mast cell degranulation, oxime derivatives are more potent than the respective ketone precursors (**6a** ≥ **2a**), and the substituent such as Me at the oxime decreased inhibitory activity (**6a** ≥ **6b**) and a longer distance between fur-quinoline and the carbonyl group was preferred (**12** ≥ **11**; **15** ≥ **13**). In addition, the inhibitory activity was decreased by the reduction of carbonyl group to an alcohol (**13** ≥ **14**), but was increased by the reduction of Cl at the 3-position (**10** ≥ **8**). Compound **6a** showed the most potent activity with IC₅₀ values of 7–12 μM for the inhibition of neutrophil degranulation.

3.2. TNF-α release

TNF-α, an early cytokine produced by activated macrophages, plays an essential role in pathological inflammatory reactions. None of compounds **2–15** had similar IC₅₀ value as dexamethasone in the inhibition of TNF-α formation in macrophage-like cell line RAW 264.7 and microglial cell line N9 cells (the brain resident macrophages) (Table 2). However, (*E*)-1-[3-(3-chlorofuro[2,3-*b*]quinolin-4-yloxy)phenyl]ethanone oxime (**9a**) showed weak potency in N9 cells (IC₅₀ values of 1.7 μM). The results also confirm the conclusion above that the substituent such as Me at the oxime decreased inhibitory activity (**9a** ≥ **9b**) (Table 2).

Table 2. IC₅₀ values of furo[2,3-*b*]quinoline derivatives on TNF-α formation

Compd	IC ₅₀ (μM) ^{a,b}	
	RAW 264.7	N9
2a	> 10.0	> 0.3
2b	> 30.0	> 30.0
3	> 30.0	> 30.0
4	> 30.0	> 30.0
5	> 10.0	> 3.0
6a	> 10.0	> 1.0
6b	> 30.0	> 3.0
7a	> 30.0	> 3.0
7b	> 30.0	> 3.0
8	> 30.0	> 30.0
9a	> 30.0	1.7 ± 0.1
9b	> 30.0	> 10.0
10	> 30.0	> 10.0
11	> 30.0	> 3.0
12	> 10.0	> 10.0
13	> 30.0	> 30.0
14	> 30.0	> 30.0
15	> 30.0	> 10.0
Dexamethasone	0.42 ± 0.12	0.074 ± 0.01

^a Values are means ± SE of at least three separate experiments.^b When compounds exhibited cytotoxicity at 30 μM, a lower concentration was applied.

3.3. Preliminary cytotoxic evaluation

Nine furo[2,3-*b*]quinoline derivatives were selected for evaluation in vitro against a three-cell line panel consisting of MCF7 (Breast), NCI-H460 (Lung), and SF-268 (CNS). In this protocol, each cell line is inoculated and preincubated on a microtiter plate. Test agents are then added at a single concentration (100 μM) and the culture incubated for 48 h. End-point determinations are made with alamar blue.²⁸ Results for each test agent are reported as the percent of growth of the treated cells when compared to the untreated control cells. Compounds which reduced the growth of any one of the cell lines to 32% or less are considered to be cytotoxic. Results indicated all of them, with the exception of compound **5** which exhibited a weak inhibitory activity on MCF7 cancer cell, are non-cytotoxic.

4. Conclusion

These results indicated that the anti-inflammatory effects of furo[2,3-*b*]quinoline derivatives were mediated, at least in part, through the suppression of chemical mediators released from mast cells, neutrophils and macrophages, and the potential of these compounds to be novel anti-inflammatory agents with no significant cytotoxicity.

5. Experimental

5.1. General

TLC: precoated (0.2 mm) silica gel 60 F₂₅₄ plates from EM Laboratories, Inc.; detection by UV light (254 nm). Mp: Electrothermal IA9100 digital melting-point appa-

ratus; uncorrected. ^1H and ^{13}C NMR spectra: Varian-Unity-400 spectrometer at 400 and 100 MHz or Varian-Gemini-200 spectrometer at 200 and 50 MHz, chemical shifts δ in ppm with SiMe_4 as an internal standard ($=0$ ppm), coupling constants J in Hz. Elemental analyses were carried out on a Heraeus CHN-O-Rapid elemental analyzer, and results were within $\pm 0.4\%$ of calculated values.

5.1.1. 4-(3-Chlorofuro[2,3-*b*]quinolin-4-ylaminobenzoic acid (2b). To a solution of 3,4-dichlorofuro[2,3-*b*]quinoline (**1**, 1.19 g, 5.00 mmol) and 4-aminobenzoic acid (1.03 g, 7.50 mmol) in EtOH/H₂O 2:1 (50 mL) was added concentrated HCl until pH 6 resulted. The mixture was refluxed for 18 h (TLC monitoring) and then the solvent evaporated in vacuo to give a residual solid, which was suspended in ice-water (250 mL) and neutralized with 1 N NaOH solution. The resulting precipitate was collected by filtration, washed with H₂O, and then purified by flash column chromatography (FC, silica gel CH₂Cl₂/MeOH 20:1) and recrystallized from EtOH to give **2b** (0.96 g, 57%). Mp 262–263°C. ^1H NMR (200 MHz, DMSO-*d*₆): 6.86 (m, 2H-C(2', 6')), 7.59 (m, H-C(6)), 7.75 (m, 2H-C(3', 5')), 7.82 (m, H-C(7)), 8.04 (dd, $J=8.4$, 0.8, H-C(8)), 8.22 (dd, $J=8.6$, 0.8, H-C(5)), 8.40 (s, H-C(2)), 9.45 (br s, NH), 12.65 (br s, COOH). ^{13}C NMR (50 MHz, DMSO-*d*₆): 110.11, 111.05, 114.84, 121.70, 122.61, 123.73, 125.18, 128.65, 130.54, 131.31, 139.98, 143.37, 146.04, 150.73, 160.88, 167.43. HRMS (EI): calcd for C₁₈H₁₁ClN₂O₃: 338.0458. Found: 338.0462.

5.1.2. 4-[4-(3-Chlorofuro[2,3-*b*]quinolin-4-ylamino)-phenyl]but-3-en-2-one (5). This compound was obtained from **1** and *N*-[4-(3-oxobut-1-enyl)phenyl]acetamide as described for **2b**, which was purified by FC (CH₂Cl₂/AcOEt 20:1) in 48% yield. Mp 207–208°C. ^1H NMR (400 MHz, CDCl₃): 2.36 (s, COMe), 6.62 (d, $J=16.4$, CH=CHCOMe), 6.93 (m, 2H-C(2', 6')), 7.31 (m, H-C(6)), 7.46 (m, 2H-C(3', 5')), CH=CHCOMe and NH), 7.71 (m, 2H-C(2, 7)), 7.89 (d, $J=8.0$, H-C(8)), 8.13 (d, $J=8.4$, H-C(5)). ^{13}C NMR (100 MHz, CDCl₃): 27.48, 108.36, 110.42, 118.41 (2C), 120.06, 124.25, 124.52, 125.46, 128.25, 128.74, 129.74 (2C), 130.50, 141.00, 141.64, 142.76, 145.55, 145.73, 160.02, 198.27. Anal. calcd for C₂₁H₁₅ClN₂O₂: C 69.52, H 4.17, N 7.72; Found: C 69.27, H 4.27, N 7.66.

5.1.3. (E)-1-[4-(3-Chlorofuro[2,3-*b*]quinolin-4-ylamino)-phenyl]ethanone oxime (6a). To a suspension of **2a** (216 mg, 0.64 mmole) in ethanol (20 mL) was added NH₂OH·HCl (89 mg, 1.28 mmol). The reaction mixture was stirred at room temperature for 1 h (TLC monitoring), then concentrated in vacuo to give a solid which was washed by H₂O (20 mL), purified by FC (CH₂Cl₂-AcOEt 20:1) and recrystallized from EtOH to give a yellow solid (140 mg, 62%). Mp 219–220°C. ^1H NMR (200 MHz, DMSO-*d*₆): 2.10 (s, Me), 6.90 (m, 2H-C(2', 6')), 7.54 (m, 3H-C(6, 3', and 5')), 7.80 (m, H-C(7)), 8.01 (d, $J=8.6$, H-C(8)), 8.26 (d, $J=8.4$, H-C(5)), 8.34 (s, H-C(2)), 9.16 (br s, NH), 10.93 (br s, NOH). ^{13}C NMR (50 MHz, DMSO-*d*₆): 11.21, 109.08, 109.90, 115.86 (2C), 121.61, 123.47, 124.36, 126.38 (2C), 128.26,

128.91, 130.01, 140.78, 142.24, 145.74, 146.44, 152.37, 160.72. Anal. calcd for C₁₉H₁₄ClN₃O₂·0.2HCl: C 63.55, H 3.99, N 11.70; Found: C 63.94, H 4.28, N 11.33.

5.1.4. (E)-1-[4-(3-Chlorofuro[2,3-*b*]quinolin-4-ylamino)-phenyl]ethanone *O*-methyloxime (6b). To a suspension of **2a** (103 mg, 0.30 mmol) in ethanol (16 mL) was added 40% NH₂Ome·HCl aqueous (126 mg, 0.6 mmol). The reaction mixture was stirred at room temperature for 1 h (TLC monitoring), then concentrated in vacuo to give a solid which was washed by H₂O (20 mL) and purified by FC (CH₂Cl₂) to give a white solid (102 mg, 93%). Mp 213–215°C. ^1H NMR (200 MHz, CDCl₃): 2.20 (s, Me), 3.98 (s, NOME), 6.89 (m, 2H-C(2', 6')), 7.20 (br s, NH), 7.31 (m, H-C(6)), 7.55 (m, 2H-C(3', 5')), 7.68 (m, 2H-C(2, 7)), 7.82 (d, $J=8.4$, H-(8)), 8.08 (d, $J=8.4$, H-C(5)). ^{13}C NMR (50 MHz, CDCl₃): 12.46, 61.86, 107.78, 110.13, 118.28 (2C), 119.88, 123.96, 124.45, 127.10 (2C), 128.92, 129.93, 130.95, 105.55, 141.75, 144.39, 146.58, 153.96, 160.57. Anal. calcd for C₂₀H₁₆ClN₃O₂·0.3HCl: C 63.76, H 4.36, N 11.15; Found: C 63.71, H 4.12, N 11.29.

5.1.5. (E)-1-[3-(3-Chlorofuro[2,3-*b*]quinolin-4-ylamino)-phenyl]ethanone oxime (7a). This compound was obtained from **3** and NH₂OH·HCl as described for **6a**, which was purified by FC (CH₂Cl₂/MeOH 100:1) in 93% yield. Mp. 250–252°C. ^1H NMR (200 MHz, DMSO-*d*₆): 2.08 (s, Me), 6.90 (m, H-C(6')), 7.13–7.26 (m, 3H-C(2', 4', 5')), 7.55 (m, H-C(6)), 7.80 (m, H-C(7)), 8.01 (d, $J=8.4$, H-C(8)), 8.30 (m, 2H-C(2, 5)), 9.10 (br s, NH), 11.11 (br s, NOH). ^{13}C NMR (50 MHz, DMSO-*d*₆): 11.44, 108.51, 109.99, 113.33, 117.02, 118.24, 121.44, 123.53, 124.35, 128.29, 129.10, 130.08, 137.73, 141.27, 142.09, 145.79, 145.88, 152.66, 160.81. Anal. calcd for C₁₉H₁₄ClN₃O₂: C 64.87, H 4.01, N 11.94; Found: C 64.83, H 4.09, N 11.73.

5.1.6. (E)-1-[3-(3-Chlorofuro[2,3-*b*]quinolin-4-ylamino)-phenyl]ethanone *O*-methyloxime (7b). This compound was obtained from **3** and 40%·NH₂Ome·HCl as described for **6b**, which was purified by FC (CH₂Cl₂) in 97% yield. Mp 151–152°C. ^1H NMR (200 MHz, DMSO-*d*₆): 2.11 (s, Me), 3.87 (s, NOME), 6.86 (m, H-C(6')), 7.17–7.34 (m, 3H-C(2', 4', 5')), 7.54 (m, H-C(6)), 7.78 (m, H-C(7)), 7.98 (d, $J=8.4$, H-C(8)), 8.30 (m, 2H-C(2, 5)), 9.13 (br s, NH). ^{13}C NMR (50 MHz, DMSO-*d*₆): 12.32, 61.46, 108.36, 109.92, 114.05, 117.06, 118.40, 121.30, 123.44, 124.26, 128.24, 129.08, 130.00, 136.68, 141.05, 142.02, 145.73, 145.84, 153.88, 160.77. Anal. calcd for C₂₀H₁₆ClN₃O₂·0.1H₂O: C 65.35, H 4.44, N 11.43; Found: C 65.28, H 4.50, N 11.08.

5.1.7. 1-[3-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)phenyl]ethanone (8). A mixture of **1** (0.72 g, 3.0 mmol), K₂CO₃ (0.62 g, 4.5 mmol), and 3-hydroxyacetophenone (0.49 g, 3.6 mmol) in acetone (50 mL) was heated at 150°C for 20 h in a steel bomb. It was cooled, the solvent evaporated in vacuo to give a residual solid, which was suspended in ice-water (100 mL). The resulting precipitate was collected by filtration, washed with H₂O, and then purified by FC (*n*-hexane/EtOAc 4:1) and recrystallized from EtOH to give a white solid **8** (0.80 g, 79%). Mp

168–169°C. ¹H NMR (200 MHz, CDCl₃): 2.56 (s, Me), 7.08 (m, H-C(6')), 7.41 (dd, *J*=8.2, 7.8, H-C(5')), 7.51 (m, 2H-C(2', 6)), 7.67 (m, H-C(4')), 7.73 (s, H-C(2)), 7.79 (m, H-C(7)), 8.08 (dd, *J*=8.4, 1.2, H-C(8)), 8.17 (d, *J*=8.4, H-C(5)). ¹³C NMR (50 MHz, CDCl₃): 26.72, 109.90, 114.82, 120.15, 121.20, 122.04, 123.15, 123.46, 125.70, 128.69, 130.13, 130.62, 138.96, 142.26, 146.75, 151.53, 159.50, 161.22, 197.18. Anal. calcd for C₁₉H₁₂ClNO₃: C 67.56, H 3.58, N 4.15; Found: C 67.52, H 3.63, N 4.13.

5.1.8. (E)-1-[3-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)-phenyl]ethanone oxime (9a). This compound was obtained from **8** and NH₂OH·HCl as described for **6a**, which was purified by FC (CH₂Cl₂/EtOAc 20:1) in 90% yield. Mp 181–183°C. ¹H NMR (400 MHz, DMSO-*d*₆): 2.11 (s, Me), 7.00 (m, H-C(6')), 7.23 (m, H-C(2')), 7.37 (m, 2H-C(4', 5')), 7.62 (m, H-C(6)), 7.88 (m, H-C(7)), 8.04 (dd, *J*=8.4, 0.8, H-C(8)), 8.13 (d, *J*=8.4, H-C(5)), 8.55 (s, H-C(2)), 11.25 (s, NOH). ¹³C NMR (100 MHz, DMSO-*d*₆): 11.46, 108.56, 110.16, 111.63, 115.83, 120.57, 120.72, 121.87, 126.01, 128.35, 130.22, 130.81, 138.89, 144.19, 146.07, 150.88, 152.22, 158.89, 160.83. Anal. calcd for C₁₉H₁₃ClN₂O₃: C 64.69, H 3.71, N 7.94; Found: C 64.33, H 3.86, N 7.60.

5.1.9. (E)-1-[3-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)-phenyl]ethanone *O*-methyloxime (9b). This compound was obtained from **8** and 40% NH₂OMe·HCl as described for **6b**, which was purified by FC (CH₂Cl₂) in 87% yield. Mp 137–139°C. ¹H NMR (200 MHz, CDCl₃): 2.18 (s, Me), 3.97 (s, NOME), 6.73 (m, H-C(6')), 7.25 (dd, *J*=8.4, 8.0, H-C(5')), 7.38 (m, 2H-C(2', 4')), 7.51 (m, H-C(6)), 7.72 (s, H-C(2)), 7.79 (m, H-C(7)), 8.14 (m, H-C(5, 8)). ¹³C NMR (50 MHz, CDCl₃): 12.67, 62.04, 110.19, 113.63, 115.70, 120.78, 121.42, 122.43, 125.58, 128.62, 129.78, 130.58, 138.78, 142.11, 142.72, 146.79, 152.08, 153.81, 159.38, 161.35. Anal. calcd for C₂₀H₁₅ClN₂O₃: C 65.49, H 4.12, N 7.64; Found: C 65.28, H 4.17, N 7.47.

5.1.10. 1-[3-(Furo[2,3-*b*]quinolin-4-yloxy)phenyl]ethanone (10). A solution of **8** (0.23 g, 0.68 mmol) in MeOH–CH₂Cl₂ (1/1, 60 mL) was hydrogenated for 1 h (TLC monitoring) under H₂ with Lindlar catalyst (0.23 g). The reaction mixture was filtered and the filtrate concentrated in vacuo to give a residual solid, which was purified by FC (*n*-hexane–EtOAc 2:1) to give **10** (0.14 g, 69%). Mp 132–133°C. ¹H NMR (200 MHz, CDCl₃): 2.60 (s, Me), 5.81 (d, *J*=2.6, H-C(3)), 7.36 (m, H-C(6')), 7.51 (d, *J*=2.6, H-C(2)), 7.57 (m, 2H-C(5', 6)), 7.76–7.89 (m, 3H-C(2', 4', 7)), 8.18 (d, *J*=8.2, H-C(8)), 8.37 (dd, *J*=8.4, 1.2, H-C(5)). ¹³C NMR (50 MHz, CDCl₃): 26.74, 103.53, 107.32, 118.84, 119.31, 122.04, 123.86, 124.84, 127.86, 130.21, 130.40, 139.13, 144.82, 145.57, 152.16, 153.56, 156.56, 163.05, 196.82. Anal. calcd for C₁₉H₁₃NO₃·0.1H₂O: C 74.80, H 4.36, N 4.59; Found: C 74.80, H 4.43, N 4.55.

5.1.11. 4-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)benzaldehyde (11). This compound was obtained from **1** and 4-hydroxybenzaldehyde in THF as described for **8**, which was purified by FC (CH₂Cl₂) in 72% yield. Mp 194–

195°C. ¹H NMR (200 MHz, CDCl₃): 7.02 (m, 2H-C(2', 6')), 7.54 (m, H-C(6)), 7.76 (s, H-C(2)), 7.81 (m, H-C(7)), 7.86 (m, 2H-C(3', 5')), 8.05 (dd, *J*=8.6, 1.6, H-C(8)), 8.19 (d, *J*=8.6, H-C(5)), 9.93 (s, CHO). ¹³C NMR (50 MHz, CDCl₃): 109.85, 110.15, 116.04 (2C), 120.96, 121.81, 125.91, 128.80, 130.79, 131.78, 132.19 (2C), 142.53, 146.77, 150.58, 161.15, 163.58, 190.51. Anal. calcd for C₁₈H₁₀ClNO₃: C 66.78, H 3.11, N 4.33; Found: C 66.63, H 3.14, N 4.34.

5.1.12. 4-[4-(3-Chlorofuro[2,3-*b*]quinolin-4-yloxy)phenyl]-but-3-en-2-one (12). This compound was obtained from **1** and 4-hydroxybenzylideneacetone in acetone as described for **8**, which was purified by FC (CH₂Cl₂) in 87% yield. Mp 185–186°C. ¹H NMR (200 MHz, CDCl₃): 2.36 (s, Me), 6.61 (d, *J*=16.0, CH=CH(C=O)Me), 6.91 (m, 2H-C(2', 6')), 7.42–7.56 (m, 3H-C(3', 5', 6) and CH=CH(C=O)Me), 7.73 (s, H-C(2)), 7.79 (m, H-C(7)), 8.07 (dd, *J*=8.0, 1.0, H-C(8)), 8.17 (d, *J*=8.6, H-C(5)). ¹³C NMR (50 MHz, CDCl₃): 27.50, 109.97, 110.17, 116.12 (2C), 121.18, 122.05, 125.75, 126.27, 128.73, 129.39, 130.15 (2C), 130.65, 142.42, 146.78, 151.45, 160.88, 161.24, 198.16. Anal. calcd for C₂₁H₁₄ClNO₃·0.1H₂O: C 68.99, H 3.91, N 3.83; Found: C 68.92, H 3.93, N 3.75.

5.1.13. 4-(Furo[2,3-*b*]quinolin-4-yloxy)benzaldehyde (13) and 4-(furo[2,3-*b*]quinolin-4-yloxy)phenylmethanol (14). Hydrogenation of **11** as described for **10**, which was purified by FC (CH₂Cl₂) to give **13** (23% yield) and **14** (34% yield).

Compound 13. Mp 129–131°C. ¹H NMR (200 MHz, CDCl₃): 6.16 (d, *J*=2.6, H-C(3)), 7.22 (m, 2H-C(2', 6')), 7.55 (m, H-C(6)), 7.61 (d, *J*=2.6, H-C(2)), 7.79 (m, H-C(7)), 7.94 (m, 2H-C(3', 5')), 8.18 (d, *J*=8.4, H-C(8)), 8.25 (dd, *J*=8.6, 1.0, H-C(5)), 10.00 (s, CHO). ¹³C NMR (50 MHz, CDCl₃): 103.43, 108.97, 118.67 (2C), 119.70, 121.87, 125.19, 128.21, 130.26, 132.08 (2C), 132.82, 145.65, 145.90, 152.11, 161.38, 163.02, 190.45. Anal. calcd for C₁₈H₁₁NO₃·0.1H₂O: C 74.28, H 3.88, N 4.81; Found: C 74.32, H 4.23, N 4.76.

Compound 14. Mp 163–164°C. ¹H NMR (400 MHz, DMSO-*d*₆): 4.57 (d, *J*=5.6, CH₂OH), 5.32 (t, *J*=5.6, OH), 5.66 (d, *J*=2.8, H-C(3)), 7.27 (m, 2H-C(2', 6')), 7.45 (m, 2H-C(3', 5')), 7.62 (m, H-C(6)), 7.83 (m, H-C(7)), 7.93 (d, *J*=2.8, H-C(2)), 8.04 (d, *J*=8.0, H-C(8)), 8.36 (dd, *J*=8.4, 0.8, H-C(5)). ¹³C NMR (100 MHz, DMSO-*d*₆): 62.27, 103.44, 106.40, 118.61, 119.52 (2C), 121.95, 124.66, 127.77, 128.27 (2C), 129.96, 139.91, 145.34, 145.66, 153.63, 154.46, 163.06. Anal. calcd for C₁₈H₁₃NO₃·0.1H₂O: C 73.76, H 4.54, N 4.78; Found: C 73.63, H 4.61, N 4.74.

5.1.14. 4-[4-(Furo[2,3-*b*]quinolin-4-yloxy)phenyl]butan-2-one (15) and 4-[4-(furo[2,3-*b*]quinolin-4-yloxy)phenyl]butan-2-ol (16). Hydrogenation of **12** as described for **10**, which was purified by FC to give **15** (eluting with CH₂Cl₂/EtOAc 20:1 and 21% yield) and **16** (eluting with CH₂Cl₂/EtOAc 4:1 and 28% yield).

Compound 15. Mp 113–115°C. ^1H NMR (200 MHz, DMSO-d_6): 2.12 (s, Me), 2.84 (br s, $\text{CH}_2\text{CH}_2(\text{C}=\text{O})\text{Me}$), 5.62 9d, $J=2.6$, H-C(3)), 7.21 (m, $2\text{H-C}(2', 6')$), 7.35 (m, $2\text{H-C}(3', 5')$), 7.61 (m, H-C(6)), 7.82 (m, H-C(7)), 7.92 (d, $J=2.6$, H-C(2)), 8.03 (d, $J=8.6$, H-C(8)), 8.34 (dd, $J=8.4, 0.8$, H-C(5)). ^{13}C NMR (50 MHz, DMSO-d_6): 28.38, 29.70, 44.09, 103.32, 106.26, 118.51, 119.66 (2C), 121.87, 124.55, 127.68, 129.87, 129.93 (2C), 138.54, 145.26, 145.55, 153.61, 153.82, 162.99, 207.48. Anal. calcd for $\text{C}_{21}\text{H}_{17}\text{NO}_3$: C 76.12, H 5.17, N 4.23; Found: C 75.92, H 5.30, N 4.20.

Compound 16. Pale-yellow oil. ^1H NMR (200 MHz, DMSO-d_6): 1.11 (d, $J=6.0$, Me), 1.66 (m, $\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{Me}$), 2.70 (m, $\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{Me}$), 3.63 (m, $-\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{Me}$), 4.51 (d, $J=5.0$, OH), 5.61 (d, $J=2.8$, H-C(3)), 7.22 (m, $2\text{H-C}(2', 6')$), 7.33 (m, $2\text{H-C}(3', 5')$), 7.61 (m, H-C(6)), 7.82 (m, H-C(7)), 7.92 (d, $J=2.8$, H-C(2)), 8.03 (dd, $J=8.4, 1.2$, H-C(8)), 8.35 (dd, $J=8.4, 1.4$, H-C(5)). ^{13}C NMR (50 MHz, DMSO-d_6): 23.63, 30.88, 40.90, 65.16, 103.36, 106.21, 118.55, 119.77 (2C), 121.95, 124.61, 127.73, 129.63, 129.99 (2C), 139.93, 145.31, 145.56, 153.64, 153.78, 163.06. Anal. calcd for $\text{C}_{21}\text{H}_{19}\text{NO}_3 \cdot 0.2\text{H}_2\text{O}$: C 74.85, H 5.80, N 4.16; Found: C 75.15, H 6.08, N 3.97.

Acknowledgements

Financial support of this work by the National Science Council of the Republic of China (NSC 91-2113-M-037-013) is gratefully acknowledged. We also thank National Cancer Institute (NCI) of the United States for the anticancer screenings and the National Center for High-Performance Computing for providing computer resources and chemical database services.

References and notes

- Wakelin, L. P. G.; Adams, A.; Denny, W. A. *J. Med. Chem.* **2002**, *45*, 894.
- Bailly, C.; Denny, W. A.; Mellor, L. E.; Wakelin, L. P. G.; Waring, M. J. *Biochemistry* **1992**, *31*, 3514.
- Gamage, S. A.; Tepsiri, N.; Wilairat, P.; Wojcik, S. J.; Figgitt, D. P.; Ralph, R. K.; Denny, W. A. *J. Med. Chem.* **1994**, *37*, 1486.
- McConnaughie, A. W.; Jenkins, T. C. *J. Med. Chem.* **1995**, *38*, 3488.
- Gamage, S. A.; Figgitt, D. F.; Wojcik, S. J.; Ralph, R. K.; Ransijn, A.; Mauel, J.; Yardley, V.; Snowdon, D.; Croft, S. L.; Denny, W. A. *J. Med. Chem.* **1997**, *40*, 2634.
- Amellal, M.; Landry, Y. *Br. J. Pharmacol.* **1983**, *80*, 365.
- Chakravarty, N.; Nielsen, E. H. *Agents Actions* **1986**, *18*, 65.
- Yu, N.; Maciejewski-Lenoir, D.; Bloom, F. E.; Magistretti, P. J. *Mol. Pharmacol.* **1995**, *48*, 550.
- Romanelli, F.; Fillo, S.; Isidori, A.; Conte, D. *Life Sci.* **1997**, *61*, 557.
- Denny, W. A.; Cain, B. F.; Hansch, G. J.; Panthanickal, A.; Leo, A. *J. Med. Chem.* **1982**, *25*, 276.
- Atwell, G. J.; Cain, B. F.; Seelye, R. N. *J. Med. Chem.* **1972**, *15*, 611.
- Zwelling, L. A.; Michaels, S.; Erickson, L. C.; Ungerleider, R. S.; Nichols, M.; Kohn, K. W. *Biochemistry* **1981**, *20*, 6553.
- Bonse, S.; Santelli-Rouvier, C.; Barbe, J.; Krauth-Siegel, R. L. *J. Med. Chem.* **1999**, *42*, 5448.
- Chen, Y.-L.; Lu, C.-M.; Chen, I.-L.; Tsao, L.-T.; Wang, J.-P. *J. Med. Chem.* **2002**, *45*, 4689.
- Chen, Y.-L.; Chen, I.-L.; Lu, C.-M.; Tzeng, C.-C.; Tsao, L.-T.; Wang, J.-P. *Bioorg. Med. Chem.* **2003**, *11*, 3921.
- Chen, I.-L.; Chen, Y.-L.; Tzeng, C.-C. *Chin. Pharm. J.* **2003**, *55*, 49.
- Chen, I.-L.; Chen, Y.-L.; Tzeng, C.-C. *Helv. Chim. Act.* **2002**, *85*, 2214.
- Huang, A.-C.; Lin, T.-P.; Kuo, S.-C.; Wang, J.-P. *J. Nat. Prod.* **1995**, *58*, 117.
- Chang, C.-P.; Lin, T.-P.; Wang, J.-P.; Kuo, S.-C. *Chin. Pharm. J.* **2001**, *53*, 239.
- Setzer, W. N.; Setzer, M. C.; Schmidt, J. M.; Moriarity, D. M.; Vogler, B.; Reeb, S.; Holmes, A. M.; Haber, W. A. *Planta Med.* **2000**, *66*, 493.
- Chen, K.-S.; Chang, Y.-L.; Teng, C.-H.; Chen, C.-F.; Wu, Y.-C. *Planta Med.* **2000**, *66*, 80.
- Chen, I.-S.; Chen, H.-F.; Cheng, M.-J.; Chang, Y.-L.; Teng, C.-M.; Tsutomu, I.; Chen, J.-J.; Tsai, I.-L. *J. Nat. Prod.* **2001**, *64*, 1143.
- Butenschon, I.; Moller, K.; Hansel, W. *J. Med. Chem.* **2001**, *44*, 1249.
- Tuppy, H.; Bohm, F. *Monatsh. Chem.* **1956**, *87*, 720.
- Rajakumar, D. V.; Rao, M. N. A. *Pharmazie* **1994**, *49*, 516.
- Smith, R. J.; Bowman, B. J.; Iden, S. S. *Immunology* **1981**, *44*, 677.
- Ochs, D. L.; Reed, P. W. *Biochem. Biophys. Res. Commun.* **1981**, *102*, 958.
- Gray, G. D.; Wickstrom, E. *BioTechniques* **1996**, *21*, 780.