

# Quantitative proteome analysis using D-labeled *N*-ethylmaleimide and $^{13}\text{C}$ -labeled iodoacetanilide by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry

Sadamu Kurono,<sup>a,b,d</sup> Tamie Kurono,<sup>a,d</sup> Naoka Komori,<sup>a</sup>  
Satomi Niwayama<sup>b,c,\*</sup> and Hiroyuki Matsumoto<sup>a,\*</sup>

<sup>a</sup>*Department of Biochemistry and Molecular Biology, University of Oklahoma Health Sciences Center, Oklahoma City, OK 73190, USA*

<sup>b</sup>*Department of Chemistry, Oklahoma State University, Stillwater, OK 74078, USA*

<sup>c</sup>*Department of Chemistry and Biochemistry, Texas Tech University, Box 41061, Lubbock, TX 79409-1061, USA*

<sup>d</sup>*IBERICA HOLDINGS Co., Ltd, Kurume University Translational Research Center, Kurume, Fukuoka 830-0011, Japan*

Received 16 August 2006; revised 12 September 2006; accepted 12 September 2006

Available online 16 October 2006

**Abstract**—A new methodology for quantitative analysis of proteins is described, applying stable-isotope labeling by small organic molecules combined with one- or two-dimensional electrophoresis and MALDI-TOF-MS, also allowing concurrent protein identification by peptide mass fingerprinting. Our method eliminates fundamental problems in other existing isotope-tagging methods requiring liquid chromatography and MS/MS, such as isotope effects, fragmentation, and solubility. It is also anticipated to be more practical and accessible than those LC-dependent methods.

© 2006 Elsevier Ltd. All rights reserved.

## 1. Introduction

Proteomics is rapidly becoming an important research area for comprehensive study of protein expression patterns under a specific set of conditions. A great deal of effort has been focused on developing methods of measuring global changes in relative protein abundance between two distinct cell systems, such as control versus treated cells. Thus, it is essential to develop efficient methodologies for quantitative analysis of proteins within complex mixtures expressed under certain physiological conditions.

Conventional methods for quantitative analysis of proteins include two-dimensional densitometry of gels<sup>1</sup> and radioisotope labeling.<sup>2</sup> While these methods are still useful for certain purposes, more recently, application of stable-isotope labeling followed by mass spectrometry

analysis has been emerging as a powerful technology for quantification and concurrent identification of proteins for proteomics research.<sup>3</sup>

Some of the earliest examples are metabolic labeling methods in which cells are cultured in isotope-enriched or normal media and the relative abundance of specific proteins is quantitatively analyzed from peak intensities from each sample in the mass spectra.<sup>4,5</sup> The disadvantage of metabolic labeling methods in general is that, although they are applicable to cultured cells or small organisms, it is rather difficult to apply to label larger organisms.

Instead, chemical modifications by covalent labeling on specific amino acid residues using isotope-labeled reagents followed by mass spectrometry analysis are expected to significantly expand the scope of isotope-labeling methods, because in theory, such methods are applicable to any protein samples.

Some of the most pioneering work has been reported by Aebersold et al., applying deuterium-labeled isotope-coded affinity tags (ICATs).<sup>6</sup> The first generation of ICAT reagents is a set of reagents

**Keywords:** Quantitative analysis; Proteomics; Stable-isotope labeling; Cysteine modifier.

\* Corresponding authors. Tel.: +1 806 742 3118; fax: +1 806 742 1289; e-mail addresses: [satomi.niwayama@ttu.edu](mailto:satomi.niwayama@ttu.edu); [hiroyuki-matsumoto@ouhsc.edu](mailto:hiroyuki-matsumoto@ouhsc.edu)

that consists of a biotin conjugate of a well-known cysteine-modifying reagent, iodoacetamide, and its deuterated version, whose molecular weight is 8 Da heavier.

In this method, proteins are modified with the cysteine-specific reactive group (iodo group) and the biotin tag in the ICAT reagent allows the specific isolation of the modified Cys-containing peptides by immobilized avidin. Changes in the relative abundance of peptides from distinct proteome samples are accomplished by the use of isotopically labeled and unlabeled ICAT reagents, which show chemically identical behavior. After the derivatized proteomes are pooled and digested with trypsin, the tryptic peptide mixtures isolated by avidin affinity chromatography can be quantitatively analyzed by liquid chromatography/electrospray ionization-tandem mass spectrometry (LC/ESI-MS/MS). Subsequently many other researchers reported other LC-dependent methods inspired by this ICAT method.<sup>7</sup> Coupled with recent development of multidimensional liquid chromatography, these LC-based methods have been playing important roles in proteomics.


However, several fundamental problems with this ICAT method have been reported, such as primary isotope effects especially for D-labeling, causing differential elution during microcapillary reversed-phase liquid chromatography ( $\mu$ LC), and fragmentation of the labels during collision-induced dissociation (CID) conditions, complicating the interpretation of tandem mass spectra. These problems are likely derived from the use of large hydrophobic organic molecules, which have decreased solubility in the aqueous media. Although several improvements have been made for the later versions of ICAT reagents, such as acid- or photo-cleavable versions,<sup>8</sup> and other different types of labeling reagents have also been reported,<sup>7</sup> these LC-dependent methods essentially require a large amount of aqueous media for purification of the derivatized peptides. For this

reason, it is imperative to develop fundamentally different approaches that use small-molecule labeling reagents and that eliminate the liquid chromatography step.


Therefore, we have been developing our own methodology for quantitative analysis of proteins by a combination of isotope-labeled and -unlabeled chemical modification of specific amino acid residues by small organic molecules followed by 2D electrophoresis and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF-MS). The concept is outlined in [Scheme 1](#).

Since our method uses 2D electrophoresis instead of LC for the purification of protein, the procedure is more convenient and more economical, and the small organic molecules are also expected to be more easily accessible as well as to have better solubility in the sample mixture. The use of small molecules and a soft ionization mass spectrometry, MALDI-MS, also eliminates the fragmentation problems. While the choice of mass spectrometry is not critical and either MALDI or electrospray ionization (ESI) is possible, we chose MALDI rather than ESI, since more laboratories tend to have MALDI and the operation and maintenance are easier, making our method more practical.

The 2D polyacrylamide gel electrophoresis (2D PAGE), which displays all separated proteins on a single gel, has been the standard tool for protein expression in proteomics. Two-dimensional gel separation using ICAT reagents has been reported.<sup>9</sup> The 2D gel electrophoresis also provides information about the molecular weights and *pI* values of proteins. Therefore our method allows us to identify each protein through peptide mass fingerprinting prior to quantitative analysis without a need for the MS/MS mode for identification of proteins, which will not only eliminate the fragmentation problems, but also make it simpler and more economical and therefore accessible to more researchers.



**Scheme 1.** Our strategy for quantitative analysis of proteins.



**Scheme 2.** Reactions of *N*-ethylmaleimide (NEM) or iodoacetanilide (IAA) with cysteine.

We previously synthesized sulfhydryl (–SH)-group specific modifiers that are small molecules for quantitative analysis of proteins, since a sulfhydryl group is one of the most nucleophilic functional groups within peptide sequences. The combinations we have synthesized are *N*-ethylmaleimide (NEM) and its  $d_5$ -deuterated versions (NEM- $d_5$ )<sup>10</sup> as well as iodoacetanilide (IAA) and  $^{13}C_6$ -labeled iodoacetanilide ( $^{13}C_6$ -IAA).<sup>11</sup> Iodoacetanilide (IAA) is a derivative of a well-known sulfhydryl group modifier, iodoacetamide, and both NEM and IAA are known to specifically react with –SH groups of cysteine residues (Scheme 2). The unlabeled *N*-ethylmaleimide is commercially available.

As both modifiers react at different pHs, these modifiers may be used according to the sample environment. The iodoacetanilides are expected not to interfere with phosphine-based reducing reagents,<sup>12</sup> and are anticipated to be more reactive than *N*-ethylmaleimides due to the existence of a reactive leaving group, an iodo group. The mass difference between IAA and  $^{13}C_6$ -IAA is 6 Da rather than 5 Da as in NEM and NEM- $d_5$ ; therefore the combination of IAA and  $^{13}C_6$ -IAA is expected to provide sharper separation of the IAA- and  $^{13}C_6$ -IAA-modified peptide peaks in the MALDI spectra rather than the NEM- and NEM- $d_5$ -modified ones, and hence more accurate quantification, while both combinations are quite visible in the mass spectra separated by 1D and 2D electrophoresis.

We have shown that both combinations are successful for quantitative analysis of various peptides.<sup>10,11</sup> Using known amounts of various peptides, we confirmed that ionization efficiencies for isotope-labeled and -unlabeled peptides are identical within experimental errors. Here we show their application to proteomic experiments, quantitative analysis of proteins.

## 2. Results and discussion

### 2.1. Quantitative analysis of standard proteins for 1D SDS-PAGE


In order to test the applicability of our method to proteins, first we applied it to quantitative analysis of four commercially available proteins:  $\alpha$ -lactalbumin, bovine serum albumin (BSA), carbonic anhydrase, and ovalbumin, each of which was purified by 1D electrophoresis. These proteins were individually reacted with NEM, NEM- $d_5$ , IAA, or  $^{13}C_6$ -IAAs in the presence of a reducing reagent, tributylphosphine, which reduces intramolecular S–S bonds to sulfhydryl groups. Figure 1a shows the MALDI spectrum of the mixture of  $\alpha$ -lactalbumin modified with NEM or NEM- $d_5$  (NEM-modified/NEM- $d_5$ -modified = 1:1) loaded onto 1D SDS-PAGE after tryptic digestion.

Seven pairs of tryptic peptides modified with NEM or NEM- $d_5$  that were 5 Da apart with nearly equal intensities were detected, which are indicated by the arrows in Figure 1a. These seven pairs of peptides reacted with NEM or NEM- $d_5$  include peptides with unknown amino acid sequences and peptides formed by cleavage of other longer peptides (data not shown). Figure 1b shows the enlarged peak area of the NEM- or NEM- $d_5$ -modified peptides of  $\alpha$ -lactalbumin consisting of residues 6–10, CEVFR, with five molar ratios, 9:1, 6:1, 3:1, 1:1, and 1:3. The correlation parameters and inclinations between the observed ratio and the theoretical ratio for the NEM- and NEM- $d_5$ -CEVFR were  $R^2 = 0.9984$  and inclination = 0.9781, as shown in Figure 1c and Table 1. Some peptides with unknown amino acid sequences may not contain any cysteines, but these data indicate that these peptides still showed reasonable correlation between the observed ratios and the theoretical ratios.

The same experiments were conducted for three other commercial proteins, and the correlation parameters and inclinations for these proteins modified with NEM or NEM- $d_5$  are summarized in Table 1 along with  $\alpha$ -lactalbumin. The three proteins other than carbonic anhydrase show the reasonable correlation parameters and inclinations. No NEM- and NEM- $d_5$ -modified peptide peaks were detected from carbonic anhydrase because this protein has no cysteine residues in its sequence.

Figure 2a shows the expanded MALDI-MS spectra of tryptic peptides digested from bovine serum albumin (BSA) unmodified (upper) and modified with IAA or  $^{13}C_6$ -IAA (IAA-modified/ $^{13}C_6$ -IAA-modified = 1:1) (lower).

In a comparison of the two spectra, many modified peaks were observed as pairs having the same intensities. Some tryptic peptides reacted with more than one equivalent of IAAs or  $^{13}C_6$ -IAA, but these IAA-modified and  $^{13}C_6$ -IAA-modified peptides still showed the same intensities. Around 90% of cysteine residues (31 out of 35) in BSA were found to be modified with IAA or  $^{13}C_6$ -IAA, and they are listed in Table 2.



**Figure 1.** (a–c) MALDI spectrum of  $\alpha$ -lactalbumin modified with NEM or NEM- $d_5$ .


**Table 1.** Results of quantitative analysis of commercial proteins with NEM and NEM- $d_5$  after 1D SDS-PAGE

Protein	Number of Cys	$R^2$	Inclination
Bovine serum albumin	35	0.9998	1.1036
Ovalbumin	6	0.9928	1.2822
Carbonic anhydrase	0	ND <sup>a</sup>	ND <sup>a</sup>
$\alpha$ -Lactalbumin	8	0.9984	0.9781

<sup>a</sup> ND, not detectable.

Figure 2b shows the enlarged peak area around  $m/z$  1800–2000 in Figure 2a. The peaks of peptides consisting of residues 508–523, RPCFSALTPDETYVPK, that were modified with IAA or  $^{13}\text{C}_6$ -IAA are clearly seen at  $m/z$  1957.0 or 1963.0, respectively, and many other modified peaks were clearly detected with high intensities. As one example, Figure 3a shows

expansion of the peak areas around 1950–1970 Da, in which the IAA- and  $^{13}\text{C}_6$ -IAA-modified peptide residues RPCFSALTPDETYVPK appear in various ratios, and Figure 3b shows the correlation between the theoretical and observed ratios calculated from these results. As in this scheme, we also observed that the modified peak intensity ratios between these pair peaks well represent



**Figure 2.** (a and b) MALDI spectrum of bovine serum albumin (BSA) modified with IAA or  $^{13}\text{C}_6$ -IAA (IAA-modified/ $^{13}\text{C}_6$ -IAA-modified = 1:1).

the five molar ratios of the loaded BSAs modified with IAA or  $^{13}\text{C}_6$ -IAA (IAA-modified/ $^{13}\text{C}_6$ -IAA-modified = 9:1, 6:1, 3:1, 1:1, and 1:3). The correlation parameters and inclinations between the observed ratio and the theoretical ratio for this peptide RPCFSALTPDETYVPK were  $R^2 = 0.9996$  and inclination = 0.9678 as shown in Figures 3a and b, and Table 3.

The correlation parameters and inclinations for the other commercial proteins modified with IAA or  $^{13}\text{C}_6$ -IAA are summarized in Table 3 as well. The three proteins other than carbonic anhydrase show the good correlation parameters and inclinations. As in the case of NEM and NEM- $d_5$ , no IAA- and  $^{13}\text{C}_6$ -IAA-modified peptide peaks were detected from carbonic anhydrase due to non-existence of a cysteine residue.

As in the peptide at  $m/z$  1895.0 in Figure 2b, sometimes acrylamide-reacted peptides are detected when SDS-PAGE is performed, however, such peaks were rarely observed in these NEM, NEM- $d_5$ , IAA, or  $^{13}\text{C}_6$ -IAA-modified proteins. While acrylamide is occasionally used as a modifier,<sup>13</sup> this observation may mean that IAA and  $^{13}\text{C}_6$ -IAA are more reactive than acrylamide toward cysteine residues and hence more suitable cysteine modifiers.

From these results we conclude that both characterization and quantification of commercially available proteins are possible by the use of NEM and NEM- $d_5$ , or IAA and  $^{13}\text{C}_6$ -IAA combined with 1D SDS-PAGE and MALDI-MS. It should also be noted that Corthals et al. have reported successful application of a combina-

tion of IAA and IAA- $d_5$  to quantitative analysis of commercially available proteins, bovine serum albumin (BSA) and lactoperoxidase, separated by 1D SDS PAGE.<sup>14</sup>

## 2.2. Quantitative analysis of protein extracts from *Drosophila* heads for 2D IEF/SDS-PAGE


Next, we applied our method to quantitative analysis of the above four proteins separated by 2D electrophoresis, and confirmed that our method is also applicable to these commercial proteins as in the experiments by 1D electrophoresis described above (data not shown). Therefore we applied our method to quantitative analysis of proteins from the extract obtained from *Drosophila* heads that were separated by 2D electrophoresis. The protein extracts from 100 *Drosophila* heads were split into a set of two solutions in the ratios of 3:1, 1:1, and 1:3, and were reacted with an excess of NEM, NEM- $d_5$ , IAA, or  $^{13}\text{C}_6$ -IAA in the presence of the reducing agent, tributylphosphine, as in the experiment with 1D electrophoresis above. We then performed quantitative analysis of many proteins in these protein samples developed on 2D IEF/SDS-PAGE described in Section 4. We also identified these proteins by peptide mass fingerprinting concurrently with or prior to the quantitative analysis. All the protein samples were developed under the same conditions. Although occasionally poor reproducibility has been pointed out for 2D electrophoresis, all the 2D gel images were highly superimposable throughout our experiments. Although recently thiourea is frequently used for performing 2D electrophoresis, we used urea instead in order to avoid

**Table 2.** Tryptic peptide peaks of BSA modified with IAA or  $^{13}\text{C}_6$ -IAA (BSA-IAA/BSA- $^{13}\text{C}_6$ -IAA = 1:1)

Number of peptides	Observed $m/z$ value of IAA-modified peptide	Observed $m/z$ value of $^{13}\text{C}_6$ -IAA-modified peptide	Theoretical <sup>a</sup> $m/z$ value of unmodified peptide	Identified peptide from database		Number of cysteines	Number of added IAAs ( $^{13}\text{C}_6$ -IAAs)
				Position on protein	Sequence		
1	655.3	661.3	522.3	Unknown	Unknown	Unknown	1
2	974.5	980.5	841.5	483–489	LCVLHEK	1	1
3	1110.5	1116.5	977.5	123–130	NECFLSHK	1	1
4	1144.5	1150.5	1011.4 (low) <sup>b</sup>	413–420	QNCDQFEK	1	1
5	1148.6	1154.5	1015.5	310–318	SHCIAVEVK	1	1
6	1157.6	1163.6	1024.5	499–507	CCTESLVNR	2	1
7	1290.6	1302.7	1024.5	499–507	CCTESLVNR	2	2
8	1183.7	1189.6	1050.5 (low)	588–597	EACFAVEGPK	1	1
9	1185.6	1191.7	1052.5 (low)	460–468	CCTKPHSER	2	1
10	1318.7	1330.6	1052.5 (low)	460–468	CCTKPESER	2	2
11	1443.8	1455.8	1177.6 (ND) <sup>c</sup>	300–309	ECCDKPLLEK	2	2
12	1615.8	1627.8	1349.5 (ND)	76–88	TCVADESHAG CEK	2	2
13	1495.9	1501.7	1362.7 (ND)	89–100	SLHILFGDELCK	1	1
14	1630.8	1642.7	1364.5 (ND)	106–117	ETYGDMAD CCEK	2	2
15	1519.8	1525.7	1386.6 (ND)	286–297	YICDNQDTISSK	1	1
16	1654.8	1666.9	1388.6 (ND)	375–386	EYEATLEE CCAK	2	2
17	1630.8	1636.7	1497.6 (ND)	387–399	DDPHACYSTVFDK.	1	1
18	1652.9	1658.8	1519.7 (ND)	139–151	LKPDPTNL CDEFK	1	1
19	1844.9	1856.8	1578.6 (ND)	267–280	ECCHGDLLE CADDR	3	2
20	1977.8	1995.8	1578.6 (ND)	267–280	ECCHGDLLE CADDR	3	3
21	1899.8	1911.8	1633.7 (ND)	184–197	YNGVFQE CCQAEDK	2	2
22	1801.0	1806.9	1667.8 (ND)	469–482	MPCTEDYLSLILNR	1	1
23	1817.0	1823.0	1683.8 (ND)	469–482 1Met-Ox <sup>d</sup>	MPCTEDYLSLILNR	1	1
24	1854.9	1860.9	1721.9 (ND)	Unknown	Unknown	Unknown	1
25	1882.9	1889.0	1749.9 (ND)	Unknown	Unknown	Unknown	1
26	1889.0	1894.8	1756.0 (ND)	Unknown	Unknown	Unknown	1
27	2155.9	2174.0	1756.7 (ND)	581–597	CCAADDKEA CFAVEGPK	3	3
28	1957.0	1963.0	1823.9	508–523	RPCFSALTPDETYVPK	1	1
29	2534.3	2540.3	2401.2 (ND)	319–340	DATPENLPPLTADFAEDKDV CK	1	1

<sup>a</sup> Theoretical  $m/z$  value of unmodified peptide: calculated from the peptide sequence or the observed  $m/z$  value of IAA/ $^{13}\text{C}_6$ -IAA-modified peptides.<sup>b</sup> Low, low intensity.<sup>c</sup> ND, not detectable.<sup>d</sup> Met-Ox, methionine is oxidized.






**Figure 3.** (a and b) Quantitative analysis of bovine serum albumin (BSA) by 1D SDS-PAGE.

**Table 3.** Results of quantitative analysis of commercial proteins with IAA and  $^{13}\text{C}_6$ -IAA after 1D SDS-PAGE

Protein	Number of Cys	$R^2$	Inclination
Bovine serum albumin	35	0.9996	0.9678
Ovalbumin	6	0.9999	1.0044
Carbonic anhydrase	0	ND <sup>a</sup>	ND <sup>a</sup>
$\alpha$ -Lactalbumin	8	0.9997	0.9931

<sup>a</sup> ND, not detectable.



**Figure 4.** 2D IEF/SDS-PAGE of *Drosophila* head proteins: mixture of IAA-modified and  $^{13}\text{C}_6$ -IAA-modified proteins in the proportion of 3:1.

potential side reactions between thiourea and these tagging reagents.


For example, **Figure 4** is the 2D IEF/SDS-PAGE gel of the mixture of the above protein samples in the ratio of 3:1 that were reacted with an excess of IAA or  $^{13}\text{C}_6$ -IAA.

The ratio reflects the amount of the sample protein reacted with IAA and  $^{13}\text{C}_6$ -IAA, namely, IAA-reacted

protein/ $^{13}\text{C}_6$ -IAA-reacted protein = 3:1 in this case. **Figure 5a** and **b** shows the MALDI spectra of the tryptic peptides digested from the protein labeled No. 3 (tropomyosin 1) in the above protein sample prepared by mixing the NEM-reacted protein extract and the NEM- $d_5$ -reacted protein extract in the ratio of 3:1 (NEM-reacted/NEM- $d_5$ -reacted) (**Fig. 5a**) and that prepared by mixing the IAA-reacted protein extract and the  $^{13}\text{C}_6$ -IAA-reacted protein extract in the ratio of 3:1 (IAA-reacted/ $^{13}\text{C}_6$ -IAA-reacted) (**Fig. 5b**), respectively.

These spectra indicate that the proteins extracted from living organisms react with NEM, NEM- $d_5$ , IAA, or  $^{13}\text{C}_6$ -IAA successfully in the presence of tributylphosphine, while some differences in reactivity between the proteins extracted from *Drosophila* heads and the commercial proteins can be pointed out. Among the proteins developed on 2D IEF/SDS-PAGE, eight proteins identified through peptide mass fingerprinting and their quantitative analysis using IAA and  $^{13}\text{C}_6$ -IAA are summarized in **Table 4**.

In a comparison of the observed and the theoretical ratios between IAA- and  $^{13}\text{C}_6$ -IAA-modified tryptic peptides, each analytical graph at each ratio shows reasonable linearity and inclination values. Among the eight proteins in **Table 4**, mitochondrial porin has no cysteine residues. Therefore as in the above results in 1D electrophoresis with commercially available proteins, no IAA- or  $^{13}\text{C}_6$ -IAA-modified peptide peaks were detected from this protein (data not shown). As in the case of the commercial proteins purified by 1D electrophoresis, acrylamide-reacted peptides were rarely observed after this 2D electrophoresis.<sup>15</sup>



**Figure 5.** (a and b) MALDI spectrum of spot 3 (tropomyosin 1). (a) NEM-modified/NEM-*d*<sub>5</sub>-modified = 3:1. (b) IAA-modified/<sup>13</sup>C<sub>6</sub>-IAA-modified = 3:1.

**Table 4.** Results of quantitative analysis of proteins from *Drosophila* heads separated by 2D IEF/SDS-PAGE

Spot no.	Protein candidate	Number of Cys	<i>R</i> <sup>2</sup>	Inclination
1	CG3612-PA	4	0.9997	1.0647
2	Enolase	4	1.0	0.9467
3	Tropomyosin 1	1	0.9989	1.092
4	Arginine kinase	5	0.9995	1.0457
5	Aldolase	4	1.0	1.0143
6	Tropomyosin 2	3	0.998	0.9542
7	Mitochondrial porin	0	ND <sup>a</sup>	ND <sup>a</sup>
8	Alcohol dehydrogenase	2	0.9953	1.017

<sup>a</sup> ND, not detectable.

Although NEM and IAA were both sufficiently reactive and hence equally efficient modifiers for quantitative analysis of model proteins in the above experiments, we observed substantial differences in performance between these two reagents when they were applied to the 2D electrophoresis experiments on biological

samples. We were able to obtain reasonable quantitative results only from tropomyosin 1 when using NEM or NEM-*d*<sub>5</sub> (data not shown). Although we tried to increase the reaction time of the proteins with NEM or NEM-*d*<sub>5</sub> to 24 and 48 h, and also to raise the reaction temperature up to 37 °C, these results did not change



significantly. While the reason is unclear, one plausible reason is that NEM and NEM- $d_5$  are less reactive than IAA and  $^{13}\text{C}_6$ -IAA to cysteine residues to begin with. It may be pointed out that these *N*-alkylmaleimides are in general known to interfere with reducing reagents more than iodoacetamide derivatives.<sup>12</sup> In other words, iodoacetamide derivatives tend to react with sulfhydryl groups more specifically. The pH conditions that are known to be suitable for reactions of cysteines and iodoacetamide are around 8–9 while those suitable for reactions of cysteines with *N*-alkylmaleimide are around 7. This slightly basic pH may also contribute to enhancing the reactivity of sulfhydryl group as a nucleophile, and hence explain the higher reactivity of the iodoacetamide derivatives, IAA or  $^{13}\text{C}_6$ -IAA, than of the *N*-alkylmaleimide derivatives, NEM or NEM- $d_5$ . Some studies also indicate that the optimal pH for reduction of S–S bonds appears to be around 8.5–8.9,<sup>16</sup> which would also account for the enhanced reactivity of the cysteines with IAA or  $^{13}\text{C}_6$ -IAA. However, it is reasonable to exclude the conditions of 2D electrophoresis as a cause, because this method was successful with either combination, NEM/NEM- $d_5$  or IAA/ $^{13}\text{C}_6$ -IAA, for the commercial proteins separated by 2D electrophoresis. Another explanation may be attributed to the buffer solutions. Since some components of the lysis buffer to prepare the protein extract from *Drosophila* heads and the buffer to prepare commercial protein solutions are different, it is possible that the components in the buffer for the protein extracts from *Drosophila* heads may have interfered with the less specific *N*-alkylmaleimides under our experimental conditions before the *N*-alkylmaleimides reacted with sulfhydryl groups. It is also possible that since commercial proteins are relatively pure the protein concentration was higher than that of the proteins from *Drosophila* heads, while the protein extracts from *Drosophila* heads have many ingredients, which may also have interfered with the less specific *N*-alkylmaleimides. We are examining the possible application of our method to other protein extract samples to obtain further information about its applicability.

From these results we conclude that both characterization and quantification of most proteins are possible by the use of IAA and  $^{13}\text{C}_6$ -IAA combined with 2D IEF/SDS–PAGE and MALDI-MS.

### 3. Summary and conclusions

We have demonstrated that the combination of isotope-labeled and -unlabeled small organic molecules, NEM and NEM- $d_5$  as well as IAA and  $^{13}\text{C}_6$ -IAA, and MALDI-MS with 1D or 2D electrophoresis is applicable to quantitative analysis of commercial proteins as well as proteins extracted from *Drosophila* heads. Ionization efficiencies of the peptides do not appear to be significantly influenced by modifications with these reagents, even though some of the peptides show slightly decreased ionization efficiency, possibly caused by their modification. Both combinations have proven to exhibit differences in the molecular mass (5 and 6 Da) sufficient to overcome overlapping problems caused by the monoisotopic peaks

from the isotope-labeled compounds and the isotopic peaks from unlabeled compounds, as we had shown in similar experiments with peptides.<sup>10,11</sup> Earlier, Sechi,<sup>17</sup> Hamdan et al.,<sup>18</sup> and Schrattenholz et al.<sup>19</sup> independently reported their methods for quantitative analysis of proteins separated by 1D or 2D electrophoresis applying  $d_0$ - and  $d_3$ -acrylamides as cysteine modifiers. Another set of small-molecule cysteine modifiers,  $d_0$ - and  $d_4$ -vinylpyridine,<sup>20</sup> is also anticipated to be equally useful in combination with 2D electrophoresis. However, from our previous study using methylmaleimide- $d_3$ ,<sup>10</sup> we conclude that at least 5 Da is necessary to sufficiently overcome this overlapping problem.<sup>21</sup> Consisting of only three carbons, acrylamide limits possibilities for further synthetic modification to yield a greater difference in the molecular mass. Our simple synthetic manipulation offers an attractive approach for further modifying various types of modifiers.

Although recently a number of proteomic methods for quantitative analysis of proteins have been reported, most methods are based on liquid chromatography for separation, purification, and observation of protein expression profiles, and proteomic methods based on the use of 2D electrophoresis as a means to obtain protein expression profiles are rather limited.<sup>22</sup> The major reasons for the preference are the ease with which LC-based methods adjust to high-throughput systems as well as the ability of these methods to deal with a wider range of proteins than 2D electrophoresis does. However, 2D PAGE has been the most mature technique for obtaining protein expression patterns, separating thousands of proteins. As there is no single proteomic procedure useful to comprehend all the necessary data, many different technologies are required for analysis of various features of proteins for proteomics.

Our method revealed several advantageous points over LC-dependent methods. First, we did not observe any isotope effects, although they have been common and fundamental problems in LC-dependent methods such as ICAT. We conclude that the reactivities of these isotope-labeled and -unlabeled modifiers are identical. Therefore, even if several cysteine residues are rather poorly reactive, they do not affect the quantitative analysis of proteins because the reactivity toward the isotope-labeled or -unlabeled modifier is identical. In fact, no particular discrepancies in the ratios were observed from different cysteine residues within the same protein. We think that this outcome is due to the use of small organic molecules which are less hydrophobic and therefore more soluble in SDS–PAGE conditions. When there is more than one cysteine residue, these residues provide multiple data points for quantitative analysis, ensuring accuracy. Furthermore, we identify each protein through peptide mass fingerprinting prior to quantitative analysis, and therefore even when non-specific binding is detected between non-sulfhydryl groups and these labeling reagents, such nonspecific binding can simply be ignored. We have demonstrated that even in the presence of potential nonspecific binding, the results of the quantitative analysis still indicated reasonable correlation between the theoretical and observed ratios in the above experiments. This is a signifi-

cant advantage over the ICAT method in which protein identification is conducted only at the end of the entire process by tandem mass spectrometry, while in our method tandem mass spectrometry is optional for confirmation of the proteins identified through peptide mass fingerprinting.

Recent comparative studies of commonly used proteomic quantitative methods indicate significant discrepancies among the quantitative data obtained by the DIGE methods and two LC-dependent methods that use cleavable ICAT or iTRAQ.<sup>23</sup> Other studies on the shotgun approach in protein identification by microcapillary liquid chromatography-tandem mass spectrometry ( $\mu$ LC/MS/MS) indicate that the efficacy of digestion heavily depends on the protocols followed in this approach.<sup>24</sup> These results also suggest the importance of an alternative approach by electrophoresis and small-molecule isotope-labeling.

Therefore, depending on the types of the sample proteins that contain cysteines and the purpose of the projects, our method is anticipated to complement other existing LC-dependent methods and to be applicable to quantitative proteomics in many types of projects, its greater applicability thus making it more readily accessible to diverse laboratories. We are currently synthesizing other modifiers that are small organic molecules of which the differences of molecular mass between isotope-labeled and -unlabeled versions are greater than 6 Da, and investigating the possibility of expanding our method toward these reagents as well as those that modify non-cysteine-containing proteins, and further studies will be reported in due course.

## 4. Experimental methods

### 4.1. Preparation of standard protein solutions for 1D SDS-PAGE (without modifiers)

Each of the following four commercial proteins (1 mg each, purchased from Sigma), bovine serum albumin (66 kDa), egg albumin (45 kDa), carbonic anhydrase (29 kDa), or  $\alpha$ -lactalbumin (14.2 kDa), was dissolved in a sample buffer (1 mL) containing Trizma<sup>®</sup> Pre-Set Crystals (purchased from Sigma), ethylenediaminetetraacetic acid (EDTA), sodium dodecyl sulfate (SDS), and  $\beta$ -mercaptoethanol. The pHs and the concentrations of Trizma<sup>®</sup>, EDTA, SDS, and  $\beta$ -mercaptoethanol were as follows. For the Trizma<sup>®</sup> Pre-Set Crystals the initial pH was 7.0 and the final concentration was 78 mM. For the EDTA, the final concentration was 6.3 mM. For the SDS, the final concentration was 2.5% (w/v). For the  $\beta$ -mercaptoethanol, the final concentration (v/v) was 1.25%. The mixed protein solution was left for 1 h at room temperature.

### 4.2. Preparation of standard protein solutions for 1D SDS-PAGE (with NEM/NEM-*d*<sub>5</sub>)

Each of the following four commercial proteins (1 mg each, purchased from Sigma), bovine serum albumin

(66 kDa), egg albumin (45 kDa), carbonic anhydrase (29 kDa), or  $\alpha$ -lactalbumin (14.2 kDa), was dissolved in a sample buffer (1 mL) containing Trizma<sup>®</sup> Pre-Set Crystals (Sigma), EDTA, SDS, and tributylphosphine (TBP). The pHs and the concentrations of the Trizma<sup>®</sup>, EDTA, SDS, and TBP were as follows. For the Trizma<sup>®</sup> Pre-Set Crystals, the initial pH and the final concentration were 7.8 and 84 mM. For the EDTA, the final concentration was 6.8 mM. For the SDS, the final concentration was 2.7% (w/v). For the tributylphosphine (TBP), the final concentration was 20 mM. The mixed protein solution was left for 1 h at room temperature for sufficient reduction with TBP.

Each of the above protein solutions (10  $\mu$ L) with different molar ratios was incubated with 1  $\mu$ L of 1 M NEM or NEM-*d*<sub>5</sub> DMSO solution, and these two solutions were mixed and incubated for 2 h at 37 °C. In this way, the following five molar ratio pairs of protein solutions were examined: 9:1, 6:1, 3:1, 1:1, and 1:3 (NEM-modified/NEM-*d*<sub>5</sub>-modified).

### 4.3. Preparation of standard protein solutions for 1D SDS-PAGE (with IAA/<sup>13</sup>C<sub>6</sub>-IAA)

Each of the following four commercial proteins (1 mg each, purchased from Sigma), bovine serum albumin (66 kDa), egg albumin (45 kDa), carbonic anhydrase (29 kDa), or  $\alpha$ -lactalbumin (14.2 kDa), was dissolved in a sample buffer (1 mL) containing Trizma<sup>®</sup> Pre-Set Crystals (Sigma), EDTA, SDS, and tributylphosphine (TBP). The pHs and the concentrations of the Trizma<sup>®</sup>, EDTA, SDS, and TBP were as follows. For the Trizma<sup>®</sup> Pre-Set Crystals, the initial pH was 9.0 and the final concentration was 84 mM. For the EDTA, the final concentration was 6.8 mM. For the SDS, the final concentration was 2.7% (w/v). For the tributylphosphine (TBP), the final concentration of was 20 mM. The mixed protein solution was left for 1 h at room temperature for sufficient reduction with TBP.

Each of the above protein solutions (10  $\mu$ L) with different molar ratios was incubated with 1  $\mu$ L of 200 mM IAA or <sup>13</sup>C<sub>6</sub>-IAA DMSO solution, and the two solutions were mixed and incubated for 2 h at 37 °C. In this way, the following five molar ratio pairs of protein solutions were examined: 9:1, 6:1, 3:1, 1:1, and 1:3 (NEM-modified/NEM-*d*<sub>5</sub>-modified or IAA-modified/<sup>13</sup>C<sub>6</sub>-IAA-modified).

### 4.4. Separation of proteins in 1D SDS-PAGE

The following SDS-PAGE gels were used: 0.8-mm thick gels (17-cm wide) that consist of 5% acrylamide in stacking gels (2-cm long) and 11% acrylamide in separating gels (13.5-cm long). Right before loading the samples, 20  $\mu$ L of the protein mixture solutions was mixed with 5  $\mu$ L of a solution consisting of 20% (v/v) glycerol and 0.1% (w/v) bromophenol blue, and the solution was subjected to SDS-PAGE. The electrophoresis conditions were as follows: 10 mA for about 1 h and 30 min, until the bromophenol blue reached the end of the stacking

gel, followed by 25 mA for about 2 h and 30 min until the bromophenol blue reached the end of the separating gel.

#### 4.5. Preparation of protein extracts from *Drosophila* heads for 2D IEF/SDS–PAGE (without modifiers)

A total of 100 precisely dissected *Drosophila* heads, excluding eyes, were homogenized using 120  $\mu$ L of a lysis buffer consisting of 8.5 M urea, 2% 3-[3-(cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS), 0.5% Bio-Lyte 3/10 ampholyte (Bio-Rad), and 5%  $\beta$ -mercaptoethanol. After 1 h, the supernatant was collected by centrifugation, and 100  $\mu$ L was loaded onto the IEF.

#### 4.6. Preparation of protein extracts from *Drosophila* heads for 2D IEF/SDS–PAGE (with NEM/NEM- $d_5$ )

A total of 100 precisely dissected *Drosophila* heads, excluding eyes, were homogenized using 80–120  $\mu$ L of a lysis buffer consisting of 47.25 mM Trizma® Pre-Set Crystals, pH 7.0, including 8.5 M urea, 2% CHAPS, 0.5% Bio-Lyte 3/10 ampholyte (Bio-Rad), and 20 mM TBP. After 1 h, the supernatant collected by centrifugation was divided into the volume ratios of 3:1, 1:1 or 1:3, and then the above lysis buffer was added to the smaller amount of supernatant in each pair to make the volumes equal. Each solution in each pair was incubated with 2  $\mu$ L of 1 M NEM or NEM- $d_5$  DMSO solution for 2 h at 37 °C, and subsequently 47.5  $\mu$ L from each solution was mixed. After 5  $\mu$ L of  $\beta$ -mercaptoethanol was added to the mixture and incubated for 10 min at room temperature to stop the reaction with NEMs, the mixture (a total of 100  $\mu$ L) was loaded onto the IEF.

#### 4.7. Preparation of protein extracts from *Drosophila* heads for 2D IEF/SDS–PAGE (with IAA/ $^{13}\text{C}_6$ -IAA)

A total of 100 precisely dissected *Drosophila* heads, excluding eyes, were homogenized using 80–120  $\mu$ L of a lysis buffer consisting of 47.25 mM Trizma® Pre-Set Crystals, pH 9.0, for incubation with IAA or  $^{13}\text{C}_6$ -IAA, including 8.5 M urea, 2% CHAPS, 0.5% Bio-Lyte 3/10 ampholyte (Bio-Rad), and 20 mM TBP. After 1 h, the supernatant collected by centrifugation was divided into the volume ratios of 3:1, 1:1 or 1:3, and then the above lysis buffer was added to the smaller amount of supernatant to make the volumes equal. Each solution in each pair was incubated with 2  $\mu$ L of 200 mM IAA or  $^{13}\text{C}_6$ -IAA DMSO solution for 2 h at 37 °C, and subsequently 47.5  $\mu$ L from each solution was mixed. After 5  $\mu$ L of  $\beta$ -mercaptoethanol was added to the mixture and incubated for 10 min at room temperature to stop the reaction with IAAs, the mixture (a total of 100  $\mu$ L) was loaded onto the IEF.

#### 4.8. Separation of proteins in 2D IEF/SDS–PAGE

For IEF analysis, the Immobiline™ DryStrips (pH 3–10, 13-cm long, 3-mm wide, and 0.5-mm thickness after rehydration, Amersham Biosciences) were rehydrated overnight with the lysis buffer including 5%

$\beta$ -mercaptoethanol (v/v) and 0.1% bromophenol blue aqueous solution. Each 90  $\mu$ L of four sets of samples was loaded into a sample cup and focused on IPGphor (Amersham Pharmacia Biotech), and then IEF was performed. The voltage was linearly increased from 0 to 500 V for 3 min, then increased up to 4000 V over 1 h and 30 min and held constant at 4000 V for an additional 30 min, and for 4 h at 8000 V with a maximum current per strip of 70  $\mu$ A. A total V h product rather than 17 kV h was used to focus the strip. The electrophoresis conditions for the second SDS–PAGE were as follows: 15 mA for about 1 h until the bromophenol blue reached the end of the stacking gel, followed by 25 mA for about 3 h, until the bromophenol blue reached the end of the separating gel.

#### 4.9. In-gel digestion

The protein bands (from 1D SDS–PAGE) or spots (from 2D IEF/SDS–PAGE) stained with Coomassie® Brilliant Blue G-250 (Bio-Rad) or Brilliant Blue R-250 (Sigma) were excised and destained with 50% acetonitrile/100 mM  $\text{NH}_4\text{HCO}_3$  (pH 8.0). After they were dried briefly, 2  $\mu$ L (1  $\mu$ L at a time, twice) of *N*-*p*-tosyl-L-phenylalanine chloromethyl ketone (TPCK)-treated trypsin (Promega) solution (0.2  $\mu$ g/ $\mu$ L in 50 mM acetic acid) was applied to each gel piece, followed by incubation in 50  $\mu$ L of 2.5 mM  $\text{NH}_4\text{HCO}_3$  overnight at 30 °C. After that, the incubated solution was collected and refrigerated. For extraction of the tryptic peptides, the gel piece was immersed in 60% acetonitrile/0.1% trifluoroacetic acid (TFA) and shaken for 1 h. The above refrigerated incubated solution and the extracted peptide solutions were mixed and dried. The dried tryptic peptides were dissolved in 5  $\mu$ L Milli-Q® water.

#### 4.10. MALDI-TOF-MS

The extracted peptide solution was mixed with a matrix ( $\alpha$ -cyano-4-hydroxycinnamic acid, CHCA) solution (prepared by dissolving 10 mg CHCA in 1 mL of 50% acetonitrile/0.1% TFA) and then subjected to MALDI-TOF-MS analysis. The MALDI spectra were obtained by a Voyager Elite BioSpectrometry Research Station, equipped with a delayed extraction option (Applied Biosystems) operated at a 20 kV accelerating voltage, 75% grid voltage, 0.05% guide wire voltage, 150 ns pulse delay time, 100 mV vertical scale, and 3% vertical offset. A pulsed nitrogen laser operating at 337 nm was used as a desorption/ionization source. Mass spectrometry was performed in a reflector with positive ion detection. The ion signal was recorded using a 500 MHz transient digitizer. The data were analyzed using GRAMS/386 (Galactic Industries Corporation).

#### 4.11. Database search

The monoisotopic *m/z* values of tryptic peptide peaks were entered into the MS-Fit in the ProteinProspector (UCSF: <http://prospector.ucsf.edu/>) for a database search for characterization of proteins.



#### 4.12. Peak area calculation

The areas of the monoisotopic peak and the following four isotopic peaks were added together for each modified peptide using GRAMS/386. The areas of the small 6- or 7–10th isotopic peaks of NEM- or IAA-modified peptide were subtracted from the monoisotopic peaks and the following isotopic peaks of NEM- $d_5$ - or  $^{13}C_6$ -IAA-modified peptide, respectively. The observed relative ratios of unlabeled and isotope-labeled peptides in the five sets of 9:1, 6:1, 3:1, 1:1, and 1:3 for 1D SDS-PAGE, or three sets of 3:1, 1:1, and 1:3 for 2D IEF/SDS-PAGE were plotted against their theoretical ratios as shown in the graphs in Section 2.

#### Acknowledgments

We thank Dr. Nobuaki Takemori and Mr. Hisao Haniu for kindly helping with the electrophoresis, and Mr. Anil Singh for providing excellent technical assistance throughout these studies. This work is supported by grants from the National Institute of Health (EY13877, EY12190, and RR17703). S.N. thanks Texas Tech University for start-up funds.

#### References and notes

- For example, (a) Matsumoto, H.; Kurien, B.; Takagi, Y.; Kahn, E. S.; Kinumi, T.; Komori, N.; Yamada, T.; Hayashi, F.; Isono, K.; Pak, W. L.; Jackson, K. W.; Tobin, S. L. *Neuron* **1994**, *12*, 997–1010; (b) Fach, E. M.; Garulacan, L. A.; Gao, J.; Xiao, Q.; Storm, S. M.; Dubaquié, Y. P.; Hefta, S. A.; Opitck, G. J. *Mol. Cell. Proteomics* **2004**, *12*, 120–1210; (c) Bernardini, G.; Renzone, G.; Comanducci, M.; Mini, R.; Arena, S.; D'Ambrosio, C.; Bambini, S.; Trabalzini, L.; Grandi, G.; Martelli, P.; Achtman, M.; Scaloni, A.; Ratti, G.; Santucci, A. *Proteomics* **2004**, *10*, 2893–2926; (d) Lamanda, A.; Zahn, A.; Roder, D.; Langen, H. *Proteomics* **2004**, *3*, 599–608.
- For example, (a) Gygi, S. P.; Rochon, Y.; Franza, B. R.; Aebersold, R. *Mol. Cell. Biol.* **1999**, *19*, 1720–1730; (b) Vuong, G. L.; Weiss, S. M.; Kammer, W.; Priemer, M.; Vingron, M.; Nordheim, A.; Cahill, M. A. *Electrophoresis* **2000**, *21*, 2594–2605.
- For example, (a) Goshe, M. B.; Smith, R. D. *Curr. Opin. Biotechnol.* **2003**, *14*, 101–109; (b) Tao, W. A.; Aebersold, R. *Curr. Opin. Biotechnol.* **2003**, *14*, 110–118; (c) Flory, M. R.; Griffin, T. J.; Martin, D.; Aebersold, R. *Trends Biotechnol.* **2002**, *20*, S23–S29; (d) Patterson, S. D. *Curr. Opin. Biotechnol.* **2000**, *11*, 413–418; (e) Griffin, T. J.; Goodlett, D. R.; Aebersold, R. *Curr. Opin. Biotechnol.* **2001**, *12*, 607–612; (f) Aebersold, R.; Goodlett, D. R. *Chem. Rev.* **2001**, *101*, 269–295; (g) Hamdan, M.; Righetti, P. G. *Mass Spectrom. Rev.* **2002**, *21*, 287–302; (h) Aebersold, R.; Mann, M. *Nature* **2003**, *422*, 198–207; (i) Julka, S.; Regnier, F. *J. Proteome Res.* **2004**, *3*, 350–363; (j) Broenstrup, M. *Expert Rev. Proteomics* **2004**, *1*, 503–512; (k) Righetti, P. G.; Campostrini, N.; Pascali, J.; Hamdan, M.; Astner, H. *Eur. J. Mass Spectrom.* **2004**, *10*, 335–348.
- For example, (a) Pasa-Tolic, L.; Jensen, P. K.; Anderson, G. A.; Lipton, M. S.; Peden, K. K.; Martinovic, S.; Tolic, N.; Bruce, J. E.; Smith, R. D. *J. Am. Chem. Soc.* **1999**, *121*, 7949–7950; (b) Oda, Y.; Huang, K.; Cross, F. R.; Cowburn, D.; Chait, B. T. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 6591–6596; (c) Veenstra, T. V.; Martinovic, S.; Anderson, G.; Pasa-Tolic, L.; Smith, R. *J. Am. Soc. Mass Spectrom.* **2000**, *11*, 78–82; (d) Chen, X.; Smith, L. M.; Bradbury, E. M. *Anal. Chem.* **2000**, *72*, 1134–1143; (e) Conrads, T. P.; Alving, K.; Veenstra, T. D.; Belov, M. E.; Anderson, G. A.; Anderson, D. J.; Lipton, M. S.; Pasa-Tolic, L.; Udseth, H. R.; Chrisler, W. B.; Thrall, B. D.; Smith, R. D. *Anal. Chem.* **2001**, *73*, 2132–2139; (f) Zhu, H.; Pan, S.; Gu, S.; Bradbury, E. M.; Chen, X. *Rapid Commun. Mass Spectrom.* **2002**, *16*, 2115–2123; (g) Berger, S. J.; Lee, S.-W.; Anderson, G. A.; Pasa-Tolic, L.; Tolic, N.; Shen, Y.; Zhao, R.; Smith, R. D. *Anal. Chem.* **2002**, *74*, 4994–5000; (h) Gu, S.; Pan, S.; Bradbury, E. M.; Chen, X. *J. Am. Soc. Mass Spectrom.* **2003**, *14*, 1–7; (i) Stump, M. J.; Jones, J. J.; Fleming, R. C.; Lay, J. O., Jr.; Wilkins, C. L. *J. Am. Soc. Mass Spectrom.* **2003**, *14*, 1306–1314; (j) Wu, C. C.; MacCoss, M. J.; Howell, K. E.; Matthews, D. E.; Yates, J. R., III *Anal. Chem.* **2004**, *76*, 4951–4959.
- Recently, Mann et al. reported a method called 'stable isotope labeling by amino acid in cell culture (SILAC),' in which isotope-labeled essential amino acids are quantitatively incorporated into cell lines. Ong, S.-E.; Blagoev, B.; Kratchmarova, I.; Kristensen, D. B.; Steen, H.; Pandey, A.; Mann, M. *Mol. Cell. Proteomics* **2002**, *1*, 376–386.
- (a) Gygi, S. P.; Rist, B.; Gerber, S. A.; Turecek, F.; Gelb, M. H.; Aebersold, R. *Nat. Biotechnol.* **1999**, *17*, 994–999; (b) Han, D. K.; Eng, J.; Zhou, H.; Aebersold, R. *Nat. Biotechnol.* **2001**, *19*, 946–951; (c) Griffin, T. J.; Aebersold, R. *J. Biol. Chem.* **2001**, *276*, 45497–45500; (d) Griffin, T. J.; Gygi, S. P.; Rist, B.; Aebersold, R.; Loboda, A.; Jilkine, A.; Ens, W.; Standing, K. G. *Anal. Chem.* **2001**, *73*, 978–986; (e) Griffin, T. J.; Han, D. K. M.; Gygi, S. P.; Rist, B.; Lee, H.; Aebersold, R.; Parker, K. C. *J. Am. Soc. Mass Spectrom.* **2001**, *12*, 1238–1246; (f) Gygi, S. P.; Rist, B.; Griffin, T. J.; Eng, J.; Aebersold, R. *J. Proteome Res.* **2002**, *1*, 47–54; (g) Ranish, J. A.; Yi, E. C.; Leslie, D. M.; Purvine, S. O.; Goodlett, D. R.; Eng, J.; Aebersold, R. *Nat. Genet.* **2003**, *33*, 349–355.
- For example, (a) Che, F. Y.; Fricker, L. D. *Anal. Chem.* **2002**, *74*, 3190–3198; (b) Ji, J.; Chakraborty, A.; Geng, M.; Zhang, X.; Amini, A.; Bina, M.; Regnier, F. *J. Chromatogr. B Biomed. Sci. Appl.* **2000**, *745*, 197–210; (c) Geng, M.; Ji, J.; Regnier, F. E. *J. Chromatogr. A* **2000**, *870*, 295–313; (d) Munchbach, M.; Quadroni, M.; Miotto, G.; James, P. *Anal. Chem.* **2000**, *72*, 4047–4057; (e) Turecek, F. *J. Mass Spectrom.* **2002**, *37*, 1–14; (f) Thompson, A.; Schäfer, J.; Kuhn, K.; Kienle, S.; Schwarz, J.; Schmidt, G.; Neumann, T.; Hamon, C. *Anal. Chem.* **2003**, *75*, 1895–1904; (g) Hsu, J.-L.; Huang, S.-Y.; Chow, N.-H.; Chen, S.-H. *Anal. Chem.* **2003**, *75*, 6843–6852; (h) Shi, Y.; Xiang, R.; Crawford, J. K.; Colangelo, C. M.; Horvath, C.; Wilkins, J. A. *J. Proteome Res.* **2004**, *3*, 104–111; (i) Ross, P. L.; Huang, Y. N.; Marchese, J. N.; Williamson, B.; Parker, K.; Hattan, S.; Khainovski, N.; Pillai, S.; Dey, S.; Daniel, S.; Purkayastha, S.; Juhasz, P.; Martin, S.; Bartlett-Jones, M.; He, F.; Jacobson, A.; Pappin, D. J. *Mol. Cell. Proteomics* **2004**, *3*, 1154–1169; (j) Kuyama, H.; Watanabe, M.; Toda, C.; Ando, E.; Tanaka, K.; Nishimura, O. *Rapid Commun. Mass Spectrom.* **2003**, *17*, 1642–1650; (k) Wang, S.; Zhang, X.; Regnier, F. E. *J. Chromatogr. A* **2002**, *949*, 153–162; (l) Weckwerth, W.; Willmitzer, L.; Fiehn, O. *Rapid Commun. Mass Spectrom.* **2000**, *14*, 1677–1681.
- For example, (a) Zhou, H.; Ranish, J. A.; Watts, J. D.; Aebersold, R. *Nat. Biotechnol.* **2002**, *20*, 512–515; (b) Goshe, M. B.; Conrads, T. P.; Panisko, E. A.; Angell, N. H.; Veenstra, T. D.; Smith, R. D. *Anal. Chem.* **2001**, *73*,

- 2578–2586; (c) Qiu, Y.; Sousa, E. A.; Hewick, R. M.; Wang, J. H. *Anal. Chem.* **2002**, *74*, 4969–4979.
9. Smolka, M.; Zhou, H.; Aebersold, R. *Mol. Cell. Proteomics* **2002**, *1*, 19–29.
10. Niwayama, S.; Kurono, S.; Matsumoto, H. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2257–2261.
11. Niwayama, S.; Kurono, S.; Matsumoto, H. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2913–2916.
12. Getz, E. B.; Xiao, M.; Chakrabarty, T.; Cooke, R.; Selvin, P. R. *Anal. Biochem.* **1999**, *273*, 73–80.
13. For example, (a) Hamdan, M.; Bordini, E.; Galvani, M.; Righetti, P. G. *Electrophoresis* **2001**, *22*, 1633–1644; (b) Brueggemeier, S. B.; Kron, S. J.; Palecek, S. P. *Anal. Biochem.* **2004**, *329*, 180–189; (c) Barber, D. S.; LoPachin, R. M. *Toxicol. Appl. Pharmacol.* **2004**, *201*, 120–136; (d) Mineki, R.; Taka, H.; Fujimura, T.; Kikkawa, M.; Shindo, N.; Murayama, K. *Proteomics* **2002**, *2*, 1672–1681.
14. Pasquarello, C.; Sanchez, J.-C.; Hochstrasser, D. F.; Corthals, G. F. *Rapid Commun. Mass Spectrom.* **2004**, *18*, 117–127.
15. In rare cases we found cysteine-containing proteins that do not react with IAA or  $^{13}\text{C}_6$ -IAA.
16. Herbert, B.; Galvani, M.; Hamdan, M.; Olivieri, E.; MacCarthy, J.; Pedersen, S.; Righetti, P. G. *Electrophoresis* **2001**, *22*, 2046–2057.
17. Sechi, S. *Rapid Commun. Mass Spectrom.* **2002**, *16*, 1416–1424.
18. Gehanne, S.; Cecconi, D.; Carboni, L.; Righetti, P. G.; Domenici, E.; Hamdan, M. *Rapid Commun. Mass Spectrom.* **2002**, *16*, 1692–1698.
19. Cahill, M. A.; Wozny, W.; Schwall, G.; Schroer, K.; Hölzer, K.; Poznanovic, S.; Hunzinger, C.; Vogt, J. A.; Stegmann, W.; Matthies, H.; Schrattenholz, A. *Rapid Commun. Mass Spectrom.* **2003**, *17*, 1283–1290.
20. Sebastiano, R.; Citterio, A.; Lapadula, M.; Righetti, P. G. *Rapid Commun. Mass Spectrom.* **2003**, *17*, 2380–2386.
21. While proteolytic  $\text{O}^{18}$  labeling using  $\text{H}_2\text{O}^{18}$  is reported as a simple quantitation method, this method creates only a 2–4 Da difference in the molecular weight. For example, (a) Mirgorodskaya, O. A.; Kozmin, Y. P.; Titov, M. I.; Körner, R.; Sönksen, C. P.; Roepstorff, P. *Rapid Commun. Mass Spectrom.* **2000**, *14*, 1226–1232; (b) Yao, X.; Freas, A.; Ramirez, J.; Demirev, P. A.; Fenselau, C. *Anal. Chem.* **2001**, *73*, 2836–2842; (c) Sakai, J.; Kojima, S.; Yanagi, K.; Kanaoka, M. *Proteomics* **2005**, *5*, 16–23; (d) Stewart, I. I.; Thomson, T.; Figeys, D. *Rapid Commun. Mass Spectrom.* **2001**, *15*, 2456–2465; (e) Hood, B. L.; Lucas, D. A.; Kim, G.; Chan, K. C.; Blonder, J.; Issaq, H. J.; Veenstra, T. D.; Conrads, T. P.; Pollet, I.; Karsan, A. *J. Am. Soc. Mass Spectrom.* **2005**, *16*, 1221–1230.
22. Other examples utilizing 2D electrophoresis and metabolic labeling, see (a) Turko, I.; Murad, F. *J. Biol. Chem.* **2003**, *278*, 35844–35849; (b) Vogt, J. A.; Schroer, K.; Hölzer, K.; Hunzinger, C.; Klemm, M.; Biefang-Arndt, K.; Schillo, S.; Cahill, M. A.; Schrattenholz, A.; Matthies, H.; Stegmann, W. *Rapid Commun. Mass Spectrom.* **2003**, *17*, 1273–1282.
23. Wu, W. W.; Wang, G.; Baek, S. J.; Shen, R.-F. *J. Proteome Res.* **2006**, *5*, 651–658.
24. Klammer, A. A.; MacCoss, M. J. *J. Proteome Res.* **2006**, *5*, 695–700.