



β-Peptides with improved affinity for hDM2 and hDMX

Elizabeth A. Harker^a, Douglas S. Daniels^{a,c}, Danielle A. Guarracino^{a,d}, Alanna Schepartz^{a,b,*}

^a Department of Chemistry, Yale University, New Haven, CT 06520, United States

^b Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520, United States

^c Novartis Institutes for BioMedical Research, Inc, Cambridge, MA 02139, United States

^d Department of Chemistry, New York University, New York, NY 10003, United States

ARTICLE INFO

Article history:

Received 20 November 2008

Revised 7 January 2009

Accepted 8 January 2009

Available online 23 January 2009

Keywords:

Protein–protein interactions

β-Peptides

Peptidomimetics

Inhibitors

Foldamer

p53

hDM2

Cancer

ABSTRACT

We previously described a series of 3_{14} -helical β-peptides that bind the hDM2 protein and inhibit its interaction with a p53-derived peptide in vitro. Here we present a detailed characterization of the interaction of these peptides with hDM2 and report two new β-peptides in which non-natural side chains have been substituted into the hDM2-recognition epitope. These peptides feature both improved affinity and inhibitory potency in fluorescence polarization and ELISA assays. Additionally, one of the new β-peptides also binds the hDM2-related protein, hDMX, which has been identified as another key therapeutic target for activation of the p53 pathway in tumors.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Short β³-peptides can adopt a helical secondary structure in water when stabilized by salt-bridges and/or macrodipole effects.^{1–8} This secondary structure, the 3_{14} -helix, is characterized by a periodicity of 3.1-residues/turn that arranges the β³-peptide side chains at 120° intervals when the structure is viewed down the helical axis. The periodicity of the 3_{14} -helix results in a reasonable, but imperfect, superposition of side chains along one face of the β-peptide helix (at positions i , $i + 3$, and $i + 6$) with those at positions i , $i + 4$, and $i + 7$ of an α-helix; this superposition can be improved by design.^{9–11} Indeed, over the past decade, we and others have reported a number of β-peptide 3_{14} -helices that bind diverse protein targets including somatostatin receptors, hDM2, and viral fusion proteins.^{11–14} Longer molecules containing both β- and α-amino acids have been shown to bind Bcl-x_L.¹⁵ In addition, oligomers of β³-amino acids convey the advantage of near complete resistance to proteolysis.^{1,16}

One target for β³-peptides that has received significant attention is hDM2, largely because of its recognized potential for novel oncology treatments.^{17,18} hDM2 is overexpressed in about 7% of human tumors, most commonly osteosarcomas and soft tissue sarcomas, and impairs the p53 stress response to allow unchecked cell

growth.^{17–20} The structure of hDM2 bound to a short α-helical fragment of p53 has provided a virtual blueprint for inhibitor design, highlighting the importance of three p53 side chains (F19, W23, and L26) that insert into a hydrophobic cleft of hDM2.²¹ A number of peptidic, peptidomimetic, and small molecule p53-mimetics have been described that inhibit this interaction in vitro^{17,22–33} and, in some cases, in vivo.^{24,27,34} In 2005, we reported a series of macrodipole- and salt bridge-stabilized 3_{14} -helical β³-peptides that each display the three critical p53 side chains along one face of the 3_{14} -helix; one of these molecules binds hDM2 with nanomolar affinity and inhibits its interaction with a p53-derived peptide.^{11,13} Importantly, high-resolution NMR analysis of these β-peptides indicates that their 3_{14} -helix is distorted—slightly unwound at the C-terminus—and that this distortion results in a significantly improved structural mimic of the p53 α-helix.^{9,11} These observations emphasize the versatility of β-peptides dominated by β³-amino acids when compared with more conformationally constrained analogs such as AHC.³⁵

This success encouraged us to evaluate the affinity of these β³-peptides for the hDM2-related protein, hDMX.^{36–38} hDMX is similar to hDM2 in that its N-terminal region binds to p53 and blocks transactivation;³⁹ however, the expression of hDMX is not regulated by p53, nor has it been shown to target p53 for degradation, so the overall mechanism of regulation is likely to differ from that of hDM2. β³-peptides capable of inhibiting both hDMX and hDM2 might offer a unique advantage with respect to p53 activation,

* Corresponding author. Tel.: +1 203 432 8276; fax: +1 203 432 3486.

E-mail address: alanna.schepartz@yale.edu (A. Schepartz).

whereas those that selectively target one of these two proteins might have use as novel research tools. Here we build upon our initial results with β^3 -peptide ligands for hDM2 to characterize further the binding interaction of these peptides with hDM2 and report two β^3 -peptides with improved affinity and inhibitory potency. Additional binding data indicates that one of these peptides also possesses sub-micromolar affinity for hDMX.

2. Results

2.1. Design of third-generation β^3 -peptide ligands for hDM2: Non-natural side chains

One of the most notable features of the previously reported X-ray structure of mDM2 (the murine homolog of hDM2) bound to a p53-derived peptide (p53AD) is the inability of the p53AD Trp₂₃ side chain to fill its binding pocket on the hDM2 surface.²¹ Indeed, replacement of Trp₂₃ with 6-chlorotryptophan in the context of an α -peptide ligand reported more than a decade ago resulted in a 60-fold increase in affinity for hDM2.²³ More modest (2–4-fold) improvements were observed when 6-chlorotryptophan was introduced into the equivalent position of either peptoid or β -hairpin ligands.^{29,30} Similarly, a trifluoromethylphenyl substituent is found in several high affinity small molecule ligands for hDM2 (IC₅₀ values ranging from 0.98 to >125 μ M for a series of 1,4-benzodiazepine-2,5-diones).^{40–42} Based upon these results, we synthesized two new variants of β^3 -8: one containing a 3-trifluoromethylphenyl side chain in place of Trp₂₃, (β^3 -12) and one containing 6-chlorotryptophan (β^3 -13) (Fig. 1).

2.2. Direct and competitive fluorescence polarization analysis of hDM2- β^3 -peptide interactions

We first used a direct fluorescence polarization assay to compare the hDM2 affinities of β^3 -12FIU and β^3 -13FIU to that of β^3 -8FIU and assess whether the non-natural side chains improved the affinity of the β^3 -peptide for hDM2. For this assay, β^3 -peptides labeled on the N terminus with fluorescein-5-EX, succinimidyl ester (Invitrogen) were equilibrated with hDM2_{1–188} at concentrations between 1 nM and 15 μ M. A fluorescently labeled α -peptide corresponding




Figure 1. Helical net representations of β^3 -1, β^3 -8, β^3 -12, and β^3 -13. β^3 -Homomino acids are identified by the one-letter code corresponding to the analogous α -amino acid.




Figure 2. Fluorescence polarization analysis of β^3 -peptide-hDM2 equilibria. (A) Direct fluorescence polarization assay: plot of the polarization of the indicated fluorescently labeled β^3 -peptides and p53AD_{15–31} as a function of [hDM2_{1–188}]. (B) Competition fluorescence polarization assay: plot of the polarization of the p53AD_{15–31} complex with hDM2_{1–188} as a function of the concentration of unlabeled peptide shown. Each value shown represents the average of at least four independent determinations, each performed in triplicate and averaged; error bars represent the standard error.

to residues 15–31 of p53 (p53AD_{15–31}) was used as a positive control.¹³ The K_d of the p53AD_{15–31}·hDM2 complex determined using this direct binding assay was 72.5 ± 9.12 nM, in agreement with previous work (Fig. 2A).^{13,17,21,43–46} The K_d values of the hDM2 complexes with β^3 -12^{FIU} and β^3 -13^{FIU} were 28.2 ± 4.79 and 30.1 ± 4.93 nM, respectively, representing a 2-fold improvement in affinity over p53AD_{15–31}. Moreover, the K_d values measured for β^3 -12^{FIU} and β^3 -13^{FIU} represent a 10-fold improvement over previously reported β^3 -peptide ligands for hDM2 ($K_d = 292 \pm 22.8$ nM for β^3 -1^{FIU} and 204 ± 17.7 nM for β^3 -8^{FIU}).^{11,13}

The relative affinities of β^3 -12 and β^3 -13 for hDM2 were also measured using a competitive fluorescence polarization assay in which unlabeled p53AD_{15–31} or β^3 -peptide competed with p53AD_{15–31} for binding to hDM2. In this assay, unlabeled p53AD_{15–31} inhibited p53AD_{15–31}·hDM2 complexation with an IC₅₀ of 5.22 ± 0.196 μ M, in line with previous reports.^{11,13,43} Under identical conditions β^3 -12 and β^3 -13 inhibited p53AD_{15–31}·hDM2 complexation with IC₅₀ values of 15.9 ± 2.72 μ M and 3.30 ± 0.681 μ M, respectively (Fig. 2B). The inhibitory activity of β^3 -13 is several-fold greater than that of previously reported β -peptides β^3 -1 and β^3 -8 (IC₅₀ = 39.2 ± 8.52 μ M and 17.8 ± 0.530 μ M, respectively).¹¹ By contrast, the IC₅₀ value for β^3 -12 is somewhat higher than expected, likely due to its poor solubility in aqueous buffer at concentrations greater than 100 μ M. While the data from both direct and competition fluorescence polarization experiments indicate that β^3 -12 and β^3 -13 possess high affinity for hDM2, the competition data is particularly valuable as it directly measures the affinity of a ligand for the p53 binding pocket on hDM2.

2.3. Exploring the binding mode of β^3 -peptide ligands for hDM2

Previous work in our lab has indicated that versions of β^3 -1 labeled with fluorescein at the N- or C-termini possess similar affinities for hDM2 as determined by direct FP analysis.¹³ Nevertheless, to rule out the possibility that the appended fluorescein moiety contributes substantially to binding affinity, we prepared a second set of β^3 -peptides labeled with Alexa Fluor[®] 594 carboxylic acid, succinimidyl ester (Invitrogen). Both the structure and charge of Alexa Fluor[®] 594 differ from that of fluorescein (Fig. 3)—Alexa Fluor[®] is larger than fluorescein and exists as a zwitterion under the conditions of the binding assays (pH \sim 7.4) while fluorescein exists as a dianion. In addition, the fluorescein label contains a hydrophilic linker whereas the Alexa Fluor[®] label does not. With these differences in mind, we synthesized two additional β^3 -peptides labeled with Alexa Fluor[®] 594 on their N-termini. One (β^3 -8^{AI}) is extended at the N-terminus by two additional β^3 -homoglycine residues, whereas the other (β^3 -12^{AI}) is not. The




Figure 3. (A) Structures of $\beta 53-8^{flu}$ and $\beta 53-8^{AI}$ ($n = 2$). $\beta 53-12^{flu}$ and $\beta 53-12^{AI}$ ($n = 0$) contain a β^3 -*m*-trifluoromethylphenylalanine residue in place of β^3W . (B) Plot illustrating the polarization of the indicated β^3 -peptide as a function of [hDM2]₁₋₁₈₈.

affinities of these two β^3 -peptides for hDM2 were evaluated by direct fluorescence polarization analysis as described above. The K_d values measured for $\beta 53-8^{AI}$ and $\beta 53-12^{AI}$ differed by only about 2-fold from the values measured for the respective fluorescein-labeled molecule despite the differences in linker and label (Fig. 3, B and C). These results suggest that the fluorophore does not contribute more than minimally to the stability of β^3 -peptide-hDM2 complexes and that the labeled β^3 -peptides provide an accurate report of hDM2 affinity.

2.4. ELISA competition

Next we employed a validated competitive ELISA to further support a functional interaction between hDM2 and $\beta 53-1$, **8**, **12**, and **13**.^{28,47} Unlabeled α - and β^3 -peptides at concentrations between 0.012 and 200 μ M were first equilibrated with 1 μ M hDM2₁₋₁₈₈, and then added to streptavidin-coated microtiter plates coated with $p53AD_{15-31}$ carrying a biotin label at the N-terminus ($p53AD_{15-31}^{bio}$). The ability of unlabeled α - and β^3 -peptides to inhibit the interaction between hDM2₁₋₁₈₈ and immobilized $p53AD_{15-31}^{bio}$ was measured by detecting bound hDM2 with an hDM2-specific primary antibody (N-20, Santa Cruz Biotechnology), a horseradish peroxidase-conjugated secondary antibody (goat anti-rabbit IgG-HRP, Santa Cruz Biotechnology) and QuantaBlu™ Fluorogenic Peroxidase Substrate (Pierce) (Fig. 4A). As expected, $p53AD_{15-31}$ itself as well as $\beta 53-1$, **8**, **12**, and **13** reduced the amount of hDM2 captured by immobilized $p53AD_{15-31}^{bio}$ in a dose-dependent manner,




Figure 4. ELISA analysis of β^3 -peptide-hDM2 affinity. (A) Plot of the fluorescence intensity of QuantaBlu™ Fluorogenic Peroxidase Substrate (Pierce) as a function of the concentration of the indicated inhibitor. Each value represents the average of at least two independent experiments of three separate trials; error bars represent the standard error. (B) Helical net representation of βNEG . β^3 -Homoamino acids are identified by the one-letter code corresponding to the analogous α -amino acid.

indicating effective inhibition of $p53$ -hDM2 complexation. A related β^3 -peptide containing β^3 -homoalanine residues in place of the β^3 -homophenylalanine, β^3 -homotryptophan, and β^3 -homoleucine residues in $\beta 53-8$ (βNEG , Fig. 4B) did not inhibit the fraction of hDM2 captured at concentrations as high as 200 μ M.¹¹ The relative IC_{50} values determined for $\beta 53-1$, **8**, **12**, and **13** and for $p53AD_{15-31}$ in the ELISA were similar to those determined using the fluorescence polarization competition assay, but the absolute values were generally 2–3-fold lower in the ELISA (Table 1). Despite these small systematic differences, the ELISA results provide further evidence for the inhibitory potency of the β^3 -peptides studied herein and support the conclusion that the fluorophore used in the fluorescence polarization assays does not play an appreciable role in the observed affinity.

2.5. Cellular entry of β^3 -peptides

Recent work has suggested that certain peptides, even those lacking appreciable positive charge, can enter the cytosol of living mammalian cells.⁴⁸ With this in mind, we investigated whether $\beta 53-12$ might show even modest uptake by HCT116 cells. $\beta 53-12$ was chosen because of its relative ease of synthesis compared to $\beta 53-13$, which is complicated by optical resolution of 6-chloro-tryptophan. $\beta 53-12^{flu}$ and fluorescein-labeled control peptides (at concentrations ranging from 5 to 20 μ M) were incubated with HCT116 cells for either 1 h or 4 h, and the resulting mean cellular fluorescence was quantified by flow cytometry (Fig. 5A and B). The $p53AD^{flu}$ peptide served as a negative control and showed only slight uptake at both time points studied. The positive controls, $(PRR)_3^{flu}$ and Tat^{flu} were taken up by the cells at both time points, but the signal for both decreased over time.^{49,50} Though the signal for $\beta 53-12^{flu}$ was lower than that observed for arginine-rich $(PRR)_3^{flu}$ and Tat^{flu} , $\beta 53-12^{flu}$ showed clearly increased uptake relative to $p53AD^{flu}$. Additionally, the fluorescence signal for $\beta 53-12^{flu}$ increased over time, highlighting its resistance to cellular degradation. Although a trypsin wash was included in the flow cytometry protocol in an effort to digest membrane proteins that could bind the arginine-rich peptides,⁵¹ live-cell confocal microscopy was used to confirm the internalization and characterize the intracellular distribution of $\beta 53-12^{flu}$. As shown in Figure 5C, addition of $\beta 53-12^{flu}$ to HCT116 cells resulted in a punctate staining pattern that colocalized with the endocytotic marker dextran, indicating internalization via endosomes. Western blot experiments reveal that $\beta 53-12^{flu}$ modestly upregulates p53 activity in HCT116 and JAR cells but not HCT116 p53-null cells (E. Harker, unpublished results), suggesting that endosomal leakage occurs. Current work is focused on the application of Kodadek's cell per-




Figure 5. Entry of $\beta 53$ -12^{flu} and controls quantified by flow cytometry (A and B) and observed by confocal microscopy (C). Flow cytometry: mean cellular fluorescence was calculated from the histogram of fluorescence intensity and was corrected for background cellular fluorescence by subtracting the geometric mean of cells treated with only PBS. Each value represents the average of three independent trials. Error bars represent the standard error. (A) HCT116 cell uptake of fluorescein-labeled peptides after 1 h. (B) Uptake after 4 h. Confocal microscopy: HCT116 cells were incubated with 20 μM fluorescein-labeled peptide (green) for 2 h in McCoy's 5A medium supplemented with 10% FBS. Endosomes were visualized using 10 μM 10 kDa dextran labeled with AlexaFluor 647 (red). Panel 1 shows $\beta 53$ -12^{flu} only, panel 2 dextran only, panel 3 bright-field cell image only, panel 4 superposition of previous 3 panels.

Table 1
Comparison of IC₅₀ values for β^3 -peptides and p53AD binding hDM2 as determined by competitive fluorescence polarization and ELISA.

Ligand	IC ₅₀ (FP) (μM)	IC ₅₀ (ELISA) (μM)
p53AD	5.22 ± 0.196	2.29 ± 0.265
β53-1	39.2 ± 8.52	13.5 ± 3.20
β53-8	17.8 ± 0.530	7.17 ± 2.52
β53-12	15.9 ± 2.72	6.32 ± 0.316
β53-13	3.30 ± 0.681	1.60 ± 0.0360

meability screen^{48,52} to identify variants of $\beta 53$ -12 that achieve higher levels of cytosolic localization.

2.6. Affinity of β^3 -peptides for hDMX

As described earlier, the p53-binding surfaces on hDM2 and the related protein hDMX are quite similar, with about 80% identity between the two.⁵³ Despite this close relationship, and the fact that α -peptides that compete with p53 to bind hDM2 also bind hDMX,^{54–56} a number of small molecule ligands for hDM2, including Nutlin-3, do not bind hDMX.^{53,57} Structural work has indeed shown that the hDMX binding pocket for p53 is somewhat obstructed compared to the hDM2 binding pocket.^{58,59} We therefore tested our existing β^3 -peptide ligands to see if their extended binding surface, similar to that of an α -helix, would render them suitable for inhibition of the p53-hDMX interaction.

We used a direct fluorescence polarization assay to measure the affinities of $\beta 53$ -1^{flu}, $\beta 53$ -8^{flu}, $\beta 53$ -12^{flu}, and $\beta 53$ -13^{flu} for hDMX_{1–200}. As a positive control, we used the α -peptide p53 12/1^{flu} which possesses higher affinity for hDMX ($K_d = 237 \pm 21.5$ nM) than p53AD^{flu} ($K_d = 841 \pm 28.8$ nM) (Fig. 6A). Interestingly, of all β^3 -peptides tested, only $\beta 53$ -12^{flu} bound hDMX with submicromolar affinity ($K_d = 518 \pm 41.3$ nM). $\beta 53$ -13^{flu}, which was comparable to $\beta 53$ -12^{flu} in affinity for hDM2, bound more weakly to hDMX ($K_d = 1600 \pm 79.9$ nM). $\beta 53$ -8^{flu} and $\beta 53$ -1^{flu} also displayed low affinity for hDMX ($K_d = 2130 \pm 139$ and 9090 ± 9937 nM, respectively).

Next we used a competitive fluorescence polarization assay to determine whether the β^3 -peptides competed with p53 12/1^{flu} for binding to hDMX. Unlabeled p53 12/1 inhibited the binding of p53 12/1^{flu} and hDMX with an IC₅₀ of 3.87 ± 0.389 μM (Fig. 6B). Similarly, $\beta 53$ -12 inhibited p53 12/1^{flu}-hDMX complexation with an IC₅₀ of




Figure 6. Fluorescence polarization analysis of β^3 -peptide:hDMX equilibria. (A) Direct fluorescence polarization assay: plot of the polarization of the indicated fluorescently labeled β^3 -peptides, p53 12/1^{flu}, and p53AD^{flu} as a function of [hDMX_{1–200}]. (B) Competition fluorescence polarization assay: plot of the polarization of the p53 12/1^{flu} complex with hDMX_{1–200} as a function of the concentration of unlabeled peptide shown. Each value shown represents the average of at least four independent determinations, each performed in triplicate and averaged; error bars represent the standard error.

10.9 ± 1.13 μM; the IC₅₀ values for $\beta 53$ -13 and $\beta 53$ -8 were 12.0 ± 0.748 and 20.1 ± 2.51 μM, respectively, whereas $\beta 53$ -1 was too weak as an inhibitor to be quantified. Overall, the data indicates that β^3 -peptides designed to bind hDM2 also efficiently bind the hDMX protein, albeit with some variations in affinity.

3. Discussion

The development of general chemical strategies to inhibit transient interactions between proteins—both inside and outside the cell—represents one of the paramount goals of modern chemistry. Historically, it has been challenging to develop drug-like inhibitors of proteins whose natural ligands are other proteins, highlighting the need for novel approaches.^{60,61} In this work we describe β^3 -peptides containing non-natural side chains that possess significantly improved affinity for hDM2, a protein widely recognized for its interaction with the tumor suppressor p53.^{17,18} Introduction of either a 3-trifluoromethylphenyl or a 6-chlorotryptophan side chain in place of the central tryptophan of the hDM2-recognition epitope resulted in molecules that bind hDM2 with K_d values of approximately 30 nM, an improvement of nearly 10-fold over previously reported β^3 -peptide ligands. Furthermore, we completed a detailed study of the binding mode of these peptides using a series

of fluorescence and ELISA assays to confirm their inhibitory activity towards the p53-hDM2 interaction. The results provide added evidence that these β^3 -peptides interact with hDM2 in much the same manner as **p53AD**. Based on previous reports, it is likely that the observed gain in affinity with the 3-trifluoromethylphenyl and 6-chlorotryptophan side chains is due to the ability of these groups to better fill the hydrophobic pocket on the hDM2 surface compared to the tryptophan side chain of the parent peptide.^{21,23,29,30} It is also interesting to note that both the trifluoromethyl and chloro groups are electron withdrawing; therefore it is possible that this electrostatic effect leads to a favorable aromatic donor–acceptor interaction with a side chain on the hDM2 surface. Structural information on the β^3 -peptide-hDM2 interaction is still needed to further explore these possibilities and would provide useful information for future designs.

Recent studies of p53 activation as a therapeutic strategy for cancer have pointed to the importance of inhibiting the interaction between p53 and the hDM2-related protein hDMX. As mentioned above, these two proteins have both been implicated in the regulation of p53, but likely function through different mechanisms.^{36,37,62} Both proteins bind to the p53 activation domain through their N-terminal regions; in fact, 10 of the 13 residues of hDM2 determined to contribute to its interaction with p53 are conserved in the hDMX protein.³⁹ Previous research has shown that optimized α -peptides designed to bind hDM2 also inhibit the p53-hDMX interaction with IC₅₀ values of 1.65 μ M (**12/1**: MRPFDYWEGLN) and 0.10 μ M (**pDI**: LTFEHYWAQLTS) reported for the most potent.^{54,55} By contrast, similar experiments have shown that potent small molecule inhibitors of the p53-hDM2 interaction (e.g., Nutlin-3, IC₅₀ of 0.09 μ M) do not inhibit p53-hDMX at concentrations as high as 30 μ M. These results indicate that although the p53-binding surfaces on hDM2 and hDMX are similar enough to allow for targeting by longer peptides, they are not readily bound by the same small molecule ligand, findings that have been recently substantiated by structural data reported for the p53-hDMX interaction.^{58,59} Interestingly, it has been demonstrated that high hDMX levels or an inability to degrade hDMX leads to Nutlin-resistance in certain cell lines.^{53,57} As a result, there exists a subset of tumors that is not responsive to treatment with Nutlin-3. We hoped that our β^3 -peptides designed to bind hDM2 would also bind hDMX based upon their structural similarity to the α -helical structure already shown to target hDMX efficiently. Excitingly, one of these β^3 -peptides (**p53-12**) does in fact bind hDMX with submicromolar affinity and effectively inhibits its interaction with a p53-derived peptide. Interestingly, we saw some variation in affinity between hDM2 and hDMX for the series of β^3 -peptides studied, which can likely be traced to the observed differences in the p53-binding pockets of each protein. Further optimization of β^3 -peptide affinity for hDMX is a clear progression from this promising starting point given the limited number of molecules available to bind this new oncology target and could provide an advantage for the application of the β^3 -peptide scaffold compared to a small molecule.

Ultimately, we are interested in determining if these peptides are able to disrupt the p53-hDM2 and/or the p53-hDMX interaction in cells. β^3 -peptides would provide a useful complement to existing molecules aimed at p53 activation since they are more proteolytically stable than α -peptides and cover more surface area than small molecules (this latter feature may be particularly advantageous in the case of hDMX). The flow cytometry and confocal microscopy data presented here clearly demonstrate a modest degree of uptake for **p53-12^{flu}**; however, compared to the positive control peptides (**PRR^{flu}**) and **Tat^{flu}** the cellular entry of **p53-12^{flu}** stands to be significantly improved upon. At this point, we are pursuing strategies of arginine incorporation into the β^3 -peptide 3₁₄-helix based upon observations that the positively charged guanid-

inium group of arginine can promote cellular entry.⁶³ This builds upon earlier work in our lab in which we incorporated arginine residues into a well-folded miniature protein to improve its cellular permeability.^{49,64} Additionally, this strategy has been implemented previously in the Gellman and Seebach labs;^{65,66} however, there has not yet been a comprehensive analysis of the effects of arginine incorporation on a functional 3₁₄-helical β^3 -peptide.

4. Conclusion

We have significantly improved upon the affinity of our first- and second-generation β^3 -peptide ligands for hDM2 through the incorporation of non-natural side chains into the 3₁₄-helical scaffold. The use of a 3-trifluoromethylphenyl or a 6-chlorotryptophan side chain in place of the central tryptophan residue of the hDM2-recognition epitope led to molecules with K_d values of 30 nM for binding hDM2, a 10-fold improvement over our previously reported β^3 -peptide ligands. A detailed analysis of the binding mode of these peptides indicates that they effectively compete with a p53-derived peptide for its binding site on the hDM2 protein. Additionally, we have demonstrated that β^3 -peptides effectively target the hDM2-related protein, hDMX, a target of growing importance in oncology. Current work is focused on improving the cellular permeability of these β^3 -peptides and on further optimization of their binding affinity, particularly in the case of hDMX for which only a limited number of potent ligands have been reported.

5. Experimental

5.1. General

Fmoc-protected α -amino acids, PYBOP®, HOBt, and Wang resin were purchased from Novabiochem (San Diego, CA). Dimethylformamide (DMF), trifluoroacetic acid (TFA), and piperidine were purchased from American Bioanalytical (Natick, MA). All other reagents were purchased from Sigma–Aldrich. Fmoc-(S)-3-amino-4-(3-trifluoromethylphenyl)butyric acid was purchased from AnaSpec, Inc. (San Jose, CA). The remaining Fmoc- β^3 -(L)-amino acids were synthesized from enantiomerically pure α -amino acids via the Arndt-Eistert procedure¹ with the exception of Fmoc-(S)-3-amino-4-(6-chloroindole)butyric acid, the enantiomeric resolution and homologation of which is reported here.

5.2. Enantiomeric resolution of 6-chlorotryptophan

D,L-6-Chlorotryptophan was resolved by esterification and selective hydrolysis in the presence of α -chymotrypsin, a procedure adapted from that for tryptophan and α -methyltryptophan.^{67,68} Thionyl chloride (1.6 mL, 22.0 mmol) was added dropwise to D,L-6-chlorotryptophan (Biosynth AG, Switzerland, 1.048 g, 4.39 mmol) in 100 mL of chilled, anhydrous methanol with stirring. The mixture was warmed to room temperature over 1 h and then brought to reflux at 65 °C. The reaction was monitored by thin layer chromatography (85:15 acetonitrile/water), and, after 19 h, cooled and solvent removed under reduced pressure. D,L-6-chlorotryptophan methyl ester was recrystallized in four crops from methanol with diethyl ether to yield, after lyophilization, 1.22 g (96%) of the hydrochloride salt. ¹H NMR (400 MHz, CD₃OD): δ 7.49 (d, J = 8.5 Hz, 1H), 7.40 (s, 1H), 7.21 (s, 1H), 7.06 (d, J = 8.5 Hz, 1H), 4.32 (t, J = 6.5 Hz, 1H), 3.79 (s, 3H), 3.40 (d, J = 5.5 Hz, 1H). Calcd Mass 252.697, obsd 253.09.

D,L-6-Chlorotryptophan methyl ester hydrochloride (1.0691 g, 3.7 mmol) was dissolved in 11 mL DMSO and diluted with

320 mL water. Sodium citrate, pH 5.5 (22.2 mL, 500 mM) was added and hydrolysis initiated by the addition of α -chymotrypsin (97.9 mg). The reaction was monitored by HPLC, and after 22 h, stopped by the addition of 150 mL of ethanol. The mixture was concentrated under reduced pressure, extracted three times with ethyl acetate and back-extracted with water. The combined aqueous layers were concentrated under reduced pressure, diluted 2-fold with acetonitrile and purified by flash chromatography (85:15 acetonitrile/water). The recovered solid was washed sparingly with cold water to yield 213 mg of L-6-chlorotryptophan, and optical purity assessed by *N*- α -(2,4-dinitro-5-fluorophenyl)-L-valine amide derivatization⁶⁹ (63% yield, 94% ee). ¹H NMR (400 MHz, CD₃OD): δ 7.66 (d, *J* = 8.5 Hz, 1H), 7.36 (s, 1H), 7.21 (s, 1H), 7.02 (d, *J* = 8.5 Hz, 1H), 3.83 (t, *J* = 4.5 Hz, 1H), 3.46 (d, *J* = 15.3 Hz, 1H), 3.17 (d, *J* = 9.2 Hz, 1H). Calcd Mass 238.670, obsd 239.00.

5.3. Synthesis of Fmoc-(S)-3-amino-4-(6-chloroindole)butyric acid

The base labile 9-fluorenylmethyloxycarbonyl (Fmoc-) amino protective group was coupled to the free N terminus of L-6-chlorotryptophan according to previously published methods.^{70,71} The amino acid (25 mmol) was dissolved in water (25 mL) in the presence of 1 equiv of triethylamine. Simultaneously, the Fmoc succinimidyl ester (24 mmol) was dissolved in acetonitrile (25 mL) and gently heated to promote dissolution. The ester solution was added to the amino acid solution in one portion and allowed to stir at room temperature for 30 min. The pH of the solution was maintained in the 8.5–9.0 range by dropwise addition of triethylamine as the pH of the reaction mixture drops sharply initially due to liberation of free *N*-hydroxysuccinimate. When complete, the solution was concentrated via rotary evaporation and the resulting residue was poured over a solution of hydrochloric acid (1.5 N). A spontaneously crystallized product was separated, washed and set aside. The oil precipitate was isolated by two ethyl acetate extractions followed by a wash with brine. The resulting solution was dried over sodium sulfate and rotary evaporated to dryness. The product was then brought up in acetonitrile and water and dried over vacuum. The yield obtained was 87%. ¹H NMR (400 MHz, CDCl₃): δ 7.78 (d, 2H), 7.30–7.52 (8H), 7.06 (d, 1H), 6.93 (s, 1H), 5.26 (d, 1H), 4.73 (s, 1H), 4.43 (t, 2H), 4.19 (d, 1H), 3.32 (s, 2H), 2.83 (s, 1H). Calcd Mass 461.00, obsd 461.30 and 483.30 (+Na).

The corresponding Fmoc-protected β -amino acid was synthesized using the Arndt-Eistert procedure to yield 45 mg (22%) of Fmoc-(S)-3-amino-4-(6-chloroindole)butyric acid.¹ ¹H NMR (400 MHz, CD₃OD): δ 9.29 (d, *J* = 7.5 Hz, 2H), 9.09 (t, *J* = 6.6 Hz, 3H), 8.89 (t, *J* = 7.3 Hz, 3H), 8.79 (dd, *J* = 8.6 Hz, 2H), 8.58 (s, 1H), 8.47 (d, *J* = 8.4 Hz, 1H), 5.77 (m, *J* = 7.2 Hz, 3H), 5.65 (t, *J* = 6.8 Hz, 1H), 4.46 (dd, *J* = 3.1 Hz, 2H), 4.02 (t, *J* = 6.6 Hz, 2H), 2.79 (s, 1H), 2.41 (s, 1H). Calcd Mass 474.7, obsd 497.4 (product + Na+).

5.4. β^3 -Peptide synthesis, general procedure

β^3 -Peptides were synthesized manually in a glass peptide-synthesis vessel with fritted glass at the top and bottom and a sidearm for addition of reagents. Peptides were synthesized on a 25 μ mol scale using standard Fmoc chemistry with Wang resin and Fmoc-protected β^3 -amino acid monomers.

5.4.1. β^3 -Peptide synthesis, Wang resin loading

Wang resin (1.0 mmol-OH/g, 380 mg, 0.38 mmol) was swelled in DMF for 30 min in a glass peptide-synthesis vessel. Excess DMF was removed by filtration. Fmoc- β^3 -hGlu(tBu) (500 mg, 1.14 mmol) was added to 10 mL dry CH₂Cl₂ at 0 °C and stirred

under argon. 1,3-Dicyclohexylcarbodiimide (78.4 mg, 0.38 mmol) was dissolved in 3 mL DMF and added to the solution. After 30 min, the solution was warmed to room temperature and concentrated to a white slurry by rotary evaporation. The slurry was dissolved in 5 mL DMF and added directly to the resin. Dimethylaminopyridine (46.5 mg, 0.38 mmol) was added and the mixture was shaken vigorously on a moving tray for 3 h. The reaction mixture was drained and washed with DMF (2 \times 30 s). Unreacted amino groups were acetylated by treatment with 6% v/v acetic anhydride and 6% v/v NMM in DMF (30 min). The capped resin was washed with DMF (8 \times 30 s), CH₂Cl₂ (8 \times 30 s), and MeOH (4 \times 30 s) and dried for 30 min under N₂. The extent of resin loading was determined as described.⁸

5.4.2. β^3 -Peptide synthesis, microwave procedure

The protocol for the synthesis of β^3 -peptides was adapted from reports by Murray and Gellman.⁷² Fmoc- β^3 -hGlu(tBu)-loaded Wang resin was placed in a glass peptide-synthesis vessel and swelled with DMF for 20 min. A solution of 20% piperidine/DMF was added to the vessel along with a stir bar for agitation, and the vessel was placed in the microwave reactor (CEM MARS) and irradiated (200 W maximum power, 70 °C, ramp 2 min, hold 4 min, cool 5 min). The vessel was removed from the microwave reactor, and the deprotected resin washed with DMF (6 \times 30 s). The appropriate β^3 -homoamino acid (3 equiv), PyBOP® (3 equiv), HOBt (3 equiv), and diisopropylethylamine (DIEA, 8 equiv) were dissolved in 1.5 mL DMF and added to the resin. The vessel was again placed in the microwave reactor and irradiated (200 W maximum power, 60 °C, ramp 2 min, hold 6 min, cool 5 min). Upon removal from the microwave reactor, the resin was washed with DMF (6 \times 30 s), and the above steps were repeated until the β^3 -peptide sequence was complete. Following removal of the final Fmoc protecting group, the resin was washed with DMF (8 \times 30 s) and CH₂Cl₂ (8 \times 30 s), dried for 20 min under N₂, and treated for 90 min with a cleavage cocktail of 3% v/v water and 3% v/v triisopropylsilane in TFA. The cleaved peptide was collected and the resin washed with the cleavage cocktail (1 \times 30 s). The resulting solution was concentrated by rotary evaporation and reconstituted in CH₃CN/H₂O (1:1).

5.4.3. Preparation of labeled variants

Peptides were prepared with an N-terminal label by synthesizing the peptide on-resin as described above. For the addition of a β^3 -hGly linker as in **β^3 53-8^{AI}**, the deprotection and coupling steps were repeated as above to add two Fmoc- β^3 -hGly residues. After removal of the final Fmoc protecting group, the resin was washed with DMF (6 \times 30 s), and fluorescein-5-EX succinimidyl ester (5 mg, Invitrogen) was added to the resin in 1.5 mL DMF with 15 μ L DIEA. The vessel was placed in the microwave reactor and irradiated (200 W maximum power, 60 °C, ramp 2 min, hold 6 min, cool 5 min). The resin was washed with DMF (8 \times 30 s) and CH₂Cl₂ (8 \times 30 s), dried for 20 min under N₂, and cleaved as described above. Alternatively, Alexa Fluor® 594 carboxylic acid,

Table 2

Theoretical and MALDI-TOF MS-observed molecular weights for **β^3 53-12** and **β^3 53-13** and labeled variants

β^3 -Peptide	Formula	Calcd mass	
		(M+H ⁺)	Obsd mass
$\beta^3$53-12	C ₇₃ H ₁₁₇ F ₃ N ₁₂ O ₁₅	1459.8	1458.7 (M+H ⁺), 1480.8 (M+Na ⁺)
$\beta^3$53-13	C ₇₄ H ₁₁₈ ClN ₁₃ O ₁₅	1465.3	1463.6 (M+H ⁺), 1485.3 (M+Na ⁺)
$\beta^3$53-12^{flu}	C ₉₈ H ₁₃₄ F ₃ N ₁₃ O ₂₂ S	1935.3	1937.9 (M+H ⁺), 1959.8 (M+Na ⁺)
$\beta^3$53-13^{flu}	C ₉₉ H ₁₃₅ ClN ₁₄ O ₂₂ S	1940.7	1939.9 (M+H ⁺), 1960.8 (M+Na ⁺)
$\beta^3$53-8^{AI}	C ₁₁₆ H ₁₆₃ N ₁₇ O ₂₇ S ₂	2277.7	2279.9 (M+H ⁺), 2292.6 (M+Na ⁺)
$\beta^3$53-12^{AI}	C ₁₀₉ H ₁₅₁ F ₃ N ₁₄ O ₂₅ S ₂	2164.5	2164.2 (M+H ⁺), 2186.5 (M+Na ⁺)

succinimidyl ester (2 mg, Invitrogen) was used in 1 mL DMF with 10 μ L DIEA following the same protocol.

5.4.4. β^3 -Peptide purification and characterization

Peptides were purified by reverse-phase HPLC. Identity and purity were assessed by analytical HPLC and MALDI-TOF (matrix-assisted laser desorption-ionization time-of-flight) mass spectrometry on a Voyager (Applied Biosystems) MALDI-TOF spectrometer with a 337 nm laser using the α -cyano-4-hydroxycinnamic acid matrix. Following purification, peptides were lyophilized, stored at -20°C , and reconstituted just prior to use. Theoretical and observed molecular weights are listed in Table 2.

5.5. Overexpression of hDMX₁₋₂₀₀

BC21 cells containing the pEGX-MDMX(1-200) plasmid were obtained from the lab of Professor Jiandong Chen. The sequence of the human MDMX protein N-terminal domain (residues 1–200) was cloned into the glutathione S-transferase fusion vector to obtain this plasmid. A single colony was used to inoculate a 1 L culture of 2XYT media containing 0.1 mg/mL carbenicillin. The culture was incubated at 37°C with shaking at 220 rpm until the optical density at 600 nm reached 0.9 absorbance units. After cooling the culture at 15°C for 30 min, protein expression was induced by addition of isopropyl β -D-thiogalactoside (IPTG) to a final concentration of 0.1 mM. Cells were harvested after 15 h by centrifugation at 5000g for 20 min, resuspended in PBS buffer (0.5 M NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4 , 1.8 mM KH_2PO_4 (pH 7.4), 1 mM EDTA, and 2 mM DTT) and lysed using a French press. Cell debris was pelleted by centrifugation for 20 min at 18,000g and the supernatant was incubated with glutathione sepharose at 4°C for 8 h to immobilize the fusion protein. The resin was washed with three bed volumes of chilled PBS buffer and two bed volumes of room temperature thrombin cleavage buffer (Novagen, 200 mM Tris-HCl, pH 8.4, 1.5 M NaCl, 25 mM CaCl_2 with 2 mM DTT added) before treatment with 16 units biotinylated thrombin for 12–15 h at room temperature to release hDMX₁₋₂₀₀ while leaving GST bound to the solid phase. The biotinylated thrombin was removed from hDMX₁₋₂₀₀ upon incubation with streptavidin-agarose and subsequent filtration. The resulting solution was characterized by SDS-PAGE and amino acid analysis to verify identity, purity, and protein concentration. Pure protein was distributed into aliquots and stored at -80°C until use.

5.5.1. Fluorescence polarization assays, direct binding

The synthesis and purification of p53AD, p53AD^{flu}, β 53-1, β 53-1^{flu}, β 53-8, β 53-8^{flu}, and β NEG as well as the overexpression and purification of hDM2₁₋₁₈₈ have been described previously.^{11,13} All binding experiments were performed in 384-well plates (Corning Life Sciences Plastic non-binding 384-well microplates, Fisher Scientific) with an Analyst[™] AD automated fluorescence-plate reader (LJL BioSystems, Inc.). Polarization was measured by excitation at 485 nm and subsequent measurement of the fluorescence emission at 530 nm using a 505 dichroic mirror.

Serial dilutions of hDM2₁₋₁₈₈ were prepared in Tris-buffered saline with detergent (10 mM Tris, 100 mM NaCl, 0.01% Tween-20, pH 7.4) or phosphate-buffered saline (138 mM NaCl, 2 mM KCl, 13 mM Na_2HPO_4 , 1.9 mM KH_2PO_4) with final concentrations ranging from 0.001 to 15 μM . An aliquot of fluorescently labeled peptide was added to a final concentration of 25 nM (20 μL total volume per well). The binding reaction was incubated for 30 min at room temperature. This was determined to be a sufficient length of time for the binding reaction to reach equilibrium, as there was no change in observed polarization values after 24 h. Measurements from three or more independent sets of samples were averaged for each determination. The average fluorescence polarization

values were plotted as a function of hDM2 concentration. The equation used to determine the K_d for each labeled peptide is⁷³

$$F = F_L + \left(\frac{F_{LP} - F_L}{2[L]_T} \right) \left([L]_T + [P]_T + K_d - \sqrt{([L]_T + [P]_T + K_d)^2 - 4[L]_T[P]_T} \right)$$

where

F	measured average fluorescence polarization
F_L	fluorescence polarization of free labeled peptide
F_{LP}	maximum fluorescence polarization of the peptide-protein complex
K_d	equilibrium dissociation constant of the peptide-protein complex
$[L]_T$	total concentration of labeled peptide
$[P]_T$	total concentration of protein

This assay was performed for p53AD^{flu}, β 53-1^{flu}, β 53-8^{flu}, β 53-12^{flu}, β 53-13^{flu}, β 53-8^{Al}, and β 53-12^{Al}.

5.5.2. Fluorescence polarization assays, competition

A solution of 250 nM p53AD^{flu} and 2.5 μM hDM2₁₋₁₈₈ (15 μL total volume per well) was prepared in Tris-buffered saline or phosphate-buffered saline and incubated for 30 min at room temperature. Serial dilutions of unlabeled peptide were prepared with final concentrations ranging from 12 nM to 200 μM and added to the initial solution (30 μL total volume per well with final concentrations of p53AD^{flu} and hDM2₁₋₁₈₈ of 25 nM and 1 μM , respectively). After 30 min, the fluorescence polarization of each well was measured as above. Measurements from three or more independent sets of samples were averaged for each determination. The average fluorescence polarization values were plotted as a function of inhibitor concentration. The equation used to determine the IC_{50} for each labeled peptide is:⁷⁴

$$F = F_L + \frac{(F_{LP} - F_L)}{1 + (\text{IC}_{50}/[I])^{-n}}$$

where

F	measured average fluorescence polarization
F_L	fluorescence polarization of free labeled peptide
F_{LP}	maximum fluorescence polarization of the peptide-protein complex
$[I]$	concentration of unlabeled peptide
n	Hill coefficient

The apparent affinity of the unlabeled peptide for hDM2 (K_i) can be calculated using the equation⁷⁵

$$K_i = \frac{\text{IC}_{50}}{1 + [L]_T/K_{d,L}}$$

where

$[L]_T$	total concentration of labeled peptide
$K_{d,L}$	measured K_d of L for hDM2

5.6. ELISA competition assay

The protocol for this assay was adapted from methods described by Stoll et al. and Sakurai et al. with all incubations at room temperature.^{28,47} The p53AD control peptide was synthesized as described previously¹³ and biotinylated using Biotin Polyethylene-oxide Iodoacetamide (Sigma). A black, Reacti-Bind[™] Streptavidin High Binding Capacity Coated 96-well plate pre-blocked with SuperBlock[®] Blocking Buffer (Pierce) was washed three times with TBS-T (Tris-buffered saline/0.05% Tween 20). To each well was added 50 μL of a 125 nM solution of biotinylated p53AD. The plate was incubated for 1 h and the wells were washed with TBS-T five

times. During this time, a solution of 1 μM hDM2 in TBS-T with 2% BSA and 10 mM DTT (15 μL per well) was incubated with varying concentrations of unlabeled peptide inhibitor in TBS-T with final concentrations ranging from 12 nM to 200 μM for 30 min. The solutions of peptide and protein were then distributed into the wells of the plate (30 μL per well) and incubated for 1 h. Each well was washed five times with TBS-T. The anti-MDM2 N20 antibody (Santa Cruz Biotechnology) was added to each well (100 μL per well) at a dilution of 1:250 in TBS-T with 1% BSA. After 1 h incubation on a microtiter shaker, the wells were washed five times with TBS-T, and 100 μL of anti-rabbit-IgG-HRP (Santa Cruz Biotechnology) diluted 1:1000 in TBS-T with 1% BSA was added to each well. While the plate was incubated for 1 h on a microtiter shaker, the QuantaBlu™ working solution was prepared by mixing 9 parts QuantaBlu™ Substrate Solution to 1 part QuantaBlu™ Stable Peroxide Solution (Pierce) and allowed to equilibrate to room temperature. At the end of the incubation, the wells were washed five times with TBS-T. The QuantaBlu™ working solution (100 μL per well) was added, and the plate was incubated for 1 h. The fluorescence intensity (counts per second, CPS) for each well was measured with an Analyst™ AD automated fluorescence-plate reader (LJL BioSystems, Inc.) by excitation at 325 nm and subsequent measurement of the fluorescence emission at 420 nm using a 50/50 dichroic mirror.

5.7. Flow cytometry

HCT116 cells (American Type Culture Collection, Manassas, VA) were grown in T-75 culture flasks containing McCoy's 5A media supplemented with 10% fetal bovine serum (Invitrogen) in a humidified environment with 5% CO_2 to ~80% confluency. The cells were then washed twice with 37 °C PBS and incubated with 10 mL of 37 °C PBS-based non-enzymatic cell dissociation solution (Chemicon International, Temecula, CA) for 15 min. Cells were centrifuged at 500g, resuspended in media, counted by hemocytometer, and diluted to 2200 cells/ μL with media. Aliquots of cells (230 μL) were added to fluorescein-labeled peptides (20 μL , 125 μM in PBS). Cells were incubated with the peptides for 1–4 h at 37 °C and then washed twice with 750 μL 37 °C PBS to remove extracellular peptide. To ensure removal of any surface-bound peptide,⁵¹ cells were then incubated with 500 μL 0.25% trypsin at 37 °C for 10 min, washed once with 750 μL 4 °C media and once with 750 μL 4 °C PBS. Cells were suspended in 500 μL PBS with 1 $\mu\text{g}/\text{mL}$ propidium iodide and analyzed on a BD FACScan (BD Biosciences, San Jose, CA) equipped with a 488 nm Argon laser. A total of 10,000 events were collected monitoring fluorescein and propidium iodide with 530/30 nm bandpass and 650 nm longpass filters, respectively. Events corresponding to cellular debris were removed by gating on forward and side scatter, while dead cells were removed by propidium iodide staining. Geometric means were then calculated from the histogram of fluorescence intensity and corrected for background cellular fluorescence by subtracting the geometric mean of cells treated only with PBS.

5.8. Confocal microscopy

Approximately 10^5 HCT116 cells were seeded in 2 mL media in 6-well plates containing cover glasses. After allowing the cells to adhere for 48 h, media was removed by aspiration and the cells were washed twice with 37 °C PBS. Inverted cover glasses were floated on 200 μL of media containing 20 μM peptide labeled with fluorescein for 2 h and/or 10 μM 10 kDa dextran labeled with Alexa Fluor® 647 (Invitrogen) for 15 min at 37 °C. Cover glasses were then washed with 37 °C media and PBS and mounted on microscope slides. Cells were imaged on an LSM 510 Meta (Carl Zeiss MicroImaging, Thornwood, NY) using a 488 nm Ar laser line with

a 525/25 nm filter or 633 nm HeNe laser line with a 680/30 nm filter for visualizing fluorescein and Alexa Fluor® 647, respectively.

Acknowledgments

We thank Dr. Bert Vogelstein for providing HCT116 $-/-$ cells and Dr. Jiandong Chen for providing the hDMX expression plasmid. This work was supported by the National Institutes of Health (GM 74756) and the National Foundation for Cancer Research.

References and notes

- Seebach, D.; Overhand, M.; Kuhnle, F. N. M.; Martinoni, B.; Oberer, L.; Hommel, U.; Widmer, H. *Helv. Chim. Acta* **1996**, 79, 913.
- Appella, D. H.; Christianson, L. A.; Karle, I. L.; Powell, D. R.; Gellman, S. H. *J. Am. Chem. Soc.* **1996**, 118, 13071.
- Seebach, D.; Ciceri, P. E.; Overhand, M.; Jaun, B.; Rigo, D.; Oberer, L.; Hommel, U.; Amstutz, R.; Widmer, H. *Helv. Chim. Acta* **1996**, 79, 2043.
- Cheng, R. P.; Gellman, S. H.; DeGrado, W. F. *Chem. Rev.* **2001**, 101, 3219.
- Raguse, T. L.; Lai, J. R.; Gellman, S. H. *J. Am. Chem. Soc.* **2003**, 125, 5592.
- Arvidsson, P. I.; Rueping, M.; Seebach, D. *Chem. Commun.* **2001**, 649.
- Cheng, R. P.; DeGrado, W. F. *J. Am. Chem. Soc.* **2001**, 123, 5162.
- Hart, S. A.; Bahadoor, A. B. F.; Matthews, E. E.; Qiu, X. J.; Schepartz, A. J. *Am. Chem. Soc.* **2003**, 125, 4022.
- Kritzer, J. A.; Hodsdon, M. E.; Schepartz, A. J. *Am. Chem. Soc.* **2005**, 127, 4118.
- Kritzer, J. A.; Stephens, O. M.; Guarracino, D. A.; Reznik, S. K.; Schepartz, A. *Bioorg. Med. Chem.* **2005**, 13, 11.
- Kritzer, J. A.; Luedtke, N. W.; Harker, E. A.; Schepartz, A. J. *Am. Chem. Soc.* **2005**, 127, 14584.
- Gademann, K.; Kimmerlin, T.; Hoyer, D.; Seebach, D. *J. Med. Chem.* **2001**, 44, 2460.
- Kritzer, J. A.; Lear, J. D.; Hodsdon, M. E.; Schepartz, A. J. *Am. Chem. Soc.* **2004**, 126, 9468.
- Stephens, O. M.; Kim, S.; Welch, B. D.; Hodsdon, M. E.; Kay, M. S.; Schepartz, A. J. *Am. Chem. Soc.* **2005**, 127, 13126.
- Sadowsky, J. D.; Fairlie, W. D.; Hadley, E. B.; Lee, H.-S.; Umezawa, N.; Nikolovska-Coleska, Z.; Wang, S.; Huang, D. C. S.; Tomita, Y.; Gellman, S. H. *J. Am. Chem. Soc.* **2007**, 129, 139.
- Frackenhof, J.; Arvidsson, P. I.; Schreiber, J. V.; Seebach, D. *ChemBioChem* **2001**, 2, 445.
- Chene, P. *Nat. Rev. Cancer* **2003**, 3, 102.
- Zheleva, D. I.; Lane, D. P.; Fischer, P. M. *Mini-Rev. Med. Chem.* **2003**, 3, 257.
- Momand, J.; Jung, D.; Wilczynski, S.; Niland, J. *Nucleic Acids Res.* **1998**, 26, 3453.
- Freedman, D. A.; Wu, L.; Levine, A. J. *Cell. Mol. Life Sci.* **1999**, 55, 96.
- Kussie, P. H.; Gorina, S.; Marchal, V.; Elenbaas, B.; Moreau, J.; Levine, A. J.; Pavletich, N. P. *Science* **1996**, 274, 948.
- Duncan, S. J.; Cooper, M. A.; Williams, D. H. *Chem. Commun.* **2003**, 316.
- Garcia-Echeverria, C.; Chene, P.; Blommers, M. J. J.; Furet, P. J. *Med. Chem.* **2000**, 43, 3205.
- Vassilev, L. T.; Vu, B. T.; Graves, B.; Carvajal, D.; Podlaski, F.; Filipovic, Z.; Kong, N.; Kammlott, U.; Lukacs, C.; Klein, C.; Fotouhi, N.; Liu, E. A. *Science* **2004**, 303, 844.
- Fotouhi, N.; Graves, B. *Curr. Top. Med. Chem.* **2005**, 5, 159.
- Vassilev, L. T. *J. Med. Chem.* **2005**, 48, 4491.
- Shangary, S. et al. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, 105, 3933.
- Sakurai, K.; Chung, H. S.; Kahne, D. J. *Am. Chem. Soc.* **2004**, 126, 16288.
- Hara, T.; Durell, S. R.; Myers, M. C.; Appella, D. H. *J. Am. Chem. Soc.* **2006**, 128, 1995.
- Fasan, R.; Dias, R. L. A.; Moehle, K.; Zerbe, O.; Obrecht, D.; Mittl, P. R. E.; Grutter, M. G.; Robinson, J. A. *ChemBioChem* **2006**, 7, 515.
- Yin, H.; Lee, G.; Park, H. S.; Payne, G. A.; Rodriguez, J. M.; Sebt, S. M.; Hamilton, A. D. *Angew. Chem., Int. Ed.* **2005**, 44, 2704.
- Bernal, F.; Tyler, A. F.; Korsmeyer, S. J.; Walensky, L. D.; Verdine, G. L. *J. Am. Chem. Soc.* **2007**, 129, 2456.
- Li, C.; Liu, M.; Monbo, J.; Zou, G.; Li, C.; Yuan, W.; Zella, D.; Lu, W.-Y.; Lu, W. J. *Am. Chem. Soc.* **2008**, 130, 13546.
- Shangary, S.; Ding, K.; Qiu, S.; Nikolovska-Coleska, Z.; Bauer, J. A.; Liu, M.; Wang, G.; Lu, Y.; McEachern, D.; Bernard, D.; Bradford, C. R.; Carey, T. E.; Wang, S. *Mol. Cancer Ther.* **2008**, 7, 1533.
- Appella, D. H.; Christianson, L. A.; Klein, D. A.; Powell, D. R.; Huang, X.; Barchi, J. J.; Gellman, S. H. *Nature* **1997**, 387, 381.
- Shvarts, A.; Steegenga, W. T.; Riteco, N.; van Laar, T.; Dekker, P.; Bazuine, M.; van Ham, R. C. A.; van der Houven van Oordt, W.; Hateboer, G.; van der Eb, A. J.; Jochemsen, A. G. *EMBO J.* **1996**, 15, 5349.
- Shvarts, A.; Bazuine, M.; Dekker, P.; Ramos, Y. F. M.; Steegenga, W. T.; Merckx, G.; van Ham, R. C. A.; van der Houven van Oordt, W.; van der Eb, A. J.; Jochemsen, A. G. *Genomics* **1997**, 43, 34.
- Danovi, D.; Meulmeester, E.; Pasini, D.; Migliorini, D.; Capra, M.; Frenk, R.; de Graaf, P.; Francoz, S.; Gasparini, P.; Gobbi, A.; Helin, K.; Pelicci, P. G.; Jochemsen, A. G.; Marine, J.-C. *Mol. Cell. Biol.* **2004**, 24, 5835.
- Toledo, F.; Wahl, G. M. *Int. J. Biochem. Cell Biol.* **2007**, 39, 1476.
- Fischer, P. M.; Lane, D. P. *Trends Pharmacol. Sci.* **2004**, 25, 343.

41. Parks, D. J.; LaFrance, L. V.; Calvo, R. R.; Milkiewicz, K. L.; Gupta, V.; Lattanze, J.; Ramachandren, K.; Carver, T. E.; Petrella, E. C.; Cummings, M. D. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 765.
42. Parks, D. J. et al *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3310.
43. Knight, S. M. G.; Umezawa, N.; Lee, H.-S.; Gellman, S. H.; Kay, B. K. *Anal. Biochem.* **2002**, *300*, 230.
44. Schon, O.; Friedler, A.; Bycroft, M.; Freund, S. M. V.; Fersht, A. R. *J. Mol. Biol.* **2002**, *323*, 491.
45. Lai, Z.; Auger, K. R.; Manubay, C. M.; Copeland, R. A. *Arch. Biochem. Biophys.* **2000**, *381*, 278.
46. Bottger, A.; Bottger, V.; Garcia-Echeverria, C.; Chene, P.; Hochkeppel, H.-K.; Sampson, W.; Ang, K.; Howard, S. F.; Picksley, S. M.; Lane, D. P. *J. Mol. Biol.* **1997**, *269*, 744.
47. Stoll, R. et al *Biochemistry* **2001**, *40*, 336.
48. Yu, P.; Liu, B.; Kodadek, T. *Nat. Biotechnol.* **2005**, *23*, 746.
49. Daniels, D. S.; Schepartz, A. J. *Am. Chem. Soc.* **2007**, *129*, 14578.
50. Vives, E.; Brodin, P.; Lebleu, B. *J. Biol. Chem.* **1997**, *272*, 16010.
51. Richard, J. P.; Melikov, K.; Vives, E.; Ramos, C.; Verbeure, B.; Gait, M. J.; Chernomordik, L. V.; Lebleu, B. *J. Biol. Chem.* **2003**, *278*, 585.
52. Tan, N. C.; Yu, P.; Kwon, Y.-U.; Kodadek, T. *Bioorg. Med. Chem.* **2008**, *16*, 5853.
53. Hu, B.; Gilkes, D. M.; Farooqi, B.; Sebt, S. M.; Chen, J. *J. Biol. Chem.* **2006**, *281*, 33030.
54. Bottger, V.; Bottger, A.; Garcia-Echeverria, C.; Ramos, Y. F. M.; van der Eb, A. J.; Jochemsen, A. G.; Lane, D. P. *Oncogene* **1999**, *18*, 189.
55. Hu, B.; Gilkes, D. M.; Chen, J. *Cancer Res.* **2007**, *67*, 8810.
56. Caputo, G. A.; Litvinov, R. I.; Li, W.; Bennett, J. S.; DeGrado, W. F.; Yin, H. *Biochemistry* **2008**, *47*, 8600.
57. Wade, M.; Wong, E. T.; Tang, M.; Stommel, J. M.; Wahl, G. M. *J. Biol. Chem.* **2006**, *281*, 33036.
58. Popowicz, G. M.; Czarna, A.; Rothweiler, U.; Szwagierczak, A.; Krajewski, M.; Weber, L.; Holak, T. A. *Cell Cycle* **2007**, *6*, 2386.
59. Popowicz, G. M.; Czarna, A.; Holak, T. A. *Cell Cycle* **2008**, *7*, 2441.
60. Cochran, A. G. *Chem. Biol.* **2000**, *7*, R85.
61. Arkin, M. R.; Wells, J. A. *Nat. Rev. Drug Discovery* **2004**, *3*, 301.
62. Toledo, F.; Krummel, K. A.; Lee, C. J.; Liu, C.-W.; Rodewald, L.-W.; Tang, M.; Wahl, G. M. *Cancer Cell* **2006**, *9*, 273.
63. Wender, P. A.; Mitchell, D. J.; Pattabiraman, K.; Pelkey, E. T.; Steinman, L.; Rothbard, J. B. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 13003.
64. Smith, B. A.; Daniels, D. S.; Coplin, A. E.; Jordan, G. E.; McGregor, L. M.; Schepartz, A. J. *Am. Chem. Soc.* **2008**, *130*, 2948.
65. Rueping, M.; Mahajan, Y. R.; Sauer, M.; Seebach, D. *ChemBioChem* **2002**, *3*, 257.
66. Umezawa, N.; Gelman, M. A.; Haigis, M. C.; Raines, R. T.; Gellman, S. H. *J. Am. Chem. Soc.* **2002**, *124*, 368.
67. Anantharamaiah, G. M.; Roeske, R. W. *Tetrahedron Lett.* **1982**, *23*, 3335.
68. Mzengeza, S.; Venkatachalam, T. K.; Rajagopal, S.; Diksic, M. *Appl. Radiat. Isot.* **1993**, *44*, 1167.
69. Bhushan, R.; Bruckner, H. *Amino Acids* **2004**, *27*, 231.
70. Chang, C. D.; Waki, M.; Ahmad, M.; Meienhofer, J.; Lundell, E. O.; Haug, J. D. *Int. J. Pept. Protein Res.* **1980**, *15*, 59.
71. Ten Kortenaar, P. B. W.; Vandijk, B. G.; Peeters, J. M.; Raaben, B. J.; Adams, P.; Tesser, G. I. *Int. J. Pept. Protein Res.* **1986**, *27*, 398.
72. Murray, J. K.; Gellman, S. H. *J. Comb. Chem.* **2005**, *8*, 58.
73. Heyduk, T.; Lee, J. C. *Proc. Natl. Acad. Sci. U.S.A.* **1990**, *87*, 1744.
74. Wohland, T.; Friedrich, K.; Hovius, R.; Vogel, H. *Biochemistry* **1999**, *38*, 8671.
75. Cheng, Y.; Prusoff, W. H. *Biochem. Pharmacol.* **1973**, *22*, 3099.