



Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Bioorganic & Medicinal Chemistry

journal homepage: www.elsevier.com/locate/bmc



Corrigendum

Corrigendum to “Chemistry of ecteinascidins. Part 3: Preparation of 2’-N-acyl derivatives of ecteinascidin 770 and evaluation of cytotoxicity” [Bioorg. Med. Chem. 19 (2011) 4421–4436]

Panithi Saktrakulkla^{a,c}, Satoru Toriumi^b, Mitsuhiro Tsujimoto^b, Chamnan Patarapanich^a,
Khanit Suwanborirux^{c,*}, Naoki Saito^{b,*}

^a Department of Food and Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Pathumwan, Bangkok 10330, Thailand

^b Graduate School of Pharmaceutical Sciences, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo 204-8588, Japan

^c Center for Bioactive Natural Products from Marine Organisms and Endophytic Fungi (BNPME), Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Pathumwan, Bangkok 10330, Thailand

In Table 1, the cytotoxicity against HCT116 of compound **5** (entry 6) was incorrectly presented as 0.45. The correct value is 0.45×10^2 .

* DOI of original article: [10.1016/j.bmc.2011.06.047](https://doi.org/10.1016/j.bmc.2011.06.047)

* Corresponding authors. Tel./fax: +66 2 218 8357 (K.S.); tel./fax: +81 42 495 8794 (N.S.).

E-mail addresses: skhanit@chula.ac.th (K. Suwanborirux), naoki@my-pharm.ac.jp (N. Saito).