

Subject Index¹

Volume 10

A

- N*-Acetylbenzotriazole
 - ionic strength effect on proton inventory for water-catalyzed hydrolysis, **10**, 243
- Achiral reducing agent
 - inorganic, asymmetric reduction of alkyl aryl ketones, **10**, 104
- Acid phosphatases
 - milk and potato, kinetics of substituted aryl phosphate hydrolysis, **10**, 90
- Activation
 - chymotrypsinogen A, recent developments (bovine), **10**, 357
- Active site
 - adenosine deaminase, adenosine acetals as probes, **10**, 82
 - enzyme, serum albumin: asymmetric reduction of alkyl aryl ketones, **10**, 104
 - liver alcohol dehydrogenase, substrate orientation (horse), **10**, 1
- o*-Acyl groups
 - from blocked deoxyribotides, selective cleavage, **10**, 161
- Adenosine
 - R*- and *S*-alkylidene acetals as adenosine deaminase probes, **10**, 82
- Adenosine deaminase
 - active site-directed probes, **10**, 82
- Albumin
 - serum, binding domain: asymmetric reduction of alkyl aryl ketones (cattle), **10**, 104
- Alcohol dehydrogenase
 - 1,2-3*H*-dihydro(3,2,1-*kl*)pyridophenothiazine-3-one reduction (fungus), **10**, 75
 - liver, substrate orientation in active site (horse), **10**, 1
 - specificity (yeast), **10**, 22
- Alkyl aryl ketones
 - asymmetric reduction, in binding domain of serum albumin, **10**, 104
- Alkylation
 - protein, model reaction of alkyl halides with thiolate anions, kinetics, **10**, 329

- Alkylcyclohexanones
 - reduction by liver alcohol dehydrogenase, substrate orientation in active site (horse), **10**, 1
- Alkyl halides
 - reaction with thiolate anions, kinetics of protein alkylation, **10**, 329
- Alkylidene acetals
 - of adenosine, *R* and *S* diastereoisomers: as adenosine deaminase probes, **10**, 82
- Amines
 - N*-methyl-substituted aromatic, oxidation catalyzed by hemeproteins, EPR study, **10**, 342
- Aminopyrene
 - oxidation to free radical, **10**, 177
- Arylhydroxylamine
 - chemical degradation, **10**, 144
- Aryl phosphate
 - substituted: hydrolysis by acid phosphatases of milk and potato, kinetics, **10**, 90
- Arylsulphydrolases
 - A and B, isolation, kinetic and inhibitor studies (avian, mammalian sources), **10**, 290

B

- Benz(*a*)anthracene
 - bioactivation, steric and electronic effects, fluoro and methyl analog metabolism (rat liver), **10**, 429
- Benzenes
 - meta*- and *para*-disubstituted, induced circular dichroism of cycloamylose complexes, **10**, 255
- Benzylic thioether diastereoisomers
 - synthesis from glutathione and styrene oxide, stereochemistry, **10**, 299
- Binding
 - p*-nitrophenol to α - and β -cyclodextrin, calorimetric studies, **10**, 388
- Binding domain
 - serum albumin, asymmetric reduction of alkyl aryl ketones, **10**, 104

¹ Boldface numbers indicate volume; lightface numbers indicate pagination.

- Bioactivation
 benz(a)anthracene fluoro and methyl metabolism, steric and electronic effects on site, 10, 429
- Biomimetic reduction
 asymmetric, alkyl aryl ketones, 10, 104
- Biotin
 -deficient rats, nonstereospecific β -methylcrotonate hydration, 10, 375
- Bovine serum albumin
 binding domain, stereoselective oxidation of aromatic sulfides and sulfoxides, 10, 311
- C
- Calorimetry
p-nitrophenol binding to α - and β -cyclodextrin, 10, 388
- Carbon dioxide
 hydration by carbonic anhydrase, mechanism, 10, 115
- Carbonic anhydrase
 CO₂ hydration, mechanism, 10, 115
- Carboxylate
 general base catalysis, importance of orientation, 10, 169
- Carboxyl groups
 intramolecular catalysis of amide hydrolysis (imidazole or pyridine derivatives of maleamic acid), 10, 266
- Carcinogenicity
N-nitroso compounds, SIMCA pattern, recognition study, 10, 29
- Catalysis
 carboxylate group as general base catalyst, importance of proton orientation, 10, 169
 intramolecular, of amide hydrolysis by carboxyl and metal ion/proton agents, 10, 266
 macromolecular, comparison of modified polyethylenimines and micelles, 10, 63
 metal ions, cytochrome *c* reduction by thioglycolate, 10, 443
 nucleophilic, hydrolysis of nitrophenyl acylates, 10, 63
- Catalytic properties
 biomimetic, comparison of modified polyethylenimines with micelles, 10, 63
- 4-Chloroaniline
 oxidation catalyzed by human myeloperoxidase, 10, 324
- Chymotrypsin
 -type serine proteases, steric aspects of peptide hydrolysis catalyzed by, 10, 46
- α -Chymotrypsin
 interaction with five- and six-membered cyclic esters, kinetic evidence for covalent enzyme intermediates, 10, 118
- Chymotrypsinogen A
 recent developments in activation process (bovine), 10, 357
- Cinnamaldehydes
 enzymatic reduction (yeast), 10, 22
- Cinnamic acids
 enzymatic decarboxylation (yeast), 10, 14
- Cinnamyl alcohols
 enzymatic reduction (yeast), 10, 22
- 2-Coumaranone
 interaction with α -chymotrypsin, kinetic evidence for covalent enzyme intermediate, 10, 118
- Crotonase
 substrate stereochemistry, nonstereospecific β -methylcrotonate hydration in biotin-deficient rats, 10, 375
- Cumene hydroperoxide
 metmyoglobin-initiated radical decomposition, EPR, 10, 177
- Cycloamylose
 complexes, induced circular dichroism with *meta*- and *para*-disubstituted benzenes, 10, 255
- α -Cyclodextrin
p-nitrophenol binding, calorimetric studies, 10, 388
- β -Cyclodextrin
p-nitrophenol binding, calorimetric studies, 10, 388
- Cytochrome *c*
 reduction by thioglycolate, metal catalyst effect, 10, 443
- D
- Deamination
 stereospecific, K_m and V_{max} , 10, 82
- Decarboxylation
 enzymatic, cinnamic acids (yeast), 10, 14
- Deoxyribotides
 blocked, selective cleavage of *o*-acyl protecting groups, 10, 161
- N',N''*-Diacylhydrazines
 synthesis, 10, 189
- N,N*-Dialkylanilines
 horseradish peroxidase-catalyzed oxidation with hydrogen peroxide and oxygen, electronic factors and lipophilicity, 10, 283

- Dichroism
induced circular, cycloamylose complexes with *meta*- and *para*-disubstituted benzenes, 10, 255
- 3,4-Dihydrocoumarin
interaction with α -chymotrypsin, kinetic evidence for covalent enzyme intermediate, 10, 118
- 1,2-3*H*-Dihydro(3,2,1-*kl*)pyridophenothiazine-3-one
reduction by alcohol dehydrogenase *in vivo* (fungus), 10, 75
- Dithiobis[9-(2-ethylenecarbamoylethylamino)-2,3-dimethoxy-6-azidoacridine]
synthesis and properties, 10, 455
- Donor-acceptor
complexes, 10, 63
- E
- Electron paramagnetic resonance
N-methyl-substituted aromatic amine oxidation catalyzed by hemeproteins, 10, 342
metmyoglobin initiation of cumene hydroperoxide radical decomposition, 10, 177
- Electron transfer
donor-acceptor, comparison of polyethylenimines and micelles as catalysts, 10, 63
- Enoyl-CoA hydratase, *see* Crotonase
- Enzyme
structure and kinetic results: substrate orientation in active site of liver alcohol dehydrogenase (horse), 10, 1
- 6-Ethyl-5-hydroxy-2,7-dimethoxy-1,4-naphthoquinone
biosynthetic study using ^{13}C labels detected by NMR and ^{14}C tracers (*Hendersonula toruloidea*), 10, 399
- F
- Fungus
Fusarium solani, glaucine oxidation, 10, 97
Rhizopus arrhizus Fisher, alcohol dehydrogenase from, 10, 75
yeast, *see* Yeast
- G
- Gastrin
big, sequence, synthesis, and immunochemistry (porcine), 10, 152
- Glaucine
stereospecific oxidation (fungus), 10, 97
- Glutathione
and styrene oxide, benzylic thioether diastereoisomer synthesis and relative stereochemistry, 10, 299
- H
- Helminthosporium maydis*
race T, host-specific pathotoxins produced in minor amounts, 10, 206
- Hemeproteins
N-methyl-substituted aromatic amine oxidation catalyzed by, EPR study, 10, 342
- Hendersonula toruloidea*
6-ethyl-5-hydroxy-2,7-dimethoxy-1,4-naphthoquinone production, biosynthetic study using ^{13}C labels and ^{14}C tracers, 10, 399
- Horseradish peroxidase
-catalyzed oxidation of *N,N*-dialkylanilines with hydrogen peroxide and oxygen, electronic factors and lipophilicity, 10, 283
- Hydration
 CO_2 by carbonic anhydrase, mechanism, 10, 115
crotonase-dependent, or β -methylcrotonyl CoA in biotin-deficient rats, 10, 375
forced electrophilic and nucleophilic behavior, 10, 189
- Hydrogen peroxide
and oxygen, horseradish peroxidase-catalyzed *N,N*-dialkylaniline oxidation, electronic factors and lipophilicity, 10, 283
- Hydrolysis
imidazole or pyridine derivatives of maleamic acid, intramolecular catalysis or carboxyl and metal ion/proton agents, 10, 266
water-catalyzed, of *N*-acetylbenzotriazole, ionic strength effect on proton inventory, 10, 243
nitrophenyl acylates, nucleophilic catalysis: comparison of modified polyethylenimines and micelles, 10, 63
peptide, catalyzed by serine proteases of chymotrypsin type: stereoelectronic control, 10, 46
substituted aryl phosphate, by acid phosphatases of milk and potato: kinetics, 10, 90
- β -*o*-Hydroxyphenylethanesulfonic acid sultone
interaction with α -chymotrypsin, kinetic evidence for enzyme binding, 10, 118

- o*-Hydroxy- α -toluenesulfonic acid sultone
interaction with α -chymotrypsin, kinetic evidence for covalent enzyme intermediate, **10**, 118

I

- Imidazole
role in carbonic anhydrase hydration of CO₂, **10**, 115
- Ionic strength
effect on proton inventory for water-catalyzed *N*-acetylbenzotriazole, **10**, 243
- Isovaleryl-CoA dehydrogenase
substrate stereochemistry of 2-Pro-R hydrogen elimination (biotin-deficient rats), **10**, 200

K

- Kinetics
alkyl halides with thiolate anions, model reaction for protein alkylation, **10**, 329

L

- Lipophilicity
horseradish peroxidase-catalyzed *N,N*-dialkylaniline oxidation with hydrogen peroxide and oxygen, **10**, 283
- Liver
alcohol dehydrogenase, substrate orientation in active site (horse), **10**, 1
- benz(*a*)anthracene fluoro and methyl analog metabolism, steric and electronic effects on bioactivation site (rat), **10**, 429

M

- Macromolecular catalysis
comparison of modified polyethylenimines and micelles, **10**, 63
- Maleamic acid
imidazole or pyridine derivatives, intramolecular catalysis of hydrolysis by carboxyl and metal ion/proton agents, **10**, 266
- Metabolism
benz(*a*)anthracene fluoro and methyl analogs, steric and electronic effects (rat liver), **10**, 429
- oxidative, by microbiological systems, **10**, 97

- β -Methylcrotonyl CoA
crotonase-dependent hydration (biotin-deficient rats), **10**, 375

- Metmyoglobin
initiation of cumene hydroperoxide radical decomposition, EPR, **10**, 177

Micelles

- biomimetic catalytic properties, comparison with modified polyethylenimines, **10**, 63

Microbiological systems

- in organic synthesis, glaucine stereospecific oxidation (fungus), **10**, 97

Milk

- acid phosphatase, kinetics of substituted aryl phosphate hydrolysis by (cattle), **10**, 90

Models

- amine-derived carcinogen chemical degradation, **10**, 144

Myeloperoxidase

- 4-chloroaniline oxidation catalyzed by (human), **10**, 324

N

Nitrobenzene

- general catalysis and formation in phenylhydroxylamine oxidation in aqueous phosphate buffer, **10**, 144

p-Nitrophenol

- binding to α - and β -cyclodextrin, calorimetric studies, **10**, 388

Nitrophenyl acylates

- hydrolysis, nucleophilic catalysis: comparison of modified polyethylenimines and micelles, **10**, 63

Nitrosobenzene

- spin-trapping reactions in biological oxidation reactions, **10**, 177

N-Nitroso compounds

- carcinogenicity: classification by SIMCA pattern recognition (rat), **10**, 29

Nuclear magnetic resonance

- 6-ethyl-5-hydroxy-2,7-dimethoxy-1,4-naphthoquinone biosynthesis (*Hendersonula toruloides*), **10**, 399

Nucleophilic catalysis

- hydrolysis of nitrophenyl acylates, **10**, 63

O

Oxidation

- 4-chloroaniline catalyzed by human myeloperoxidase, **10**, 324

- N*-methyl-substituted aromatic amines catalyzed by heme proteins, EPR study, **10**, 342
- stereoselective, aromatic sulfides and sulfoxides in binding domain of bovine serum albumine, **10**, 311
- stereospecific and quantitative, glaucine (fungus), **10**, 97
- Oxygen
- and hydrogen peroxide, horseradish peroxidase-catalyzed *N,N*-dialkylaniline oxidation, electronic factors and lipophilicity, **10**, 283
- P
- Papain
- acylated crude, stereoselectivity, stereospecificity, **10**, 189
 - crude, forced stereochemical control, **10**, 189
- Pathotoxins
- host-specific, produced by *Helminthosporium maydis* race T, **10**, 206
- Pattern
- recognition, SIMCA: carcinogenicity of *N*-nitroso compounds (rat), **10**, 29
- Peptide
- highly acidic, synthesis by improved solid-phase method from nucleolar nonhistone protein C23, **10**, 219
 - hydrolysis, catalyzed by serine proteases of chymotrypsin type: stereoelectronic control, **10**, 46
- Phenylhydroxylamine
- oxidation in aqueous phosphate buffer, general catalysis and formation of nitrobenzene, **10**, 144
- Polyethylenimines
- modified, biomimetic catalytic properties: comparison with micelles, **10**, 63
- Potato
- acid phosphatase, kinetics of substituted aryl phosphate hydrolysis (cattle), **10**, 90
- Probes
- stereochemical, for active site of adenosine deaminase, **10**, 82
- Prohormones
- synthesis (porcine), **10**, 152
- Protein
- acyl carrier, semisynthesis, **10**, 277
- Protein C23
- nucleolar nonhistone, highly acidic peptide synthesis by improved solid-phase method, **10**, 219
- Proton
- inventory, ionic strength effect for water-catalyzed *N*-acetylbenzotriazole hydrolysis, **10**, 243
 - orientation, importance in general base catalysis by carboxylate, **10**, 169
- R
- Radicals
- oxidation by cumene hydroperoxide initiated by heme proteins, **10**, 177
- Radioimmunoassays
- big gastrin (porcine), **10**, 152
- Reducing agent
- achiral inorganic, asymmetric reduction of alkyl aryl ketones, **10**, 104
- Reduction
- asymmetric, in binding domain of serum albumin: alkyl aryl ketones, **10**, 104
 - enzymatic, cinnamyl alcohols and cinnamaldehydes (yeast), **10**, 22
- Resolution
- alkaloidal recemate, by microbiological systems, **10**, 97
- S
- Saccharomyces cerevisiae*, see Yeast
- Saxitoxin
- derivatives: preparation, spectral characterization, chromatographic and biological properties, **10**, 412
- Serine proteases
- chymotrypsin type, steric aspects of peptide hydrolysis catalyzed by, **10**, 46
- Stereochemistry
- crotonase-dependent β -methylcrotonyl CoA hydration (biotin-deficient rats), **10**, 375
- Structure
- activity relationships, carcinogenicity of *N*-nitroso compounds (rat), **10**, 29
- Styrenes
- formation from cinnamic acid, decarboxylation by yeast, **10**, 14
- Styrene oxide
- and glutathione, benzylic thioether diastereoisomer synthesis and relative stereochemistry, **10**, 299
- Substrate
- orientation in active site of liver alcohol dehydrogenase (horse), **10**, 1

Sulfides

aromatic, stereoselective oxidation in binding
domain of bovine serum albumin, **10**, 311

Sulfonyl- α -chymotrypsin

isolation and characterization, **10**, 133

Sulfoxides

aromatic, stereoselective oxidation in binding
domain of bovine serum albumin, **10**, 311

Sulfur

atom of coenzyme thiamine pyrophosphate,
catalytic role in stabilizing negative charge
developing in transition states of biological
reactions, **10**, 233

Synthesis

organic, and microbiological systems, **10**, 97

T

Tetrahydrofurans

C₃₇-C₄₅, structure/toxicity relationship, **10**,
206

Thiamine pyrophosphate

catalytic role of thiazolium sulfur atom in
stabilizing negative charge developing in
transition states of biological reactions,
10, 233

Thioglycolate

cytochrome *c* reduction, metal catalyst effect,
10, 443

Thiolate anions

reaction with alkyl halides, kinetics of protein
alkylation, **10**, 329

Y

Yeast

Saccharomyces cerevisiae

reduction of cinnamyl alcohols and cinnamaldehydes by, **10**, 22

styrene formation from cinnamic acid, decarboxylation by, **10**, 14