

The thermodynamic mechanism of protein stabilization by trehalose

Guifu Xie, Serge N. Timasheff *

Graduate Department of Biochemistry, Brandeis University, Waltham, Massachusetts 02254, USA

Received 14 May 1996; revised 16 August 1996; accepted 16 August 1996

Abstract

The stabilization of ribonuclease A by α - α -trehalose was studied by preferential interaction and thermal unfolding. The protein is stabilized by trehalose at pH 2.8 and pH 5.5. Wyman linkage analysis showed increased exclusion of trehalose from the protein domain on denaturation. Preferential interaction measurements were carried out at 52°C at pH 5.5 and 2.8, where the protein is native and unfolded, respectively, and at 20°C where the protein is native at both pH values. At the low temperature, the interaction was preferential exclusion. At 52°C the interaction was that of preferential binding, greater to the native than the unfolded protein, the increment on denaturation being identical to that deduced from the Wyman analysis. The stabilizing effect of trehalose can be fully accounted by the change in transfer free energy on unfolding. The temperature dependence of the preferential interactions of 0.5 M trehalose with ribonuclease A showed that it is the smaller preferential binding to the unfolded protein than to the native one which gives rise to the stabilization. A thermodynamic analysis of the data led to approximate values of the transfer enthalpies and transfer entropies for the trehalose–ribonuclease A system.

Keywords: Trehalose; Preferential interactions; Transfer free energy; Transfer enthalpy; Transfer entropy; Protein stabilization; Co-solvents

1. Introduction

It is well known that the native conformation of many proteins and enzymes can be stabilized by polyhydric molecules such as glycerol and sucrose at high concentration (≥ 1 M) [1–15]; and Xie and Timasheff, manuscript in preparation. Studies on the mechanism of interaction of these compounds with proteins have shown that, at room temperature, they are preferentially excluded from contact with the protein surface [10,12–15]. This means that the addition of these co-solvents to an aqueous protein solu-

tion is thermodynamically unfavorable since it raises the chemical potential of both the protein and the additive.

The sugar, α , α -trehalose, α -D-glucopyranosyl-(1 \rightarrow 1)- α -D-glucopyranoside, is a carbohydrate found in living organisms. It is a nonreducing glucose disaccharide commonly found at high concentrations in anhydrobiotic organisms (as high as 20% of the dry weight) [16–19]. It has been identified as being of potential biotechnological importance, due to its effectiveness in stabilizing membrane structure in the dry state, inhibiting biological damage at low temperature and stabilizing protein structure during freezing and freeze drying [19–22]. The detailed

* Corresponding author.

mechanism by which trehalose produces these stabilizing effects has not yet been established. It is known that trehalose binds to the head group of lipids in dry bilayers [23], but the nature of the molecular interactions in solution is not yet known [24–26]. Trehalose has a greater ability to preserve membranes in anhydrobiotic organisms and to stabilize enzymes during the freeze-drying procedure than glycerol, glucose, sucrose and galactose [19,20,22].

Thermodynamic analyses of the interactions of sucrose, glucose and lactose [10,12] have led to the conclusion that these sugars stabilize the native structure of proteins by inducing a greater preferential hydration of denatured, unfolded proteins than of the native structure. Yet this hypothesis has not been rigorously tested. It seemed of interest, therefore, to examine the detailed mechanism of this stabilization by carrying out parallel preferential interaction measurements of the protein with sugar solutions in both the native and denatured states. Trehalose was selected due to its chemically inert nature. The results of these studies are presented in this paper.

2. Materials and methods

2.1. Materials

α,α -Trehalose dihydrate was purchased from Pfanstiehl Laboratories Inc. The pure product is a white fine powder with an absorbance at 277 nm of a 0.5 M aqueous solution lower than 0.04. Some lots were rejected, since the absorbance value was as high as 0.10 and the solution was yellow. Because of its high cost, the trehalose was recycled after use by recrystallizing it three times from distilled water. Ribonuclease A (RNase A) was purchased from Sigma (type II A from bovine pancreas). This protein was further purified on a Sulfoethyl-Sephadex c-25 column in 0.1 M, pH 6.47 phosphate buffer (Xie and Timasheff, manuscript in preparation), following essentially the procedure of Crestfield et al. [27,28]. Glycine was the reagent grade product from Sigma. The molecular-porous membrane tubing (23 \times 100, M.W. cutoff: 6,000–8,000) was purchased from Spectrum. Fisher standardized 1 N HCl solution was used for the pH adjustment of solutions. All solu-

tions contained 0.04 M glycine at pH 2.8, or 0.04 M sodium acetate at pH 5.5. The protein concentrations were determined by UV absorbance using an extinction coefficient of RNase A of 7.06 dL g⁻¹ cm⁻¹ at pH 2.8 and 7.12 dL g⁻¹ cm⁻¹ at pH 5.5 at 277 nm. The absorbance values of the protein in concentrated trehalose can be expected to be somewhat different from those in dilute buffer. For 0.9 M trehalose, the protein absorbance values were 7.10 dL g⁻¹ cm⁻¹ and 7.20 dL g⁻¹ cm⁻¹ for pH 2.8 and pH 5.5, respectively. While this may affect slightly the absolute values of the partial specific volumes, it should have no effect on the preferential interaction parameters.

2.2. Thermodynamics of thermal denaturation

The change in protein absorbance with changing temperature was followed on a Gilford Response II UV/Vis spectrophotometer at 287 nm. In all experiments, 1 mg ml⁻¹ RNase was used. The solutions were degassed for 3–5 min by using a water pump to prevent the production of bubbles on heating. The temperature change steps were programmed in such a way that the rate of heating was $\sim 0.35^\circ\text{C min}^{-1}$. This had the consequence that in the full heating-cooling cycles the protein solutions spent up to 4 h at high temperature ($> 40^\circ\text{C}$), since heating was continued up to 85°C before being reversed. In all cases, the cooling curves were mirror images of the heating curves. Since the interaction measurements required the protein solutions to be kept at 52°C for 20 h, control experiments were carried out following this regime. The transition curves obtained after cooling for 2 h were normal, which indicated that no irreversible changes had taken place.

2.3. Preferential interactions

The preferential interactions of the solvent components with the protein were obtained from the partial specific volumes of the protein measured with a Precision DMA-02 density meter (Anton Paar, Gratz). All measurements were made at 20°C following the procedures published previously [12,29–31]. Buffer solutions containing a given trehalose concen-

tration were filtered through a sintered-glass filter. Each preferential interaction point requires that the apparent partial specific volume be measured with and without dialysis against the solvent at the experimental temperature, giving, in turn, values at constant chemical potential, ϕ_2' , and constant molality, ϕ_2 , of co-solvent. To this end, for each point, parallel experiments were carried out. In one set, the protein solutions were dialyzed for 20 h at the experimental temperature, say 52°C. In the other set, the protein solutions were kept for 20 h at the same temperature without dialysis. Since the density measurements, which are a co-solvent concentration measurement, were carried out at 20°C, both solutions were then cooled to 20°C for 2–3 h, the one that had been dialyzed at high temperature having been transferred out of the dialysis bag. Specifically, for each point in constant molality experiments, six samples (6–20 mg) of RNase were dried in small glass test tubes over phosphorus pentoxide at 40°C for 2 days in a vacuum oven. After cooling to room temperature, about 1 ml of solution was added, sealed tightly immediately with Parafilm and left to stand overnight at the experimental temperature. When the samples had to be kept in a water bath, the tubes were sealed by 5 layers of Parafilm and 2 layers of Saran Wrap. Following this, the samples were kept at 20°C for 3–4 h prior to the densimetry experiments. In constant chemical potential experiments, i.e., at dialysis equilibrium with the co-solvent, each of seven samples of protein (8–25 mg) was dissolved in about 1.1 ml solvent solution and transferred into a dialysis bag, the molecular-porous membrane tubing, which had been boiled and rinsed in deionized water. The samples were dialyzed at the desired temperature for about 20 h against 800–1000 ml of co-solvent solution and then kept for 3–4 h at 20°C in small tubes that had been sealed tightly to avoid evaporation. Following the density measurement, each protein solution was diluted gravimetrically with solvent to a final optical density of about 0.7 ($\sim 1 \text{ mg ml}^{-1}$). The concentrations were determined spectrophotometrically with a Perkin Elmer Lambda 3B UV/Vis spectrophotometer. Individual ϕ_2^o and $\phi_2'^o$ points were measured in up to four independent experiments. The results were found to be highly reproducible both at low and high temperature.

2.4. Calculation of preferential interaction parameters

The partial specific volume of a protein is obtained from density measurements with the equation [32]

$$\phi = 1/\rho_o \times [1 - (\rho - \rho_o)/c] \quad (1)$$

where ϕ is the apparent specific volume, ρ and ρ_o are the densities of the protein solution and reference solvent, respectively, and c is the protein concentration in g ml^{-1} . ϕ_2^o and $\phi_2'^o$ were determined under conditions at which the molalities of the solvent components and their chemical potentials were, in turn, kept identical in the protein solution and in the reference solvent. Using the notation of Scatchard [33] and Stockmayer [34], component 1 = water, component 2 = protein, component 3 = additive (here trehalose), the preferential binding parameter, $(\partial g_3/\partial g_2)_{T,\mu_1,\mu_3}$, was calculated by [35]

$$(\partial g_3/\partial g_2)_{T,\mu_1,\mu_3} = \rho_o(\phi_2^o - \phi_2'^o)/(1 - \rho_o \bar{v}_3) \quad (2)$$

where g_i is the concentration of component i in grams per gram of water, T is the thermodynamic (Kelvin) temperature, μ_i is the chemical potential of component i , \bar{v}_3 is the partial specific volume of component 3. The superscript o indicates extrapolation to zero protein concentration¹. The values of ρ_o and \bar{v}_3 used are given in Table 1. The preferential binding parameter, $(\partial g_3/\partial g_2)_{T,\mu_1,\mu_3}$, expresses the excess of component 3 in the immediate domain of the protein with reference to its concentration in the bulk solvent. A positive value of this parameter means an excess of component 3, while a negative

¹ The measured values of ϕ_2 and ϕ_2' were found to be independent of protein concentration, whether the protein solution was heated or not. This is a strong indication that any effects observed are not the consequence of protein self-association or aggregation, which, by the Law of Mass Action, must increase with protein concentration. For a reversible self-association, its effects would be, furthermore, eliminated by extrapolation to zero protein concentration. Random, non-specific irreversible aggregation (coagulation) has, in previous experience, led to erratic, poorly reproducible values of the apparent partial specific volumes. Furthermore heating at 52° did not generate any measurable turbidity at 450 nm.

Table 1
Characteristic parameters of trehalose solutions

Trehalose/M	$g_3/(g/g)$	$m_3/(\text{mol}/1000\text{g H}_2\text{O})$	$\rho_0/\text{g ml}^{-1}$	$\bar{v}_3/\text{ml g}^{-1}$	γ_3	$(\partial \ln \gamma_3 / \partial m_3)$
0	0	0	0.9998 ± 0.0001			
0.2	0.0714	0.209	1.0272 ± 0.0002	0.600	1.038	0.177
0.3	0.1095	0.320	1.0407 ± 0.0002	0.602	1.058	0.177
0.4	0.1493	0.436	1.0542 ± 0.0001	0.603	1.080	0.177
0.5	0.1909	0.558	1.0676 ± 0.0001	0.604	1.104	0.177
0.7	0.2804	0.819	1.0942 ± 0.0001	0.606	1.156	0.177

value indicates its deficiency in the domain of the protein, i.e., an excess of component 1, water. The corresponding preferential hydration parameter, $(\partial g_1 / \partial g_2)_{T, \mu_1, \mu_3}$, is given by [10,36,37]

$$(\partial g_1 / \partial g_2)_{T, \mu_1, \mu_3} = -(\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3} / g_3 \quad (3)$$

In molal units, preferential binding is given by

$$(\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3} = (M_2 / M_3) (\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3} \quad (4)$$

where m_i is the molality of component i and M_i is its molecular weight. The preferential binding parameter is a direct expression of the mutual perturbation of the chemical potentials of components 2 and 3 [38–40]

$$\begin{aligned} (\partial \mu_2 / \partial m_3)_{T, P, m_2} &= (\partial \mu_3 / \partial m_2)_{T, P, m_3} \\ &= -(\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3} (\partial \mu_3 / \partial m_3)_{T, P, m_2} \end{aligned} \quad (5)$$

$$\begin{aligned} &= -(\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3} \\ &\quad \times [nRT / m_3 + nRT (\partial \ln \gamma_3 / \partial m_3)_{T, P, m_2}] \end{aligned} \quad (6)$$

where R is the universal gas constant, n is 1 for trehalose, and γ_i is the activity coefficient of component i . The values listed in Table 1 were calculated from osmotic coefficient data of Lin and Timasheff (to be published) using the relation

$$\ln \gamma_3 = (\phi_3 - 1) + \int_0^{m_3} [(\phi_3 - 1) / m_3] dm_3 \quad (7)$$

where ϕ_3 is the osmotic coefficient of the trehalose solution. The value of

$$(\partial \ln \gamma_3 / \partial m_3)_{T, P, m_2}$$

was found to be constant at 0.177 for the entire trehalose concentration range used in this study.

3. Results

3.1. Thermal stabilization

The thermal transition curves, both heating and cooling, of RNase A in varying amounts of trehalose at pH 2.8 (A and B) and pH 5.5 (C and D) are shown in Fig. 1. It is clear that addition of trehalose causes an elevation of the thermal transition temperature of RNase A. Taking the midpoint of the transition as the transition temperature (T_m), it can be seen that the thermal transition temperature increased monotonely with trehalose concentration, as shown in Table 2, the increase being greater at pH 2.8 than at pH 5.5: at 0.9 M trehalose, the transition temperature was raised by 7.8°C at pH 2.8, but only by 5.5°C at pH 5.5. The curves of Fig. 1 were analyzed in terms of a two-state transition, $N \rightleftharpoons D$. The relative amounts of the protein in the native (α) and unfolded states ($1 - \alpha$) were estimated from the UV absorbance, following the procedure of Biltonen and Lumry [41], and the equilibrium constant

$$K = \alpha / (1 - \alpha)$$

was calculated at each trehalose concentration as a function of temperature. From these values, the enthalpy of unfolding as a function of solvent composition was obtained from fitting to a truncated form of the integrated Van't Hoff equation [12,42–44]

$$\ln K = a + b(1/T) + c \ln T$$

$$\Delta H^\circ = R(cT - b) \quad (8)$$

$$\Delta C_p = Rc$$




Fig. 1. Thermal transitions of RNase A. (A), pH 2.8, 0.04 M glycine buffer, heating. (B), same as (A), cooling. (C), pH 5.5, 0.04 M sodium acetate buffer, heating. (D), same as (C), cooling. The trehalose concentrations are listed on the figure.

Table 2

Thermodynamic parameters of thermal denaturation of RNase A at pH 2.8 and pH 5.5 in trehalose–water systems

Treha- lose/M	T_m /°C Heating (unfolding)	T_m /°C Cooling (refolding)	ΔH° at T_m / kcal mol ⁻¹	ΔG° (20°C)/ kcal mol ⁻¹	$\delta(\Delta G^\circ)$ (20°C)/ kcal mol ⁻¹	$\delta(\Delta G^\circ)$ (52°C)/ kcal mol ⁻¹
pH 2.8						
0.0	40.9 ± 0.3	40.5 ± 0.4	82.7 ± 5	5.51 ± 0.3	0 ± 0.3	0 ± 0.3 ^a
0.1	41.8	41.6	82.8	5.72	0.21	0.23
0.2	42.2	42.0	83.6	5.92	0.41	0.37
0.3	43.4	43.1	83.9	6.19	0.68	0.62
0.4	44.1	43.8	84.5	6.43	0.92	0.83
0.5	45.0	45.0	85.3	6.69	1.18	1.03
0.6	45.9	45.9	86.8	7.04	1.53	1.25
0.7	47.0	46.8	88.1	7.38	1.87	1.40
0.8	47.7	47.6	89.8	7.75	2.24	1.72
0.9	48.7	48.5	90.4	8.06	2.55	2.00
pH 5.5						
0.0	60.7 ± 0.3	59.4 ± 0.4	108.3 ± 4	13.15 ± 0.4	0 ± 0.4	0 ± 0.4 ^a
0.1	61.1	60.1	107.0	13.20	0.05	0.07
0.2	61.7	61.1	108.4	13.47	0.32	0.34
0.3	62.3	61.8	107.7	13.50	0.35	0.44
0.4	62.9	62.1	106.0	13.71	0.56	0.67
0.5	63.3	62.6	105.6	13.78	0.67	0.73
0.6	64.0	62.9	104.7	13.86	0.71	0.92
0.7	64.8	64.2	107.0	14.14	0.99	1.25
0.8	65.4	64.8	106.9	14.30	1.15	1.42
0.9	66.2	65.8	110.4	14.99	1.84	1.81

^a The values of ΔG° in water at 52°C were -2.91 at pH 2.8 and 2.81 at pH 5.5.

where ΔH° is the standard enthalpy change for thermal unfolding. The values of ΔH° for RNase A appear to have little dependence on trehalose concentration both at pH 2.8 (A) and pH 5.5 (B), as shown in Table 2. There was no detectable curvature in the Van't Hoff plots. This is similar to the results obtained for RNase A in sorbitol solutions at pH 2.0 and pH 5.5 (Xie and Timasheff, manuscript in preparation). The resulting thermodynamic parameters are summarized in Table 2. The stabilizing effect of trehalose was expressed by $\delta(\Delta G^\circ) = \Delta G^\circ(\text{sorbitol solution}) - \Delta G^\circ(\text{dilute buffer})$. It is seen that the values calculated at two temperatures at both pH 2.8 and 5.5 show a consistently significant stabilization that increases with an increase in trehalose concentration.

3.2. Preferential interactions between trehalose and RNase A in the native and denatured states

The contribution of the interaction of solvent components with a protein to its stabilization can be established by preferential binding measurements to the protein in the two states at identical temperature

[45]. This, by necessity, requires that the two studies be carried out at different values of pH. The thermal transition data of Fig. 1 and Table 2 permitted the identification of such conditions as 52°C where RNase A is native at pH 5.5 and unfolded at pH 2.8. To ascertain the suitability of these conditions, it was necessary to establish first that the preferential interactions with the native protein are identical at the two pH values. This was done by carrying out dialysis equilibrium experiments with the native protein at 20°C at pH 2.8 and 5.5. The results are presented in Tables 3 and 4 and Fig. 2A. It is quite clear that, at room temperature, the preferential binding values, $(\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3}$, are essentially identical at pH 5.5 and 2.8. With the assumption that this identity of preferential interactions at room temperature is also valid at 52°C, it was possible to compare directly the measurements at the high temperature at the two pH values as those of the two states of the protein. This was further supported by the temperature study described later in which the preferential interactions remained identical up to the onset of the unfolding at pH 2.8.

Examination of Tables 3 and 4 shows that at 20°C

Table 3
Preferential interaction parameters of RNase A with aqueous trehalose solutions at pH 2.8

Treha- lose/M	$\phi_2^\circ/\text{ml g}^{-1}$	$\phi_2^\circ/\text{ml g}^{-1}$	$(\frac{\partial g_3}{\partial g_2})_{T, \mu_1, \mu_3}/$ (g/g)	$(\frac{\partial g_1}{\partial g_2})_{T, \mu_1, \mu_3}/$ (g/g)	$(\frac{\partial m_3}{\partial m_2})_{T, \mu_1, \mu_3}/$ (mol/mol)	$(\frac{\partial m_1}{\partial m_2})_{T, \mu_1, \mu_3}/$ (mol/mol)	$(\frac{\partial \mu_2}{\partial m_3})_{T, P, m_2}^{a,b}$
20°C							
0.2	0.700 ± 0.001	0.713 ± 0.001	-0.0348 ± 0.0027	+0.487 ± 0.037	-1.39 ± 0.11	+371 ± 29	+4017 ± 309
0.3	0.700 ± 0.001	0.722 ± 0.001	-0.0613 ± 0.0028	+0.560 ± 0.025	-2.45 ± 0.11	+426 ± 19	+4711 ± 214
0.4	0.700 ± 0.001	0.727 ± 0.002	-0.0781 ± 0.0043	+0.523 ± 0.029	-3.13 ± 0.17	+398 ± 22	+4504 ± 250
0.5	0.700 ± 0.002	0.736 ± 0.003	-0.1082 ± 0.0075	+0.567 ± 0.039	-4.33 ± 0.30	+431 ± 30	+4967 ± 345
0.7	0.701 ± 0.001	0.743 ± 0.003	-0.1364 ± 0.0065	+0.486 ± 0.023	-5.46 ± 0.26	+370 ± 18	+4444 ± 212
52°C							
0.2	0.703 ± 0.003	0.703 ± 0.003	0	0	0	0	0
0.3	0.703 ± 0.002	0.702 ± 0.005	+0.0028 ± 0.0098	-0.026 ± 0.089	+0.11 ± 0.39	-19 ± 67	-235 ± 822
0.4	0.704 ± 0.002	0.696 ± 0.006	+0.0232 ± 0.0116	-0.155 ± 0.078	+0.93 ± 0.47	-118 ± 59	-1484 ± 742
0.5	0.704 ± 0.002	0.691 ± 0.008	+0.0391 ± 0.0150	-0.205 ± 0.079	+1.56 ± 0.60	-156 ± 60	-1984 ± 763

^a cal (mol of trehalose)⁻¹ (mol of protein)⁻¹ in 1000 g H₂O.

^b The values of $(\frac{\partial \mu_3}{\partial m_3})_{T, P, m_2}$ at 20°C: 2890, 1923, 1439, 1147 and 814 cal (mol of trehalose)⁻² in 1000 g H₂O for 0.2, 0.3, 0.4, 0.5 and 0.7 M trehalose solutions, respectively. And the values of $(\frac{\partial \mu_3}{\partial m_3})_{T, P, m_2}$ at 52°C: 3206, 2133, 1596 and 1272 cal (mol trehalose)⁻² in 1000 g H₂O for 0.2, 0.3, 0.4 and 0.5 M trehalose solutions, respectively.

Table 4

Preferential interaction parameters of RNase A with aqueous trehalose solutions at pH 5.5

Treha- lose/M	$\phi_2^0/\text{ml g}^{-1}$	$\phi_2^0/\text{ml g}^{-1}$	$(\frac{\partial g_3}{\partial g_2})_{T,\mu_1,\mu_3}$ (g/g)	$(\frac{\partial g_1}{\partial g_2})_{T,\mu_1,\mu_3}/$ (g/g)	$(\frac{\partial m_3}{\partial m_2})_{T,\mu_1,\mu_3}/$ (mol/mol)	$(\frac{\partial m_1}{\partial m_2})_{T,\mu_1,\mu_3}/$ (mol/mol)	$(\frac{\partial \mu_2}{\partial m_3})_{T,P,m_2}^{a,b}$
20°C							
0.2	0.699 ± 0.001	0.712 ± 0.001	−0.0348 ± 0.0027	+0.487 ± 0.037	−1.39 ± 0.11	+371 ± 29	+4017 ± 309
0.3	0.698 ± 0.001	0.720 ± 0.001	−0.0613 ± 0.0028	+0.560 ± 0.025	−2.45 ± 0.11	+426 ± 19	+4711 ± 214
0.4	0.698 ± 0.001	0.724 ± 0.002	−0.0752 ± 0.0043	+0.504 ± 0.029	−3.01 ± 0.17	+383 ± 22	+4331 ± 250
0.5	0.698 ± 0.001	0.735 ± 0.001	−0.1112 ± 0.0030	+0.583 ± 0.016	−4.45 ± 0.12	+443 ± 12	+5104 ± 138
0.7	0.698 ± 0.002	0.740 ± 0.001	−0.1364 ± 0.0049	+0.486 ± 0.017	−5.46 ± 0.20	+370 ± 13	+4444 ± 159
52°C							
0.2	0.700 ± 0.002	0.698 ± 0.004	+0.0054 ± 0.0080	−0.075 ± 0.112	+0.21 ± 0.31	−57 ± 85	−673 ± 1009
0.3	0.700 ± 0.004	0.695 ± 0.003	+0.0139 ± 0.0098	−0.127 ± 0.089	+0.56 ± 0.39	−97 ± 67	−1194 ± 836
0.4	0.699 ± 0.003	0.684 ± 0.005	+0.0434 ± 0.0116	−0.291 ± 0.078	+1.74 ± 0.47	−221 ± 59	−2777 ± 740
0.5	0.699 ± 0.003	0.674 ± 0.009	+0.0752 ± 0.0180	−0.394 ± 0.094	+3.01 ± 0.72	−300 ± 72	−3829 ± 919

^a cal (mol of trehalose)^{−1} (mol of protein)^{−1} in 1000 g H₂O.^b The values of $(\frac{\partial \mu_3}{\partial m_3})_{T,P,m_2}$ at 20°C: 2890, 1923, 1439, 1147 and 814 cal (mol of trehalose)^{−2} in 1000 g H₂O for 0.2, 0.3, 0.4, 0.5 and 0.7 M trehalose solutions, respectively. And the values of $(\frac{\partial \mu_3}{\partial m_3})_{T,P,m_2}$ at 52°C: 3206, 2133, 1596 and 1272 cal (mol of trehalose)^{−2} in 1000 g H₂O for 0.2, 0.3, 0.4 and 0.5 M trehalose solutions, respectively.

the preferential interactions assume progressively negative values with an increase in trehalose concentration, the values at the two pH values remaining essentially identical. Thus, at 0.2 M trehalose,

$(\partial m_3/\partial m_2)_{T,\mu_1,\mu_3}$ is −1.4 moles sugar per mole protein and decreases to −5.46 moles sugar per mole protein at 0.7 M trehalose. At 52°C, however, the interaction is one of preferential binding at all




Fig. 2. Trehalose concentration dependence of: (A) the preferential binding of co-solvent to RNase A; (B) the preferential hydration of RNase A. (○) pH 5.5, 20°C; (●) pH 2.8, 20°C; (▽) pH 5.5, 52°C; (▼) pH 2.8, 52°C.

trehalose concentrations. The binding is stronger to the native protein than to the unfolded form. At 0.5 M trehalose, it has values of 3.0 moles of sugar per mole protein at pH 5.5 and 1.56 moles sugar per mole protein at pH 2.8. Comparison of the interactions in terms of preferential hydration, shown in Fig. 2B, demonstrates strikingly that the protein maintains a level of preferential hydration which is identical at pH 2.8 and 5.5 and does not vary significantly with trehalose concentration. The value of the preferential hydration (0.49–0.58 grams water per gram protein) is similar to values determined by Bull and Breese [46] and Kuntz [47] for a number of proteins. Its constancy indicates essentially total exclusion of trehalose from contact with the native protein. Quantitatively, this can be expressed by the relation between preferential binding and site occupancy by water and co-solvent [37,48–50]

$$(\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3} = A_3 - g_3 A_1 \quad (9)$$

where A_i is the amount of component i that is bound to (occupies sites on) component 2 as a time average, expressed in grams of i per gram of component 2. Thermodynamically, A_1 and A_3 are related by the exchange equilibria implicit in $(\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3}$ [45,51–53]. The linear plot, shown in Fig. 2A extrapolates to $A_3 = 0.005$ grams trehalose per gram of protein, i.e. to essentially no binding. At high temperature the situation is different as the data do not fall on a straight line. At all trehalose concentrations the preferential hydration is negative, more so for the native protein than for the denatured form. This indicates that the immediate domain of the protein is richer in trehalose than the bulk solvent. In molecular terms, some sugar molecules make non-thermodynamically neutral contacts with loci on the protein surface, with the number of contacts being diminished relative to those with water when the protein unfolds. The formation of such contacts does not imply any kind of specific interaction at sites or complexation of the sugar to the protein molecule. The interactions remain very weak. Preferential interactions at loci on a protein mean exchange with weakly interacting water molecules [51,52]. The exchange equilibrium constants are of the order 0.02 m^{-1} and the corresponding free energies of interaction per site are $< |0.05| \text{ kcal mol}^{-1}$ [45,53]. Therefore, a minor shift in the

relative affinities of water or trehalose for some loci on the protein surface (by $\sim 0.05 \text{ kcal mol}^{-1}$) can lead to a shift in the measured preferential interactions including a change in sign.

3.3. Transfer free energy

The free energy of interaction between protein and co-solvent, with water as reference state, is the transfer free energy, $\Delta \mu_2 = \mu_2(\text{solvent of a given composition}) - \mu_2(\text{water})$. This is obtained by integration of the preferential interaction parameter, $(\partial \mu_2 / \partial m_3)_{T, P, m_2}$, which is related by Eq. (5) to the binding measured by dialysis equilibrium

$$\begin{aligned} \Delta \mu_2 &= \int_0^{m_3} (\partial \mu_2 / \partial m_3)_{T, P, m_2} dm_3 \\ &= - \int_0^{m_3} (\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3} \\ &\quad \times (\partial \mu_3 / \partial m_3)_{T, P, m_2} dm_3 \end{aligned} \quad (10)$$

The values of the preferential interaction parameter, $(\partial \mu_2 / \partial m_3)_{T, P, m_2}$, presented in Fig. 3, were fitted to quadratic forms in trehalose concentration and integrated according to Eq. (10). The resulting data-fitting equations are listed:

Native protein (pH 2.8, 20°C):

$$\begin{aligned} (\partial \mu_2 / \partial m_3)_{T, P, m_2} &= 3.13 + 5.57 m_3 \\ &\quad - 4.31 m_3^2 (\text{r.c.:0.99}) \end{aligned} \quad (11)$$

$$\Delta \mu_2^N = 3.13 m_3 + 2.79 m_3^2 - 1.44 m_3^3$$

Native protein (pH 5.5, 20°C):

$$\begin{aligned} (\partial \mu_2 / \partial m_3)_{T, P, m_2} &= 3.80 + 1.30 m_3 \\ &\quad + 1.53 m_3^2 (\text{r.c.:0.99}) \end{aligned} \quad (12)$$

$$\Delta \mu_2^N = 3.80 m_3 + 0.65 m_3^2 + 0.51 m_3^3$$

Native protein (pH 5.5, 52°C):

$$\begin{aligned} (\partial \mu_2 / \partial m_3)_{T, P, m_2} &= 0.53 - 3.78 m_3 \\ &\quad - 7.46 m_3^2 (\text{r.c.:0.99}) \end{aligned} \quad (13)$$

$$\Delta \mu_2^N = 0.53 m_3 - 1.89 m_3^2 - 2.49 m_3^3$$

Denatured protein (pH 2.8, 52°C):

$$\begin{aligned} (\partial \mu_2 / \partial m_3)_{T, P, m_2} &= 1.02 - 3.76 m_3 \\ &\quad - 3.18 m_3^2 (\text{r.c.:0.99}) \end{aligned} \quad (14)$$

$$\Delta \mu_2^D = 1.02 m_3 - 1.88 m_3^2 - 1.06 m_3^3$$




Fig. 3. Variation of the preferential interaction parameter, $(\partial\mu_2/\partial m_3)_{T,P,m_2}$, with trehalose concentration for RNase A. (○) pH 5.5, 20°C; (●) pH 2.8, 20°C; (▽) pH 5.5, 52°C; (▼) pH 2.8, 52°C.

The transfer free energies of RNase A from water to the sugar solution, $\Delta\mu_2 = \mu_{2,s} - \mu_{2,w}$, are presented in Fig. 4 as a function of trehalose concentration. It is evident that at 20°C the interactions of the




Fig. 4. Chemical potential change of RNase A on transferring it from water to an aqueous solution of trehalose. (○) pH 5.5, 20°C; (●) pH 2.8, 20°C; (▽) pH 5.5, 52°C; (▼) pH 2.8, 52°C.

native protein with trehalose solutions at both pH values are thermodynamically unfavorable relative to pure water, as $\Delta\mu_2$ assumes progressively increasing positive values with increasing sugar concentration. At the high temperature, however, 52°C, the situation is reversed. For the native protein (pH 5.5), $\Delta\mu_2$ is essentially zero up to 0.3 M trehalose, which means that the protein is thermodynamically indifferent to contact with water or sugar solution. Above this solvent composition, the transfer free energy becomes negative, and attains a value of $-5.1 \text{ kcal mol}^{-1}$ at 0.9 M, i.e., contact between the protein and aqueous trehalose becomes highly favorable. The unfolded protein, however, exhibits a less favorable pattern of interactions with the sugar solutions. In fact, after assuming a pattern of weak repulsion, $\Delta\mu_2 = 0.1 \text{ kcal mol}^{-1}$ at 0.3 M trehalose, the interaction becomes increasingly favorable above 0.45 M, but attains a value of only $-2.6 \text{ kcal mol}^{-1}$ at 0.9 M. It is this less favorable interaction of trehalose with denatured than with native protein which causes it to stabilize proteins against unfolding, when water is taken as the reference state.

3.4. Temperature dependence of preferential interactions

The measurement of the transfer free energies of the protein in the native and denatured states into a

cosolvent system of a given composition gives the stabilizing ability of the given co-solvent. A full understanding of the thermodynamic mechanism also requires knowledge of the corresponding enthalpic and entropic parameters. This requires that the preferential interactions be measured as a function of temperature. Such measurements were, therefore, performed on the interactions of RNase A with 0.5 M trehalose at pH 2.8 and 5.5. The thermal transition data of Fig. 1 show that at pH 5.5, the protein was in the native state at temperatures below 54°C (the transition takes place between 54.0°C and 72.4°C). However, at pH 2.8, denaturation set in above 35°C (the transition region is between 34.2°C and 54.8°C). Therefore, dialysis equilibrium experiments were carried out on RNase A in 0.5 M trehalose between 4.0°C and 58.0°C at both pH values. At the high pH this gave values of the preferential interactions with

native protein as a function of temperature. At pH 2.8, however, the preferential interaction measurements scanned over the transition of the protein, native below 34°C, and denatured above 52°C, after passing through the zone of transition.

The results of these measurements, presented in Table 5, show that at isomolal conditions, the apparent partial specific volume remains essentially invariant with temperature at pH 5.5. At pH 2.8, it also was constant below the zone of transition, but increased slightly as the protein unfolded. At isopotential conditions, this parameter, ϕ_2° , changed drastically with temperature, as its value decreased sharply. As shown in Fig. 5A, the decrease in ϕ_2° was stronger at pH 5.5 than at pH 2.8. At low temperature, the values of ϕ_2° were identical at the two pH values up to 30°C. Above this temperature, ϕ_2° of the native protein started to decline sharply. At the

Table 5

Temperature dependence of the preferential interaction parameters of RNase A with 0.5 M trehalose solutions at pH 2.8 and pH 5.5

Temperature/°C	$\phi_2^{\circ}/\text{ml g}^{-1}$	$\phi_2^{\circ}/\text{ml g}^{-1}$	$(\frac{\partial g_3}{\partial g_2})_{T,\mu_1,\mu_3}/$ (g/g)	$(\frac{\partial g_1}{\partial g_2})_{T,\mu_1,\mu_3}/$ (g/g)	$(\frac{\partial m_3}{\partial m_2})_{T,\mu_1,\mu_3}/$ (mol/mol)	$(\frac{\partial m_1}{\partial m_2})_{T,\mu_1,\mu_3}/$ (mol/mol)	$(\frac{\partial \mu_2}{\partial m_3})_{T,P,m_2}$ ^{a,b}
pH 2.8							
4	0.700 ± 0.001	0.737 ± 0.002	−0.1112 ± 0.0045	+0.583 ± 0.024	−4.45 ± 0.18	+443 ± 18	+4824 ± 196
15	0.700 ± 0.001	0.736 ± 0.003	−0.1082 ± 0.0060	+0.567 ± 0.031	−4.33 ± 0.24	+431 ± 24	+4880 ± 271
20	0.700 ± 0.002	0.736 ± 0.003	−0.1082 ± 0.0075	+0.567 ± 0.039	−4.33 ± 0.30	+431 ± 30	+4967 ± 345
25	0.701 ± 0.001	0.736 ± 0.003	−0.1052 ± 0.0060	+0.551 ± 0.031	−4.21 ± 0.24	+419 ± 24	+4913 ± 281
35	0.702 ± 0.002	0.735 ± 0.005	−0.0992 ± 0.0105	+0.520 ± 0.055	−3.97 ± 0.42	+395 ± 42	+4788 ± 508
45	0.702 ± 0.002	0.734 ± 0.008	−0.0962 ± 0.0150	+0.504 ± 0.079	−3.85 ± 0.60	+384 ± 60	+4793 ± 749
48.5	0.703 ± 0.002	0.726 ± 0.005	−0.0692 ± 0.0105	+0.362 ± 0.055	−2.77 ± 0.42	+276 ± 42	+3485 ± 530
52	0.704 ± 0.002	0.691 ± 0.008	+0.0391 ± 0.0150	−0.205 ± 0.079	+1.56 ± 0.60	−156 ± 60	−1984 ± 763
58	0.704 ± 0.002	0.661 ± 0.005	+0.1293 ± 0.0105	−0.677 ± 0.055	+5.17 ± 0.42	−516 ± 42	−6700 ± 545
pH 5.5							
4	0.698 ± 0.001	0.735 ± 0.002	−0.1112 ± 0.0045	+0.583 ± 0.024	−4.45 ± 0.18	+443 ± 18	+4824 ± 196
15	0.698 ± 0.001	0.735 ± 0.002	−0.1112 ± 0.0045	+0.583 ± 0.024	−4.45 ± 0.18	+443 ± 18	+5015 ± 196
20	0.698 ± 0.001	0.735 ± 0.001	−0.1112 ± 0.0030	+0.583 ± 0.016	−4.45 ± 0.12	+443 ± 12	+5104 ± 138
30.5	0.698 ± 0.002	0.734 ± 0.001	−0.1082 ± 0.0045	+0.567 ± 0.024	−4.33 ± 0.18	+431 ± 18	+5144 ± 214
37	0.698 ± 0.002	0.725 ± 0.004	−0.0812 ± 0.0090	+0.425 ± 0.047	−3.25 ± 0.36	+324 ± 36	+3946 ± 438
43	0.699 ± 0.002	0.714 ± 0.007	−0.0451 ± 0.0135	+0.236 ± 0.071	−1.80 ± 0.54	+140 ± 42	+2227 ± 668
47.5	0.699 ± 0.002	0.692 ± 0.008	+0.0210 ± 0.0150	−0.110 ± 0.079	+0.84 ± 0.60	−84 ± 60	−1054 ± 753
52	0.699 ± 0.003	0.674 ± 0.009	+0.0752 ± 0.0180	−0.394 ± 0.094	+3.01 ± 0.72	−300 ± 72	−3829 ± 919
58	0.699 ± 0.003	0.638 ± 0.009	+0.1834 ± 0.0180	−0.961 ± 0.094	+7.34 ± 0.72	−731 ± 72	−9513 ± 936

^a cal (mol of trehalose)^{−1} (mol of protein)^{−1} in 1000 g H₂O.

^b The values of $(\frac{\partial \mu_3}{\partial m_3})_{T,P,m_2}$ were 1084, 1127, 1147, 1167, 1188, 1206, 1214, 1237, 1245, 1255, 1258, 1272 and 1296 cal (mol of trehalose)^{−2} in 1000 g H₂O for 0.5 trehalose at 4, 15, 20, 25, 30.5, 35, 37, 43, 45, 47.5, 48.5, 52 and 58°C, respectively.




Fig. 5. Temperature dependence of: (A) the apparent partial specific volume of RNase A at dialysis equilibrium; (B) the preferential binding of co-solvent to RNase A in 0.5 M trehalose solution. (○) pH 5.5; (●) pH 2.8.

acid pH, this decline set in only above 45°C, i.e. after the onset of the unfolding transition.

The preferential interaction parameters deduced from these measurements are presented in Table 5. An identical pattern of a decrease in preferential exclusion and preferential hydration with rising temperature is quite evident. The preferential trehalose

binding, $(\partial g_3 / \partial g_2)_{T, \mu_1, \mu_3}$, is compared for the two pH values in Fig. 5B. The values are negative and identical at the two pH values below 35°C, where they are essentially constant. Above this temperature they diverge as the values measured at pH 5.5 start increasing and become positive at temperatures above 45°C. The same is true at pH 2.8, but only above




Fig. 6. Temperature dependence of: (A) the preferential hydration of RNase A in 0.5 M trehalose solution; (B) the total binding (site occupancy) of co-solvent to RNase A in 0.5 M trehalose solution. (○) pH 5.5; (●) pH 2.8.

52°C. This means that the interaction of the co-solvent with the protein is greater with the native protein than with the denatured form. Expressed in terms of preferential hydration, shown in Fig. 6A, these results indicate that hydration of the protein remains constant below 35°C. Analysis in terms of Eq. (9) gives the results of Fig. 6B. These show that below 35°C there is total exclusion of trehalose from the immediate domain of the protein. Using a value of 0.49 grams water per gram protein for A_1 , which is the average measured at 20°C (Tables 3 and 4), results in a slightly negative value of the total binding of trehalose to RNase A. Above 35°C, however, there sets in a sharp increase for the native protein which at 52°C attains a value of A_3 of 0.17 grams trehalose per gram protein for the native protein, i.e. 6.8 molecules of trehalose penetrate the hydration layer and enter into contact with the protein even though the hydration remains at a constant high value. For the denatured protein, pH 2.8, at 52°C A_3 attains a value of only 0.13 grams trehalose per gram protein, or 5.3 molecules of trehalose that make contact with one molecule of protein.

3.5. Thermodynamic parameters of the preferential interactions

The transfer free energy, $\Delta\mu_2$, which is the change in the partial molal free energy, \bar{G}_2 , of the protein when transferred from water to the co-solvent system, is the sum of the contributions of the partial molal enthalpy, \bar{H}_2 , and the partial molal entropy, \bar{S}_2 . Therefore, knowledge of the concentration dependence of the preferential interaction parameter as a function temperature should permit the deduction of the necessary enthalpy and entropy parameters, since

$$\left(\frac{\partial\mu_2}{\partial m_3}\right)_{T,P,m_2} = \left(\frac{\partial\bar{H}_2}{\partial m_3}\right)_{T,P,m_2} - T\left(\frac{\partial\bar{S}_2}{\partial m_3}\right)_{T,P,m_2} \quad (15)$$

The temperature dependence of $(\partial\mu_2/\partial m_3)_{T,P,m_2}$ at the two pH values is given in Fig. 7. Its analysis in terms of Eq. (8), in which $\ln K$ is replaced by

$$-\frac{1}{RT}\left(\frac{\partial\mu_2}{\partial m_3}\right), \text{ yields } \left(\frac{\partial\bar{H}_2}{\partial m_3}\right)_{T,P,m_2} \text{ and } \left(\frac{\partial\bar{S}_2}{\partial m_3}\right)_{T,P,m_2}. \text{ The results of the calculations for native RNase A (pH 5.5) are shown in Fig. 8. Panel}$$

A gives the Van't Hoff plot of the preferential interaction parameter, while Fig. 8B gives the temperature dependence of the variation of the partial molal enthalpy with trehalose concentration obtained by applying Eq. (8) to the data of Fig. 8A. This parameter is found to increase with temperature, with values of $-87 \text{ kcal mol}^{-2}$ at 4°C and $236 \text{ kcal mol}^{-2}$ at 58°C, which gives a strongly positive value of the variation with co-solvent concentration of the partial molal heat capacity, $6.0 \text{ kcal deg}^{-1} (\text{mol prot})^{-1} (\text{mol co-solvent})^{-1}$. It is interesting to note that these values are similar to those reported for the closely related osmolyte sorbitol. At 25°C, the value of $(\partial\bar{H}_2/\partial m_3)_{T,P,m_2}$ for the interaction of 0.5 M trehalose with RNase A is 30 kcal mol^{-2} . For 30% sorbitol, it is 28 kcal mol^{-2} when the protein is RNase A (Xie and Timasheff, to be published), and 48 kcal mol^{-2} when the protein is bovine serum albumin [53a].

The transfer enthalpy, $\Delta\bar{H}_{2,m_3}$, and transfer entropy, $\Delta\bar{S}_{2,m_3}$ can be calculated by integrating $(\partial\bar{H}_2/\partial m_3)_{T,P,m_2}$ over dm_3 , since

$$\begin{aligned} \Delta\bar{H}_{2,\text{tr}} &= \Delta\bar{H}_{2,m_3} = \bar{H}_{2,m_3} - \bar{H}_{2,\text{w}} \\ &= \int_0^{m_3} \left(\frac{\partial\bar{H}_2}{\partial m_3} \right)_{T,P,m_3} dm_3 \end{aligned} \quad (16a)$$

$$T\Delta\bar{S}_{2,m_3} = \Delta\bar{H}_{2,m_3} - \Delta\mu_{2,\text{tr}} \quad (16b)$$




Fig. 7. Temperature dependence of the preferential interaction parameter, $(\partial\mu_2/\partial m_3)_{T,P,m_2}$, in 0.5 M trehalose solution, (○), pH 5.5; (●), pH 2.8. The dotted line is the parameter calculated at 52°C for the denatured protein (see text).




Fig. 8. Thermodynamics of the preferential interactions of aqueous trehalose solutions with native RNase A. (A) Van't Hoff plot of the transfer free energy variation with trehalose concentration: (∇) 0.2 M trehalose; (\blacktriangledown) 0.3 M trehalose; (\square) 0.4 M trehalose; (\bullet) 0.5 M trehalose. (B) Temperature dependence of the variation of the transfer enthalpy with trehalose concentration at 0.2 M (∇), 0.3 M (\blacktriangledown), 0.4 M (\square), 0.5 M (\bullet) trehalose. The numbers on the figure are the slopes, which represent $(\partial\bar{C}_{P,2}/\partial m_3)_{T,P,m_2}$ in $\text{cal deg}^{-1} (\text{mol prot})^{-1} (\text{mol co-solvent})^{-1}$. C. Dependence on trehalose concentration of the transfer enthalpy variation with trehalose concentration at 20°C and 52°C. The dotted line is the parameter calculated at 52°C for the denatured protein. D. Dependence of the transfer enthalpy on trehalose concentration at 20°C and 52°C. The dotted line is the parameter calculated at 52°C for the denatured protein. E. Temperature dependence of the variation of the transfer entropy with trehalose concentration at 0.5 M trehalose. F. Dependence of the transfer entropy on trehalose concentration at 20°C and 52°C.

In the absence of detailed measurements of the temperature dependence of preferential interactions at several co-solvent concentrations, an approximate calculation was carried out with the assumption that the form of the temperature dependence of

$(\partial\mu_2/\partial m_3)$ is the same at all co-solvent concentrations. To that end the data at 20°C and 52°C at 0.2, 0.3 and 0.4 M trehalose were plotted in a manner similar to Fig. 8A and subjected to the Van't Hoff analysis of Eq. (8). The resulting values are shown in

Fig. 8B as a function of temperature and in Fig. 8C as a function of trehalose concentration. Integration according to Eq. (16a) gave the values $\Delta H_{2, \text{tr}}$ presented in Fig. 8D. Application of Eq. (16b) to the data of Fig. 7 and Fig. 8B, as well as those of Fig. 8D and Fig. 4 gave the entropy parameters of Fig. 8E and F. While the numbers shown in Fig. 8C, D, and F are highly approximate and should be regarded as indicative of trends, nevertheless they are useful in permitting some qualitative conclusions. With this caution, it can be stated that the transfer of RNase A from water into aqueous trehalose at high temperature is accompanied by large concentration-dependent positive changes in enthalpy and entropy which are close to compensating, while at room temperatures these changes are much smaller, reflecting the positive value of $\Delta \bar{C}_{p,2}$.

4. Discussion

In the absence of some non-identified and non-detected, highly reproducible artifact or secondary effect, which mimics exactly the linkage between protein unfolding and changes in preferential interactions, the current results give a full, even though in parts highly approximate, description of the thermodynamics of the stabilization of a protein (RNase A) by an osmolyte (trehalose) in thermal unfolding. Let us examine this in terms of the linkage concept taking as reference state first the solvent of a given composition and then dilute aqueous buffer, i.e. water.

4.1. Wyman analysis of the stabilization

The effect of a ligand on any equilibrium at a given concentration of ligand, which is the reference state, is expressed by the Wyman linkage equation [45,53–56]

$$\frac{\partial \ln K}{\partial \ln a_3} = \left(\frac{\partial m_3}{\partial m_2} \right)_{T,P,\mu_3}^D - \left(\frac{\partial m_3}{\partial m_2} \right)_{T,P,\mu_3}^N \quad (17)$$

where a_3 is the activity of component 3, the co-solvent, and D and N refer specifically to the denaturation equilibrium $N \rightleftharpoons D$. Therefore, the slope of a

log–log plot of the equilibrium constant versus the activity of the co-solvent, trehalose in the present case, gives the change in preferential binding of the ligand to the protein that accompanies the transformation of the protein. A plot of the data of Fig. 1 in terms of Eq. (17) is given in Fig. 9. Values of slopes are written on the figure. The negative values show that denaturation of RNase A in the presence of trehalose is accompanied by a decrease in the preferential binding of trehalose to the protein, and that this effect increases with trehalose concentration. Specifically, at 52°C, for 0.5 M trehalose, the slope has a value of -1.40 ± 0.05 moles trehalose per mole protein. This should be equal to the difference between the preferential bindings measured directly on the two end states of the reaction. The data of Tables 3 and 4 give values of $(\partial m_3 / \partial m_2)_{T,P,\mu_3}$ of 3.01 moles trehalose per mole protein for the native protein (pH 5.5) and 1.56 moles of trehalose per




Fig. 9. Linkage plots of the effect of trehalose on the thermal denaturation of RNase A at (A) pH 2.8 and (B) pH 5.5. The numbers on the figures are the slopes at the given points.

mole of protein for the unfolded form (pH 2.8). Their difference, $1.56 - 3.01 = -1.45$ moles trehalose per mole protein, is essentially identical with that given by the Wyman slope. This identity shows that stabilization of RNase A by trehalose is modulated by general weak protein–co-solvent interactions, expressed through preferential binding. The small value of the change in preferential interactions in fact reflects the change in the actual number of trehalose–protein contacts which have much greater numerical values. Coming back to the decomposition of the preferential interactions into total protein surface occupancy by water and trehalose, as expressed through Eq. (9), gives $(A_3^D - A_3^N) = 5.32 - 6.76 = -1.44$ molecules of trehalose occupying sites on a protein molecule. The much larger values of A_3 than $(\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3}$ reflect the fact that the latter is a pure thermodynamic number that encompasses water–trehalose exchange at loci on the protein surface, as well as non-exchangeable water [45,51–53]. In addition to A_3 , $(\partial m_3 / \partial m_2)_{T, \mu_1, \mu_3}$ contains some 450 molecules of water that occupy sites on the protein surface. On the other hand, site-occupancy values are non-thermodynamic quantities that reflect protein–ligand contacts that may be detected by some particular techniques, such as calorimetry or spectroscopy, but which, by neglecting exchange with water, do not provide a measure of the global interactions of the protein surface with its environment [45,53]. Finally, the results of the present study clearly demonstrate that stabilization of the native protein structure does not require preferential exclusion of the co-solvent from the protein surface. In the present example, at 52°C trehalose is preferentially bound to both the native and denatured proteins. What provides the stabilization is the difference between these two quantities. Preferential binding to the denatured protein is smaller than that to the native form. Hence, there is stabilization. At lower temperatures, trehalose is preferentially excluded from native protein. For example, at 43°C, the preferential interaction with native protein (pH 5.5) is -1.80 moles of sugar per mole of protein. Wyman analysis gives a slope of -1.72 moles sugar per mole of protein. This predicts that the preferential interaction with denatured protein would be one of exclusion at a level of -3.5 moles of sugar per mole of protein. The finding that at low temperature stabi-

lization is determined by an increase in preferential exclusion of trehalose on denaturation of RNase A, while at 52°C it is determined by a decrease in preferential binding of trehalose to the protein, does not imply in any way a change in the mechanism of stabilization. These observations are a trivial consequence of the fact that preferential interaction, exclusion or binding, is simply a measure of the gradient with co-solvent concentration of the transfer free energy (see Eq. (5) and Eq. (10)). Preferential interaction indicates, therefore, whether $\Delta\mu_2$ is increasing or decreasing at the given co-solvent concentration. Stabilization with respect to water, which is the commonly sought parameter, is given by the difference between the transfer free energies of the denatured and native proteins. (For a detailed discussion, see Refs. [45,53]).

4.2. Stabilization relative to pure aqueous medium

Integration of the Wyman equation (Eq. (17)) shows that, with water as the reference state, stabilization expressed as the increment of the standard free energy of denaturation in a given medium over that in pure water, $\delta\Delta G^\circ = \Delta G_s^\circ - \Delta G_w^\circ$, is equal to the increment in transfer free energy upon denaturation, $\delta\Delta\mu_2 = \Delta\mu_2^D - \Delta\mu_2^N$, i.e.

$$\delta\Delta G^\circ = \Delta G_s^\circ - \Delta G_w^\circ = \Delta\mu_2^D - \Delta\mu_2^N = \delta\Delta\mu_2 \quad (18)$$

The thermal unfolding results given in Fig. 1 and Table 2 were combined with the transfer free energies calculated for native and denatured protein at 52°C from the preferential interaction data of Tables 3 and 4 with the use of Eq. (13) and Eq. (14). The results are summarized in Table 6. It is seen that, at all trehalose concentrations, the two sides of Eq. (18), after rearrangement into a thermodynamic box, are identical within experimental error. Therefore, the change in transfer free energy is sufficient to account for the stabilization. This means that there is no need to invoke any specific interactions, such as binding of the sugar at particular structure stabilizing loci, to account for its action. General, non-specific thermodynamic interaction is sufficient.

Table 6

Closing of the thermodynamic box, $(\Delta G_w^o + \Delta \mu_2^D) = (\Delta G_s^o + \Delta \mu_2^N)$, in the stabilization of RNase A by trehalose at 52°C

Trehalose/ M	$\Delta G_w^o + \Delta \mu_2^D /$ kcal mol ⁻¹	$\Delta G_s^o + \Delta \mu_2^N /$ kcal mol ⁻¹
0.1	-2.91 + 0.08 = -2.8	-2.68 + 0.03 = -2.7
0.2	-2.91 + 0.12 = -2.8	-2.54 + 0.01 = -2.5
0.3	-2.91 + 0.10 = -2.8	-2.29 - 0.11 = -2.4
0.4	-2.91 + 0.00 = -2.9	-2.08 - 0.34 = -2.4
0.5	-2.91 - 0.20 = -3.1	-1.88 - 0.73 = -2.6
0.6	-2.91 - 0.52 = -3.4	-1.66 - 1.33 = -3.0
0.7	-2.91 - 1.01 = -3.9	-1.51 - 2.20 = -3.7
0.8	-2.91 - 1.69 = -4.6	-1.19 - 3.44 = -4.6
0.9	-2.91 - 2.62 = -5.5	-0.91 - 5.12 = -6.0

The stabilization, expressed by the change in transfer free energy upon denaturation, is a resultant of such changes in the transfer enthalpy and transfer entropy during denaturation, since

$$\delta \Delta \mu_{2, \text{tr}} = \Delta \bar{H}_{2, \text{tr}}^D - T \Delta \bar{S}_{2, \text{tr}}^D - \Delta H_{2, \text{tr}}^N + T \Delta S_{2, \text{tr}}^N \quad (19)$$

The analysis described in Fig. 8 has given approximate $\Delta \bar{H}_{2, \text{tr}}$ and $\Delta \bar{S}_{2, \text{tr}}$ values for the native protein. For denatured protein, data are even more scarce. Nevertheless, approximate values for the denatured protein below the transition temperature were generated by extrapolating the $(\partial \mu_2 / \partial m_3)_{T, P, m_2}$ curve at

pH 2.8 to the low temperature zone. This extrapolation was based on two assumptions: 1) that $(\partial m_3 / \partial m_2)_{T, P, \mu_3}^D - (\partial m_3 / \partial m_2)_{T, P, \mu_3}^N$ is independent of temperature, and 2) that, in the transition region, the preferential interactions are the weighted average of the parameters for the native and denatured states of the protein, i.e.

$$\left(\frac{\partial \mu_2}{\partial m_3} \right)_{T, P, m_3}^{\text{Exper}} = f_N \left(\frac{\partial \mu_2}{\partial m_3} \right)_{T, P, \mu_3}^N + f_D \left(\frac{\partial \mu_2}{\partial m_3} \right)_{T, P, \mu_3}^D \quad (20)$$

where f_N and f_D are the fractions of the protein in the native and denatured states. The resulting calculated temperature dependence of $(\partial \mu_2 / \partial m_3)_{T, P, \mu_3}^D$ for the denatured protein below the transition is shown by the dotted line of Fig. 7. Construction of a Van't Hoff plot, given in Fig. 10A and application of Eq. (8) permitted the calculation of the temperature dependence of the partial molal enthalpy and entropy variations with trehalose concentration. The results are plotted in Fig. 10B and C. Both parameters are seen to increase with temperature, as had been found for the native protein. While the highly approximate nature of these calculations requires that the resulting numbers be considered only as estimates, nevertheless it would seem that the thermodynamic parameters for denatured RNase A follow a dependence on




Fig. 10. Temperature dependence of the variation with trehalose concentration of the calculated transfer free energy (A), transfer enthalpy (B) and transfer entropy (C) of denatured RNase A in 0.5 M aqueous trehalose solution.

Table 7

Transfer enthalpies of native and denatured RNase A at 52°C in aqueous trehalose solutions

Trehalose/M	m_3 /(mol/1000 g H ₂ O)	$\Delta H_{m_3}^o - \Delta H_w^o$ ^{a,c}	$\Delta \bar{H}_{2, \text{tr}}^N$ ^a	$\Delta \bar{H}_{2, \text{tr}}^D$ ^{a,c}	$(\frac{\partial \Delta \bar{H} - 2, \text{tr}^D}{\partial m_3})_{T, P, m_2}$ ^{b,c}
0.2	0.209	0.1 (0.8)	23.3	23.4 (24.1)	127.4(140.6)
0.3	0.320	−0.6 (1.2)	39.6	39.0 (40.8)	147.4(161.9)
0.4	0.436	−2.4 (1.8)	59.2	56.8 (61.0)	168.3(184.1)
0.5	0.558	−3.3 (2.5)	82.3	79.0 (84.8)	190.3(207.4)

^a kcal mol protein^{−1}.^b kcal mol protein^{−1} mol trehalose^{−1}.^c The numbers listed are at pH 5.5; numbers in parentheses are at pH 2.8.

trehalose concentration that is qualitatively similar to that of the native protein.

Finally, the availability of these values, even though estimated and therefore highly approximate, has permitted us to evaluate the magnitude of the contribution of interactions with the co-solvent to the measured standard enthalpy and entropy of denaturation. This contribution can be expressed by relations similar to Eq. (18) for the free energy. For the standard enthalpy change, the increment, $\delta\Delta H^o$, is given by

$$\begin{aligned}\delta\Delta H^o &= \Delta H_{m_3}^o - \Delta H_w^o = \Delta \bar{H}_{2, \text{tr}}^D - \Delta \bar{H}_{2, \text{tr}}^N \\ &= \delta\Delta \bar{H}_{2, \text{tr}}^{N-D}\end{aligned}\quad (21)$$

The calculations are summarized in Table 7, where $\Delta \bar{H}_{2, \text{tr}}^D$ was calculated from $(\Delta H_{m_3}^o - \Delta H_w^o + \Delta \bar{H}_{2, \text{tr}}^N)$ (Eq. (21)), using the data of Table 2 and Fig. 8D. The corresponding values of the slope, e.g. 199 ± 10 kcal mol protein^{−1} mol trehalose^{−1} at 0.5 M trehalose, agree well with those calculated from the extrapolated values of $(\partial\mu_2/\partial m_3)_{T, P, m_2}$, namely 196 kcal mol protein^{−1} mol trehalose^{−1}, presented in Fig. 10B.

Examination of the values of the various measured and calculated (even though very approximately) parameters leads to the observations that: 1) the increments of the transfer free energy during denaturation are of the same magnitude as the values of $\Delta\mu_{2, \text{tr}}$ of the native or denatured proteins; for example, at 52°C, 0.5 M trehalose, $\Delta\mu_{2, \text{tr}}$ has values of -0.73 and -0.20 kcal (mol protein)^{−1} (mol trehalose)^{−1} at pH 5.5 and 2.8, respectively, and a

stabilizing contribution ($\delta\Delta G^o$) of 1.0 ± 0.3 kcal mol^{−1}). 2) The corresponding enthalpic contributions, however, differ greatly. At the same condition, $\Delta \bar{H}_{2, \text{tr}}^N$ has a value of 82.3 kcal (mol protein)^{−1} (mol trehalose)^{−1} and ΔH^o of denaturation has values 85.3 and 105.6 kcal (mol protein)^{−1} (mol trehalose)^{−1} at pH 2.8 and 5.5 respectively. The enthalpy increment during denaturation, however, is very small, $\delta\Delta H^o = 0 \pm 3$ kcal mol^{−1}. This leads to the conclusions that: 1) The very small free energy values result from frequently large enthalpies that must be close to compensated by equally large entropic contributions, which for the transfer parameters are shown in Fig. 8. 2) The enthalpic contributions of interactions with the co-solvent system to the stabilization during denaturation are very small relative to the standard enthalpy change in the absence of the stabilizer, as well as to the absolute values of the transfer enthalpies. This means that, on denaturation, the interactions with the newly exposed structural elements of the protein must be similar in nature to those with the surface of the native protein. This is consistent with recent analysis by Liu and Bolen [57] of the interactions of osmolytes with amino acid side chains and peptide groups. These authors have shown that sucrose and sarcosine have a weak affinity for both polar and some non-polar residues, but a lower affinity than water for peptide groups. Since protein unfolding exposes to contact with solvent a large number of previously buried peptide groups, this report leads to the anticipation that, in the presence of osmolytes, preferential interaction should be less positive (lower binding or

higher exclusion) with unfolded protein than with its native form. This is exactly the observation made in the present study and permits the reduction in preferential binding on unfolding to be ascribed to the exposure of peptide groups previously not available to solvent molecules.

Acknowledgements

Communication No. 1805 from the Graduate Department of Biochemistry, Brandeis University, Waltham, MA 02254. This work was supported by GM14603.

References

- [1] J. Jarabak, J.A. Adams, H.C. Williams-Achmen and P. Talalay, *J. Biol. Chem.*, 237 (1962) 234.
- [2] S.L. Bradbury and W.B. Jakoby, *Proc. Natl. Acad. Sci. USA*, 69 (1972) 2373.
- [3] J.S. Meyers and W.B. Jakoby, *Biochem. Biophys. Res. Commun.*, 51 (1973) 631.
- [4] J.S. Meyers and W.B. Jakoby, *J. Biol. Chem.*, 250 (1975) 3785.
- [5] S.Y. Gerlsma, *J. Biol. Chem.*, 243 (1968) 957.
- [6] S.Y. Gerlsma, *Eur. J. Biochem.*, 14 (1970) 150.
- [7] S.Y. Gerlsma and E.R. Stuur, *Im. J. Peptide Protein Res.*, 4 (1972) 377.
- [8] S.Y. Gerlsma and E.R. Stuur, *Im. J. Peptide Protein Res.*, 6 (1974) 65.
- [9] S.N. Timasheff, J.C. Lee, E.P. Pittz and N. Tweedy, *J. Colloid Interface Sci.*, 55 (1976) 658.
- [10] T. Arakawa and S.N. Timasheff, *Biochemistry*, 21 (1982) 6536.
- [11] K.C. Aune and S.N. Timasheff, *Biochemistry*, 9 (1970) 1480.
- [12] J.C. Lee and S.N. Timasheff, *J. Biol. Biochem.*, 256 (1981) 7193.
- [13] K. Gekko and S.N. Timasheff, *Biochemistry*, 20 (1981) 4667.
- [14] K. Gekko and S.N. Timasheff, *Biochemistry*, 20 (1981) 4677.
- [15] T.-Y. Lin and S.N. Timasheff, *Protein Science*, 5 (1996) 372.
- [16] A.S. Sussman and B.T. Lingappa, *Nature (London)*, 130 (1959) 1343.
- [17] J.S. Clegg, *Comp. Biochem. Physiol.*, 14 (1965) 135.
- [18] C. Womersley, *Comp. Biochem. Physiol. B*, 70 (1981) 669.
- [19] J.H. Crowe, L.M. Crowe and D. Chapman, *Science*, 223 (1984) 701.
- [20] J.F. Carpenter, L.M. Crowe and J.H. Crowe, *Biochim. Biophys. Acta*, 923 (1987) 109.
- [21] T.E. Honadel and G.J. Killian, *Cryobiology*, 25 (1988) 331.
- [22] J.H. Crowe, J.F. Carpenter, L.M. Crowe and T.J. Anchordoguy, *Cryobiol.*, 27 (1990) 219.
- [23] C.W.B. Lee, J.S. Waugh and R.G. Griffin, *Biochemistry*, 25 (1986) 3737.
- [24] E.M. Arnett, N. Harvey, E.A. Johnson, D.S. Johnston, and D. Chapman, *Biochemistry*, 25 (1986) 5239.
- [25] G. Strauss and H. Hauser, *Proc. Natl. Acad. Sci. USA*, 83 (1986) 2422.
- [26] J.H. Crowe, B.J. Spargo and L.M. Crowe, *Proc. Natl. Acad. Sci. USA*, 84 (1987) 1537.
- [27] A.M. Crestfield, W.H. Stein and S. Moore, *Arch. Biochem. Biophys. Suppl.*, 1 (1962) 217.
- [28] A.M. Crestfield, W.H. Stein and S. Moore, *J. Biol. Chem.*, 236 (1963) 618.
- [29] J.C. Lee and S.N. Timasheff, *Biochemistry*, 13 (1974) 257.
- [30] J.C. Lee, K. Gekko and S.N. Timasheff, *Methods Enzymol.*, 61 (1979) 26.
- [31] T. Arakawa, R. Bhat and S.N. Timasheff, *Biochemistry*, 29 (1990) 1914.
- [32] W.W. Kielley and W.F. Harrington, *Biochim. Biophys. Acta*, 41 (1960) 401.
- [33] G. Scatchard, *J. Am. Chem. Soc.*, 68 (1946) 2315.
- [34] W.H. Stockmayer, *J. Chem. Phys.*, 18 (1950) 58.
- [35] G. Cohen and H. Eisenberg, *Biopolymers*, 6 (1968) 1077.
- [36] S.N. Timasheff and M.J. Kronman, *Arch. Biochem. Biophys.*, 83 (1959) 60.
- [37] E. Reisler, Y. Haik and H. Eisenberg, *Biochemistry*, 16 (1977) 197.
- [38] J.G. Kirkwood and R.J. Goldberg, *J. Chem. Phys.*, 18 (1950) 54.
- [39] E.F. Casassa and H. Eisenberg, *J. Phys. Chem.*, 65 (1961) 427.
- [40] E.F. Casassa and H. Eisenberg, *Adv. Protein Chem.*, 19 (1964) 287.
- [41] R. Biltonen and R. Lumry, *J. Am. Chem. Soc.*, 91 (1969) 4256.
- [42] S. Glasstone, *Thermodynamics for Chemists*, Van Nostrand, New York, 1947, pp. 292–295.
- [43] T. Kiefhaber, F.X. Schmid, M. Renner, J.-H. Hinz, U. Hahn and R. Quaas, *Biochemistry*, 29 (1990) 8250.
- [44] T.-Y. Lin and S.N. Timasheff, *Biochemistry*, 33 (1994) 12695.
- [45] S.N. Timasheff, *Annu. Rev. Biophys. Biomol. Struct.*, 22 (1993) 67.
- [46] H.B. Bull and K. Breese, *Arch. Biochem. Biophys.*, 128 (1968) 488.
- [47] I.D. Kuntz, *J. Am. Chem. Soc.*, 93 (1971) 514.
- [48] S.N. Timasheff and H. Inoue, *Biochemistry*, 7 (1968) 2501.
- [49] H. Inoue and S.N. Timasheff, *Biopolymers*, 11 (1972) 737.
- [50] D.W. Kupke, in S.J. Leach (Ed.), *Physical Principles and Techniques of Protein Chemistry, Part C*, Academic Press, New York, 1973, pp. 1–75.
- [51] J.A. Schellman, *Biophys. Chem.*, 37 (1990) 121.

- [52] J.A. Schellman, *Biophys. Chem.*, 45 (1993) 273.
- [53] S.N. Timasheff, *Biochemistry*, 31 (1992) 9857.
- [53a] K. Gekko and T. Morikawa, *J. Biochem.*, 90 (1981) 39.
- [54] J. Wyman, *Adv. Protein Chem.* 19 (1964) 223.
- [55] C. Tanford, *J. Mol. Biol.*, 39 (1969) 539.
- [56] J. Wyman and S.J. Gill, *Binding and Linkage. Functional Chemistry of Biological Macromolecules*, Univ. Science Books, Mill Valley, CA, 1990.
- [57] Y. Liu and D.S. Bolen, *Biochemistry*, 34 (1995) 12884.