

# Vesicular $\text{Ca}^{2+}$ -induced secretion promoted by intracellular pH-gradient disruption

Christy L. Haynes, Leah A. Buhler, R. Mark Wightman \*

*University of North Carolina, Chapel Hill, Department of Chemistry, Venable Hall, CB 3290, Chapel Hill, NC 27599, USA*

Received 6 February 2006; received in revised form 12 April 2006; accepted 12 April 2006

Available online 26 April 2006

## Abstract

The actions of the protonophore CCCP on intracellular  $\text{Ca}^{2+}$  regulation and exocytosis in chromaffin cells have been examined. Simultaneous fura-2 imaging and amperometry reveal that exposure to CCCP not only perturbs mitochondrial function but that it also alters vesicular storage of  $\text{Ca}^{2+}$  and catecholamines. By disrupting the pH gradient of the secretory vesicle membrane, the protonophore allows both  $\text{Ca}^{2+}$  and catecholamine to leak into the cytosol. Unlike the high cytosolic  $\text{Ca}^{2+}$  concentrations resulting from mitochondrial membrane disruption,  $\text{Ca}^{2+}$  leakage from secretory vesicles may initiate exocytotic release. In conjunction with previous studies, this work reveals that catalytic and self-sustained vesicular  $\text{Ca}^{2+}$ -induced exocytosis occurs with extended exposure to weak acid or base protonophores.

© 2006 Published by Elsevier B.V.

**Keywords:** Chromaffin cell; Amperometry; Fura-2; CCCP;  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release

## 1. Introduction

Elevated intracellular  $\text{Ca}^{2+}$  triggers exocytosis from secretory vesicles. The primary route for its elevation is entry of external  $\text{Ca}^{2+}$  via voltage-gated  $\text{Ca}^{2+}$  channels [1]. However, intracellular stores of  $\text{Ca}^{2+}$  can also promote exocytosis [2–5]. The endoplasmic reticulum is often designated as the key dynamic  $\text{Ca}^{2+}$  reservoir within cells, participating in  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release and using  $\text{Ca}^{2+}$ -regulated enzymes [6–8]. Another source of intracellular  $\text{Ca}^{2+}$  is the secretory vesicles where  $\text{Ca}^{2+}$  is sequestered into the acidic compartment and stored at high concentration by virtue of an association with chromogranin A [9,10]. In fact, nearly 60% of  $\text{Ca}^{2+}$  in chromaffin cells resides in the vesicular pools [11] although >99% of the vesicular  $\text{Ca}^{2+}$  is highly associated with the other vesicular contents [12]. Weak bases such as methylamine and methamphetamine, as well as VMAT inhibitors, disrupt vesicular storage, promoting  $\text{Ca}^{2+}$  leakage from the vesicles and evoking exocytosis [13,14].  $\text{Ca}^{2+}$  release from vesicles in many cell types such as insulin-secreting cells, buccal ganglion neurons, caudate nucleus synaptosomes, and neurohypophysial

nerve terminals has been demonstrated [3–5]. While high affinity  $\text{Ca}^{2+}$  probes localized within vesicles of PC-12 cells revealed that the vesicular compartment tracks cytosolic  $\text{Ca}^{2+}$  levels [15], other vesicular compartments are known to have  $\text{Ca}^{2+}$  concentrations that are highly sensitive to the vesicular pH. This vesicular  $\text{Ca}^{2+}$  source is particularly intriguing because it suggests that vesicles can directly induce their own release [16].

A third intracellular store of  $\text{Ca}^{2+}$  is mitochondria. Mitochondria accumulate  $\text{Ca}^{2+}$  due to a pH gradient between the inner mitochondrial membrane (pH=6.8) and the mitochondrial matrix (pH=8) [17]. Several investigators have shown that mitochondria are the chief organelle that restores elevated levels of intracellular  $\text{Ca}^{2+}$  to the resting concentration [18–21]. A common strategy to probe mitochondrial function is to use uncouplers, substances that dissipate the pH gradient, such as carbonyl cyanide *m*-chlorophenylhydrazone (CCCP). In this work, we examine the actions of the protonophore CCCP on intracellular  $\text{Ca}^{2+}$  regulation and exocytosis in chromaffin cells. CCCP has one extractable proton (pKa ~6) [22] and is able to cross lipophilic membranes in both the protonated and deprotonated forms. CCCP acts as a proton shuttle by bringing protons from the cytosol to the more basic mitochondrial matrix, thus collapsing the pH gradient that acts as the driving force for

\* Corresponding author. Tel.: +1 919 843 8164; fax: +1 919 962 2388.

E-mail address: [rmw@email.unc.edu](mailto:rmw@email.unc.edu) (R.M. Wightman).

mitochondrial  $\text{Ca}^{2+}$  uptake. While CCCP can be used to investigate the role of mitochondrial  $\text{Ca}^{2+}$  regulation, recent evidence indicates that structurally similar compounds can also cause  $\text{Ca}^{2+}$  leakage from vesicles in PC-12 cells [12]. Since vesicular  $\text{Ca}^{2+}$  leakage is often accompanied by dissipation of the proton gradient and self-catalyzed exocytosis [13,14], we felt it was imperative to evaluate whether CCCP could evoke exocytosis as well.

## 2. Methods

Acutely dissociated bovine adrenal medullary chromaffin cells were prepared as previously described [23]. The cells were loaded with the cytosolic  $\text{Ca}^{2+}$ -chelating dye, fura-2 before simultaneously recording fluorescence and release with amperometry at carbon-fiber microelectrodes [23]. The cells were exposed to 2  $\mu\text{M}$  CCCP either by transient pressure ejection for 3 s or incubation for 5 or 15 min. Except where noted, cells were maintained in a standard buffer solution containing 10 mM Tris-HCl, 150 mM NaCl, 5 mM KCl, 2 mM  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 1.2 mM  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , and 5 mM glucose, adjusted to pH 7.4. Carbon-fiber disk microelectrodes were beveled at  $45^\circ$ . Amperometry employed an Axopatch 200A potentiostat with an applied voltage of +650 mV versus a Ag/AgCl reference electrode. Release was stimulated with 3 s applications of 60 mM  $\text{K}^+$ . Cytosolic catecholamine measurements were performed in  $\text{Ca}^{2+}$ -free buffer using transient applications of 10  $\mu\text{M}$  digitonin [14].

## 3. Results and discussion

### 3.1. Transient CCCP exposure can evoke cytosolic $\text{Ca}^{2+}$ and exocytosis

To examine the direct, short-term effects of the protonophore, CCCP was pressure ejected onto individual chromaffin cells in  $\text{Ca}^{2+}$ -free buffer. As shown for a single example in Fig. 1, a 3-s application of CCCP evoked elevated cytosolic  $\text{Ca}^{2+}$ , measured via fura-2 fluorescence, in all cases. One source of this  $\text{Ca}^{2+}$  is undoubtedly intracellular mitochondria where CCCP uncouples the proton gradient that controls mitochondrial stores of  $\text{Ca}^{2+}$ . Recent results have shown that compounds similar to CCCP can also displace intravesicular  $\text{Ca}^{2+}$ , a process that appears to be controlled by a  $\text{H}^+/\text{Ca}^{2+}$  exchanger [12]. Thus, raising the vesicular pH via CCCP would generate conditions where the chromogranin A matrix could begin to unravel, allowing vesicular  $\text{Ca}^{2+}$  to escape into the cytosol via the exchanger. The elevated cytosolic  $\text{Ca}^{2+}$  concentration was sufficient to cause vesicular release, measured by amperometry, in two out of ten cells tested in this way (bottom amperometric trace in Fig. 1).

### 3.2. Long-term CCCP exposure increases cytosolic $\text{Ca}^{2+}$ clearance and release duration

When examined following a 5-min incubation in CCCP (2  $\mu\text{M}$ ) in buffer containing  $\text{Ca}^{2+}$ , levels of intracellular  $\text{Ca}^{2+}$




Fig. 1. Amperometric and fura-2 signal traces in response to a transient application of CCCP. A 3-s application of 2  $\mu\text{M}$  CCCP caused  $\text{Ca}^{2+}$  elevation (top). In 8 out of 10 cells, no release was observed (middle). In 2 out of 10 cells, the elevated  $\text{Ca}^{2+}$  concentration was sufficient to evoke amperometric release (bottom). The left y-axis corresponds to the upper trace while the scale on the right corresponds to both amperometric traces. Cells were maintained in  $\text{Ca}^{2+}$ -free buffer.

were returned to levels similar to control cells. However, spontaneous exocytotic events were occasionally observed during the CCCP incubation period (data not shown). These spontaneous events occurred at a lower frequency than stimulated events and were more common at the beginning of the incubation period than the end. More dramatic effects were seen on  $\text{Ca}^{2+}$  clearance following membrane depolarization. Incubation in CCCP increased both the cytosolic  $\text{Ca}^{2+}$  clearance time and the catecholamine release duration evoked by transient  $\text{K}^+$ , consistent with previous results [18,21]. Specifically, responses to pressure ejection (3 s) of 60 mM  $\text{K}^+$  at cells incubated for 5 min in CCCP (2  $\mu\text{M}$ ,  $n=6$ ) were compared to control cells ( $n=6$ ) stimulated in the same way (Fig. 2). The cytosolic  $\text{Ca}^{2+}$  clearance in the cells exposed to CCCP had a half-width of  $20.3 \pm 2.3$  s. This was almost twice as long as the  $\text{Ca}^{2+}$  clearance in the control cells, with a fura-2 half-width of  $9.4 \pm 1.2$  s (Fig. 2). Similarly, the  $21.9 \pm 1.9$  s duration of vesicular release from cells maintained in buffer containing CCCP was more than twice the  $5.6 \pm 0.7$  s duration observed in control cells. The increase in both the calcium clearance time and catecholamine release duration was significantly different from the control values (Mann-Whitney  $U$ -test,  $p < 0.01$ ). Thus, during  $\text{K}^+$  evoked release, we largely attribute the effect of CCCP on elevated  $\text{Ca}^{2+}$  and release duration to its effects on mitochondria, where its uncoupling impairs the ability to restore cytosolic  $\text{Ca}^{2+}$  levels. This conclusion is consistent with that of Yang et al. who reported that the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, endoplasmic reticulum, and mitochondria work cooperatively to regulate  $\text{Ca}^{2+}$  near exocytotic sites but that the mitochondrial uptake is dominant when the cytosolic  $\text{Ca}^{2+}$  diffuses away from exocytotic sites [18].




Fig. 2. Amperometric and fura-2 signal traces in the presence and absence of the protonophore, CCCP. Each cell was exposed to a 3-s pressure ejection of 60 mM  $K^+$  in  $Ca^{2+}$  containing buffer. Overlays of representative amperometric (right y-axis) and fura-2 (left y-axis) traces for (A) untreated cells and (B) treated cells. Exposure to CCCP increased both the release duration and the width at half-maximal height of the fura-2 signal.

### 3.3. Quantal size decreases with long protonophore exposure time

The role of CCCP in secretory vesicle leakage was examined by monitoring the quantal size ( $Q$ ) of exocytosed catecholamines with different CCCP exposure times. Each cell was stimulated 4 times by pressure ejection of high  $K^+$  with 2-min intervals between each stimulation, and the resulting exocytotic events were recorded. Next, the cell was incubated in CCCP (2  $\mu$ M) for 5 or 15 min. After washing with CCCP-free buffer, the same stimulation pattern with high  $K^+$  was applied again. To account for the variability that can exist between cells, each cell was used as its own control by normalizing amperometric spikes obtained post-incubation with the spikes obtained prior to the CCCP incubation. In control cells ( $n=10$ ), neither 5 nor 15 min incubations in CCCP-free buffer caused a significant change in spike half-width ( $t_{1/2}$ ) or  $Q$  as indicated by post/pre ratios near 1.0 for all cases (Fig. 3B and C, black bars). Ratios calculated from amperometric traces recorded before and after a 5 min incubation in CCCP ( $n=4$ ) also did not yield significantly different  $t_{1/2}$  compared to control cells (Fig. 3B, white bars). The  $Q$  ratios were slightly lower, but the result was not significant ( $p>0.05$ ). In contrast,  $Q$  ratios calculated for cells that were incubated with CCCP for 15 min ( $n=5$ ) are significantly smaller (0.51 post/pre-incubation) than control cell  $Q$  ratios (Fig. 3C, white bars) while  $t_{1/2}$  ratios remain unaffected. Because there was not a significant difference between the  $t_{1/2}$  ratios obtained following incubation with CCCP, it is clear that the lowered amounts released per exocytotic event are not due to an increased affinity between catecholamine and chromogranin A.




Fig. 3. Amperometric detection of time-dependent depletion of epinephrine from vesicles. Individual chromaffin cells were exposed 4 times to a 3-s pressure ejection of 60 mM  $K^+$  in a  $Ca^{2+}$  containing buffer. The cells were then incubated in buffer containing 2  $\mu$ M CCCP for the indicated time. After the incubation period, the cells were washed with buffer and stimulated again using the same  $K^+$  ejection pattern. Average spike parameters were then normalized by the pre-incubation values for each cell so that each cell acted as its own control. (A) A representative amperometric trace of a 15-min CCCP incubation. (B) 5-min incubation in CCCP caused no significant change in the average current spike half-width or quantal size compared with pre-incubation averages. (C) 15-min incubation causes a significant decrease in quantal size ( $p<0.001$  by Mann-Whitney  $U$ -test).




Fig. 4. Cytosolic catecholamine content evaluated by digitonin permeabilization. The cells were incubated with 2  $\mu$ M CCCP in  $\text{Ca}^{2+}$ -containing buffer for either (A) 5 min or (B) 15 min. After washing and transferring to  $\text{Ca}^{2+}$ -free buffer, 10  $\mu$ M digitonin was pressure ejected onto the cell. The resulting envelope is an indication of the amount of catecholamine that was present in the cytosol of the cell. (C) The 15-min incubation resulted in approximately double the amount of catecholamine molecules than were detected after a 5-min incubation ( $p < 0.05$  by Mann-Whitney  $U$ -test).

Rather, the results are consistent with a disruption of vesicular storage following incubation with CCCP as a consequence of collapsing intracellular pH gradients, allowing catecholamine molecules and  $\text{Ca}^{2+}$  leakage. This would result in the reduced amount released in each exocytotic event, as observed.

#### 3.4. Cytosolic catecholamine increases dramatically with long CCCP exposure time

Because both catecholamine and  $\text{Ca}^{2+}$  are highly associated within the acidic intravesicular matrix, leakage of catecholamine into the cytosol during CCCP incubation is consistent with the simultaneous release of intravesicular  $\text{Ca}^{2+}$  reported previously [12]. This leakage was verified by examining cytosolic catecholamine molecules after permeabilizing CCCP-exposed cells with 10  $\mu$ M digitonin in  $\text{Ca}^{2+}$ -free buffer. Unlike the millisecond timescale of exocytotic release, cytosolic catecholamine liberated following digitonin permeation has a time course of several seconds (Fig. 4). In control cells where CCCP was not included in the buffer, no catecholamine was detected when the cells were exposed to digitonin (data not shown), indicating low cytoplasmic catecholamine levels. However, digitonin permeabilization after 5-min CCCP (2  $\mu$ M) incubation resulted in detection of  $2.1 \pm 0.5 \times 10^7$  cytoplasmic catecholamine molecules ( $n=7$ ) and  $5.5 \pm 1.0 \times 10^7$  cytoplasmic catecholamine molecules were detected from cells incubated for 15 min with the same concentration of CCCP ( $n=6$ ). These cytosolic catecholamine amounts are comparable to those measured when chromaffin cells were incubated with the VMAT inhibitor, reserpine [14]. Thus, the cytosolic catecholamine measurements clearly illustrate that the intravesicular contents can be displaced into the cytosol following CCCP. This extensive vesicular perturbation after 15 min CCCP incubation is manifested by

the decreased amounts released in each exocytotic event (Fig. 3C).

#### 4. Conclusions

The protonophore CCCP has been a critical tool in establishing the predominance of mitochondria in restoring high cytosolic  $\text{Ca}^{2+}$  to resting levels [20,21,24]. Simultaneous fura-2 imaging and amperometry used in this work reveal that incubations in CCCP also alter vesicular storage of  $\text{Ca}^{2+}$  and catecholamine. These data clearly indicate that CCCP promotes leakage of vesicle contents into the cytosol. Indeed, the magnitude of the change in intracellular  $\text{Ca}^{2+}$  concentration following transient exposure to CCCP is similar to when the cells were depolarized (compare Figs. 1 and 2). Despite evoking large cytosolic  $\text{Ca}^{2+}$  concentrations, exposure to CCCP did not always cause secretion, and it did so primarily upon the initial exposure. Elevated intracellular  $\text{Ca}^{2+}$  can only cause exocytosis if it occurs in close proximity to both the cell membrane and a fusion competent vesicle. Because CCCP disrupts pH gradients, and thus  $\text{Ca}^{2+}$  storage, at multiple intracellular organelles, most of which are not proximal to the cell membrane, overall intracellular  $\text{Ca}^{2+}$  rises but only a small portion of it is optimally positioned to induce exocytosis. Based on this work, previous results achieved with amine weak bases [14] can be further generalized to state that the weak acids as well as weak base protonophores can induce unraveling of vesicular contents. The results also show that the vesicular  $\text{Ca}^{2+}$  likely catalyzes and promotes self-sustained exocytosis.

#### Acknowledgements

This work was supported by National Institutes of Health Grant NS 38879.

#### References

- [1] R.W. Tsien, Calcium channels in excitable cell membranes, *Annu. Rev. Physiol.* 45 (1983) 341–358.
- [2] S.H. Yoo, J.P. Albanesi,  $\text{Ca}^{2+}$ -induced conformational change and aggregation of chromogranin A, *J. Biol. Chem.* 265 (1990) 14414–14421.
- [3] J.-D. Troade, S. Thirion, J.-P. Laugier, G. Nicaise, Calcium-induced calcium increase in secretory vesicles of permeabilized rat neurohypophyseal nerve terminals, *Biol. Cell* 90 (1998) 339–349.
- [4] W.J.J.M. Scheenen, C.B. Wollheim, T. Pozzan, C. Fasolato,  $\text{Ca}^{2+}$  depletion from granules inhibits exocytosis: a study with insulin-secreting cells, *J. Biol. Chem.* 273 (1998) 19002–19008.
- [5] P. Fossier, et al., Control of the calcium concentration involved in acetylcholine release and its facilitation: an additional role for synaptic vesicles? *Neuroscience* 85 (1998) 85–91.
- [6] M.F. Rossier, J.W. Putney Jr., The identity of the calcium-storing, inositol 1,4,5-trisphosphate-sensitive organelle in non-muscle cells: calciosome, endoplasmic reticulum... or both? *Trends Neurosci.* 14 (1991) 310–314.
- [7] A. Verkhratsky, E.C. Toescu, Endoplasmic reticulum  $\text{Ca}^{2+}$  homeostasis and neuronal death, *J. Cell. Mol. Med.* 7 (2003) 351–361.
- [8] G. Kang, G.G. Holz, Amplification of exocytosis by  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release in INS-1 pancreatic  $\beta$  cells, *J. Phys.* 546 (2003) 175–189.
- [9] J.S. Videen, M.S. Mezger, Y.M. Chang, D.T. O'Connor, Calcium and catecholamine interactions with adrenal chromogranins. Comparison of driving forces in binding and aggregation, *J. Biol. Chem.* 267 (1992) 3066–3073.

- [10] S.H. Yoo, M.S. Lewis, Effects of pH and  $\text{Ca}^{2+}$  on monomer-dimer and monomer-tetramer equilibria of chromogranin A, *J. Biol. Chem.* 267 (1992) 11236–11241.
- [11] J.R. Haigh, R. Parris, J.H. Phillips, Free concentrations of sodium, potassium, and calcium in chromaffin granules, *Biochem. J.* 259 (1989) 485–491.
- [12] N.R. Mahapatra, et al., A dynamic pool of calcium in catecholamine storage vesicles, *J. Biol. Chem.* 279 (2004) 51107–51121.
- [13] M.L. Mundorf, S.E. Hochstetler, R.M. Wightman, Amine weak bases disrupt vesicular storage and promote exocytosis in chromaffin cells, *J. Neurochem.* 73 (1999) 2397–2405.
- [14] M.L. Mundorf, K.P. Troyer, S.E. Hochstetler, J.A. Near, R.M. Wightman, Vesicular  $\text{Ca}^{2+}$  participates in the catalysis of exocytosis, *J. Biol. Chem.* 275 (2000) 9136–9142.
- [15] A. Moreno, et al., Calcium dynamics in catecholamine-containing secretory vesicles cell, *Cal* 37 (2005) 555–564.
- [16] K.J. Mitchell, et al., Dense core secretory vesicles revealed as a dynamic  $\text{Ca}^{2+}$  store in neuroendocrine cells with a vesicle-associated membrane protein aequorin chimera, *J. Cell Biol.* 155 (2001) 45–51.
- [17] M.F. Cano Abad, G. Di Benedetto, P.J. Magalhaes, L. Filippin, T. Pozzan, Mitochondrial pH monitored by a new engineered green fluorescent protein mutant, *J. Biol. Chem.* 279 (2004) 11521–11529.
- [18] D.-M. Yang, L.-S. Kao, Relative contribution of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, mitochondria and endoplasmic reticulum in the regulation of cytosolic  $\text{Ca}^{2+}$  and catecholamine secretion of bovine adrenal chromaffin cells, *J. Neurochem.* 76 (2001) 210–216.
- [19] J.D. Johnson, J.P. Chang, Calcium buffering activity of mitochondria controls basal growth hormone secretion and modulates specific neuropeptide signaling cell, *Cal* 37 (2005) 573–581.
- [20] T. Xu, M. Naraghi, H. Kang, E. Neher, Kinetic studies of  $\text{Ca}^{2+}$  binding and  $\text{Ca}^{2+}$  clearance in the cytosol of adrenal chromaffin cells, *Biophys. J.* 73 (1997) 532–545.
- [21] J. Herrington, Y.B. Park, D.F. Babcock, B. Hille, Dominant role of mitochondria in clearance of large  $\text{Ca}^{2+}$  loads from rat adrenal chromaffin cells, *Neuron* 16 (1996) 219–228.
- [22] E. Sturdik, et al., Relationships among structure, reactivity towards thiols, and basicity of phenylhydrazonopropanedinitriles collection of Czechoslovak, *Chem. Commun.* 50 (1985) 2065–2076.
- [23] J.M. Finnegan, R. Borges, R.M. Wightman, Comparison of cytosolic  $\text{Ca}^{2+}$  and exocytosis responses from single rat and bovine chromaffin cells, *Neuroscience* 71 (1996) 833–843.
- [24] Y.B. Park, J. Herrington, D.F. Babcock, B. Hille,  $\text{Ca}^{2+}$  clearance mechanisms in isolated rat adrenal chromaffin cells, *J. Phys.* 492 (1996) 329–346.