



Structural effects in vitro of the anti-inflammatory drug diclofenac on human erythrocytes and molecular models of cell membranes

Mario Suwalsky^{a,*}, Marcela Manrique^{a,b}, Fernando Villena^c, Carlos P. Sotomayor^d

^a Faculty of Chemical Sciences, University of Concepción, Concepción, Chile

^b University of Antioquia, Faculty of Exact and Natural Sciences, Medellín, Colombia

^c Faculty of Biological Sciences, University of Concepción, Concepción, Chile

^d Institute of Chemistry, Catholic University of Valparaíso, Valparaíso, Chile

ARTICLE INFO

Article history:

Received 12 November 2008

Received in revised form 18 December 2008

Accepted 19 December 2008

Available online 31 December 2008

Keywords:

Diclofenac

Nonsteroidal anti-inflammatory drugs

Human erythrocytes

Cell membrane

Lipid bilayer

ABSTRACT

Diclofenac, a nonsteroidal anti-inflammatory drug (NSAID), has been widely investigated in terms of its pharmacological action, but less is known about its effects on cell membranes and particularly on those of human erythrocytes. In the present work, the structural effects on the human erythrocyte membrane and molecular models have been investigated and reported. This report presents the following evidence that diclofenac interacts with red cell membranes: a) X-ray diffraction and fluorescence spectroscopy of phospholipid bilayers showed that diclofenac interacted with a class of lipids found in the outer moiety of the erythrocyte membrane; b) in isolated unsealed human erythrocyte membranes (IUM) the drug induced a disordering effect on the acyl chains of the membrane lipid bilayer; c) in scanning electron microscopy (SEM) studies on human erythrocytes it was observed that the drug induced changes different from the normal biconcave morphology of most red blood cells. This is the first time in which structural effects of diclofenac on the human erythrocyte membrane have been described.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Nonsteroidal anti-inflammatory drugs (NSAIDs), widely used for anti-inflammatory, antipyretic and analgesic purposes, are one of the most frequently used kinds of medicines, accounting for nearly 5% of all prescribed medications [1]. Their therapeutic effects result mostly from the inhibition of cyclooxygenase (COX), an enzyme involved in the production of prostaglandins which have a strong propensity for inducing inflammation [2,3]. NSAIDs are a chemically heterogeneous group of compounds, often chemically unrelated, which nevertheless share certain therapeutic actions and adverse effects. Diclofenac, (2-[(2,6-dichlorophenyl)amino]benzeneacetic acid, Fig. 1) is a non-steroidal anti-inflammatory drug with good analgesic properties; however, it may cause side effects including gastrointestinal disorders when administered by oral route and cutaneous lesions by intramuscular injection [4]. It also has cytotoxic effects and induces apoptosis in various cultured cell lines. Both toxic and apoptotic effects of diclofenac might be involved in drug-induced Reye's syndrome, renal toxicity, hepatotoxi-

city, and pancytopenia [5]. Its effects on patient blood are significantly associated with aplastic and hemolytic anemia, thrombocytopenia and agranulocytosis [6]. The primary target of NSAIDs in relation to their direct cytotoxicity remains unknown [2]. One such target candidate is the cell membrane. It has been reported that NSAIDs interact with phospholipids, and that phospholipid liposomes reduce the direct cytotoxicity of in vivo NSAIDs [7–9]. On the other hand, it has been suggested that mitochondrial membrane permeability transition is a possible mechanism of diclofenac-induced apoptosis in hepatocytes [5]. It has been found that diclofenac induced membrane permeability transition in isolated rat liver mitochondria and in primary cultured hepatocytes [10]. The cell membrane is a diffusion barrier that protects the interior of the cell, and thus, its structure and functions are susceptible to alterations as a consequence of interactions with chemical species.

In the course of in vitro systems search for the toxicity screening of biologically relevant chemicals, different cellular models have been applied in order to examine their adverse effects. Intended to better understand the molecular mechanisms of diclofenac interaction with cell membranes, we have utilized human erythrocytes and molecular models of cell membranes. Human erythrocytes were chosen because since they have only one membrane and no internal organelles it is an ideal cell system for studying basic drug–biomembrane interactions [11]. Additionally, although less specialized than many other cell membranes, they carry on enough similar functions, such as active and passive transport and the production of ionic and electric gradients,

Abbreviations: NSAIDs, nonsteroidal anti-inflammatory drugs; DMPC, dimyristoylphosphatidylcholine; DMPE, dimyristoylphosphatidylethanolamine; SEM, scanning electron microscopy; LUV, large unilamellar vesicles; r, fluorescence anisotropy; GP, generalized polarization; DPH, 1,6-diphenyl-1,3,5-hexatriene; laurdan, 6-dodecanoyl-2-dimethylaminonaphthalene; IUM, isolated unsealed human erythrocyte membrane; RBCS, red blood cells suspension.

* Corresponding author. Tel.: +56 41 2204171; fax: +56 41 2245974.

E-mail address: msuwalsk@udec.cl (M. Suwalsky).




Fig. 1. Structural formula of diclofenac.

in order to be considered representative of the plasma membrane in general. The molecular models consisted of bilayers built-up of dimyristoylphosphatidylcholine (DMPC) and dimyristoylphosphatidylethanolamine (DMPE), where DMPC represents phospholipid classes located in the outer monolayer of cell membranes, particularly of the human erythrocyte, whereas DMPE represents those preferentially sited in the inner monolayers [12,13]. The diclofenac capacity of perturbing the multilayer structures of DMPC and DMPE was evaluated by X-ray diffraction. Large unilamellar DMPC vesicles (LUV) were studied by fluorescence spectroscopy, and human erythrocytes were observed by means of scanning electron microscopy. These systems and techniques have been used in our laboratories in order to determine the interaction with and the membrane-perturbing effects of other therapeutic drugs [14–16].

2. Materials and methods

2.1. X-ray diffraction studies of DMPC and DMPE multilayers

The capacity of diclofenac to perturb the structures of DMPC and DMPE multilayers was evaluated by X-ray diffraction. Synthetic DMPC (lot. 14OPC-241, MW 677.9), DMPE (lot. 14OPE-58, MW 635.9), and sodium diclofenac (lot. 0751896, MW 318.1) from Sigma (St. Louis, MO) were used without further purification. About 2 mg of each phospholipid was mixed in Eppendorf tubes with 200 μ l of (a) distilled water and (b) aqueous solutions of diclofenac in a range of concentrations (0.01 mM to 2.0 mM).

The specimens were incubated for 30 min at 37 °C and 60 °C with DMPC and DMPE respectively, and centrifuged for 10 min at 2500 rpm. The samples were then transferred to 1.5 mm dia special glass capillaries (Glas-Technik & Konstruktion, Berlin, Germany) and X-ray diffracted. Specimen-to-film distances were 8 and 14 cm, standardized by sprinkling calcite powder on the capillary surface. Ni-filtered CuK α radiation from a Bruker Kristalloflex 760 (Karlsruhe, Germany) X-ray generator was used. The relative reflection intensities were obtained in an MBraun PSD-50 M linear position-sensitive detector system (Garching, Germany) and no correction factors were applied. The experiments were performed at 18 ± 1 °C, which is below the main phase transition temperature of both DMPC (24.3 °C) and DMPE (50.2 °C) [17,18]. Higher temperatures would have induced transitions to more fluid phases making the detection of structural changes harder. Each experiment was performed in triplicate and in case of doubts, additional experiments were carried out.

2.2. Fluorescence measurements of isolated unsealed human erythrocyte membranes (IUM) and large unilamellar vesicles (LUV)

The influence of diclofenac on the physical properties of IUM and DMPC LUV was examined by fluorescence spectroscopy using DPH and laurdan (Molecular Probe, Eugene, OR, USA) fluorescent probes. DPH is widely used as a probe for the hydrophobic regions of the phospholipid bilayers because of its favorable spectral properties. Their steady-state fluorescence anisotropy measurements were used to investigate the structural properties of IUM and DMPC LUV as it provides a measure of the rotational diffusion of the fluorophor, restricted within a certain region such as a cone due to the lipid acyl chain packing order. Laurdan, an amphiphilic probe, has high excitation sensitivity and emission spectra to the physical state of membranes. With the fluorescent moiety within a shallow position in the bilayer, laurdan provides information about the polarity and/or molecular dynamics at the level of the phospholipid glycerol backbone. The quantification of the laurdan fluorescence spectral shift was effected by means of the general polarization (GP) concept [19]. Erythrocytes were separated from heparinized venous blood samples obtained from normal casual donors by centrifugation and washing procedures. IUM were prepared by lysis, according to Dodge et al. [20].




Fig. 2. Microdensitograms from X-ray diffraction patterns of (A) dimyristoylphosphatidylcholine (DMPC) and (B) dimyristoylphosphatidylethanolamine (DMPE) in water and aqueous solutions of sodium diclofenac; (a) low-angle and (b) wide-angle reflections.




Fig. 3. Effects of sodium diclofenac on the anisotropy (r) of DPH and on the general polarization (GP) of lauridan embedded in isolated unsealed human erythrocyte membranes (IUM) at 37 °C.

DMPC LUV suspended in water were prepared by extrusion of frozen and thawed multilamellar liposome suspensions (final lipid concentration 0.4 mM) through two stacked polycarbonate filters of 400 nm pore size (Nucleopore, Corning Costar Corp., MA, USA) under nitrogen pressure at 10 °C above the lipid phase transition temperature. DPH and lauridan were incorporated into IUM and LUV by addition of 2 μ L aliquots of 0.5 mM solutions of the probe in dimethylformamide and ethanol, respectively, in order to obtain final analytical concentrations of 1×10^{-3} mM, and incubated them at 37 °C for 45 min. Fluorescence spectra and anisotropy measurements were performed in a phase shift and modulation K₂ steady-state and time resolved spectrofluorometer (ISS, Inc., Champaign, IL, USA) interfaced to computer. Software from ISS was used for both data collection and analysis. LUV suspensions measurements were carried out at 18 °C and 37 °C, and IUM measurements were made at 37 °C using 10 mm path-length square quartz cuvettes. Sample temperature was controlled by an external bath circulator (Cole-Parmer, Chicago, IL, USA)

and monitored before and after each measurement using an Omega digital thermometer (Omega Engineering, Inc., Stamford, CT, USA). Anisotropy measurements were made in the L configuration using Glan Thompson prism polarizers (I.S.S., Inc.) in both exciting and emitting beams. The emission was measured using a WG-420 Schott high-pass filter (Schott WG-420, Mainz, Germany) with negligible fluorescence. DPH fluorescence anisotropy (r) was calculated according to the definition: $r = (I_{\parallel} - I_{\perp}) / (I_{\parallel} + 2I_{\perp})$, where I_{\parallel} and I_{\perp} are the corresponding parallel and perpendicular emission fluorescence intensities with respect to the vertically polarized excitation light [21]. Lauridan fluorescence spectral shifts were quantitatively evaluated using the GP concept (see above) which is defined by the expression $GP = (I_b - I_r) / (I_b + I_r)$, where I_b and I_r are the emission intensities at the blue and red edges of the emission spectrum, respectively. These intensities have been measured at the emission wavelengths of 440 and 490 nm, which correspond to the emission maxima of lauridan in both gel and liquid crystalline phases, respectively [22]. Diclofenac was




Fig. 4. Effects of sodium diclofenac on the generalized polarization (GP) of lauridan embedded in DMPC large unilamellar vesicles (LUV) at 18 and 37 °C.

incorporated in IUM and LUV suspensions by addition of adequate (10 mM) aliquots of a concentrated diclofenac in order to obtain the different concentrations used in this work. Samples thus prepared were then incubated at 37 °C, for ca. 15 min and measured at 18 and 37 °C; at 18 °C because the X-ray experiments were performed at about this temperature, and at 37 °C because that is the normal temperature at which erythrocytes circulate in humans. Blank subtraction was performed in all measurements using unlabeled samples without probes. Data presented in Figs. 3–5 represent mean values and standard error of ten measurements in two independent samples. Unpaired Student's *t*-test was used for statistical calculations.

2.3. Scanning Electron Microscope (SEM) studies of human erythrocytes

Blood was obtained from healthy human male donor not receiving any pharmacological treatment. Blood samples (0.1 ml) were obtained by puncture of the ear lobule and received in an Eppendorff tube containing 10 µl of heparin (5000 UI/ml) in 0.9 ml of saline solution (NaCl 0.9%, pH 7.4). The sample was centrifuged (1000 rpm × 10 min) and the supernatant was discarded and replaced by the same volume of saline solution; the whole process was repeated three times. The sedimentary red blood cells were suspended in 0.9 mL of saline solution and fractions of this stock of red blood cells suspension (RBCS) and saline solution were placed in Eppendorf tubes to prepare (a) the control, by mixing 100 µl of saline solution plus 100 µl RBCS, and (b) a range of concentrations of diclofenac (0.01 mM–2 mM) by mixing 100 µl of RBCS with 100 µl of adequate diclofenac stock solutions in saline solution. All the samples were then incubated for 1 h at 37 °C. After the incubation, samples were centrifuged (1000 rpm × 10 min) and the supernatant was discarded. Then, they were fixed overnight at 4 °C by adding 500 µl of 2.5% glutaraldehyde to each one. The fixed samples were washed with distilled water, placed over Al glass cover stubs, air dried at 37 °C for 30 min to 1 h, and gold-coated for 3 min at 10⁻¹ Torr in a sputter device (Edwards S150, Sussex, England). Resulting specimens were examined in a Jeol SEM (JSM 6380 LB, Japan).

3. Results

3.1. X-ray diffraction studies of DMPC and DMPE multilayers

Fig. 2A exhibits the results obtained by incubating DMPC with water and diclofenac. As expected, water altered the DMPC structure: its

bilayer repeat (bilayer width plus the width of the water layer between bilayers) increased from about 55 Å in its dry crystalline form [23] to 64.5 Å when immersed in water) and its low-angle reflections, which correspond to DMPC polar terminal groups, were reduced to only the first two orders of the bilayer repeat. On the other hand, only one strong reflection of 4.2 Å showed up in the wide-angle region, which corresponds to the average distance between fully extended acyl chains organized with rotational disorder in hexagonal packing. These results were indicative of the fluid state reached by DMPC bilayers. Fig. 2A discloses that after exposure to 0.1 mM diclofenac there was a weakening of the low- and wide-angle lipid reflection intensities (indicated as (a) and (b) in the figure, respectively); addition of diclofenac in increasing concentrations caused a monotonically decrease in the phospholipid reflection intensities, until they practically disappeared at 2 mM. From these results, it can be concluded that diclofenac produced a significant structural perturbation of DMPC bilayers. Fig. 2B shows the results of the X-ray diffraction analysis of DMPE bilayers incubated with water and diclofenac. As reported elsewhere, water did not significantly affect the bilayer structure of DMPE [23]. Fig. 2B also shows that increasing concentrations of diclofenac did not cause a weakening in DMPE reflection intensities, all of which still remained practically unaffected at the highest assayed diclofenac concentration. From these results, it can be concluded that diclofenac only induced structural perturbations to DMPC bilayers.

3.2. Fluorescence measurements of isolated unsealed human erythrocyte membranes (IUM) and large unilamellar vesicles (LUV)

The concentration-dependent effect of diclofenac on IUM was explored at two different depths of the erythrocyte membrane lipid bilayer: at the hydrophilic/hydrophobic level, estimated from the laurdan fluorescence spectral shift through the GP parameter, and in the deep hydrophobic core determined by the DPH steady-state fluorescence anisotropy (*r*). As shown in Fig. 3, diclofenac incorporation at increasing concentrations up to 2 mM only induced a mild increase of the laurdan GP and a mild decrease of DPH anisotropy up to 0.3 mM. These results imply a moderate rigidifying of the bilayer polar group's organization, and a moderate disorder of the hydrophobic acyl chains. Figs. 4 and 5 show that the incorporation of diclofenac to DMPC LUV in the same range of increasing concentrations induced a mild decrease of the GP values at 18 °C and a mild increase at 37 °C. Similar measurements of DPH




Fig. 5. Effects of sodium diclofenac on the anisotropy of DPH embedded in DMPC large unilamellar vesicles (LUV) at 18 and 37 °C.

fluorescence anisotropy (r) showed that while at 18 °C remained practically constant, there was a mild increase at 37 °C, implying a mild ordering of DMPC acyl chains. It should be taken into account that at 37 °C this lipid is in a much more fluid state than at 18 °C.

3.3. Scanning Electron Microscope (SEM) studies of human erythrocytes

SEM examinations of human erythrocytes incubated with diclofenac in the range of 0.01–2 mM indicated that the drug induced changes different from the normal biconcave morphology of most red blood cells (Fig. 6A). With 0.01 mM diclofenac (Fig. 6B), about a third of the cells changed their discoid normal shape into stomatocytes (a cup-shaped form with evagination of one surface and a deep invagination of the opposite face), and another third into knizocytes (red blood cells with two or three concavities due to indentations in the cell membrane), with 0.1 mM (Fig. 6C) the majority of the cells are stomatocytes, with 0.3 mM (Fig. 6D) the majority of the cells are knizocytes, with 0.5 mM (Fig. 6E) there are many stomatocytes and spherostomatocytes (cells with a visible change towards spheroid morphology with lightly or minor cupped profiles), and with 2 mM (Fig. 6F) the majority of the cells are knizocytes.

4. Discussion

Studies on the effects of diclofenac on cell membranes are not too frequent, but a few of them have indicated that the drug produced some functional effects. Thus, Uyemura et al. [24] reported that diclofenac induced membrane permeability transition to rat renal mitochondrial, and uncoupled oxidative phosphorylation [25], increased the fluidity of mouse splenocyte membrane [26], caused cell death in human hepatocytes through mitochondrial permeabilization [27], and activates K^+ channels exerting its effect at both inner and outer sides of rat cerebellar granule cells [28]. However, it becomes amazing the lack of information on the effects induced by diclofenac into human erythrocytes. One of the few reports indicates that diclofenac at the therapeutic and higher concentrations exerts in vitro an inhibition on H_2O_2 forced erythrocytic membrane lipid peroxidation as well as increased hemolysis [29]. The present study presents the following evidence that diclofenac affects human erythrocytes and molecular models of its cell membrane: X-ray diffraction experiments were performed on bilayers made up of DMPC and DMPE, classes of the major phospholipids present in the outer and inner erythrocyte membrane, respectively. Results showed that diclofenac




Fig. 6. Effects of sodium diclofenac on the morphology of human erythrocytes. SEM images of (A) untreated erythrocytes; incubated with (B) 0.01 mM; (C) 0.1 mM; (D) 0.3 mM; (E) 0.5 mM, and (F) 2 mM sodium diclofenac.

interacted practically only with DMPC, affecting both its polar head and acyl chain regions. DMPC and DMPE differ only in their terminal amino groups, being these $^*N(CH_3)_3$ in DMPC and *NH_3 in DMPE. Moreover, both molecular conformations are very similar in their dry crystalline phases [23] with the hydrocarbon chains mostly parallel and extended and the polar head groups lying perpendicularly to them. However, the gradual hydration of DMPC results in water filling the highly polar interbilayer spaces with the resulting width increase. This phenomenon allows the incorporation of diclofenac into DMPC bilayers with the resulting disruption of DMPC bilayer structure. Similar results were observed on unilamellar liposomes and floating Langmuir monolayers of soya phosphatidylcholine [30]. Given the amphiphilic nature of sodium diclofenac it is very likely that it locates into DMPC bilayer in such a way that its negatively charged carboxyl group electrostatically interacts with the positively charged DMPC terminal $^*N(CH_3)_3$ group, while the apolar ring sits in the neighborhood of the hydrophobic acyl region of DMPC. A somewhat similar conclusion has been suggested by Lichtenberger [7] and reported on the basis of ^{31}P NMR spectra of phosphatidylcholine liposomes [31]. As diclofenac is negatively charged its location in the polar region of DMPC modifies the electrostatic interactions between the lipid phosphate and terminal amino groups disrupting its bilayer structure. However, other explanations cannot be disregarded. One of them considers the hydration, which plays an important role in the stability of phospholipids bilayers. In both gel and liquid crystalline phases water molecules are bound to DMPC head group. The molecules are oriented and form a hydration shell through hydrogen bonding with the polar groups. In the presence of diclofenac the amount of water molecules, their orientation and hydrogen bonds can be affected. Thus, the presence of diclofenac negatively charged carboxyl group located near the phosphate region can compete for the formation of hydrogen bonds with the water molecules of the hydration shell. The weakening of the water binding to the phosphate region would modify DMPC affinity for water and consequently the packing of its bilayer. On the other hand, DMPE is not significantly affected by diclofenac, the explanation lying in that the molecules pack tighter than those of DMPC due to their smaller polar groups and higher effective charge, resulting in a very stable bilayer system held by electrostatic interactions and hydrogen bonds [23].

The toxic effect due to diclofenac treatment was also observed on erythrocytes in the same concentration range. In fact, scanning electron microscopy (SEM) observations of human erythrocytes indicated that they underwent a morphological alteration as their discoid normal shape changed with increasing diclofenac concentrations. According to the bilayer-couple hypothesis, the preferential interaction of diclofenac with DMPC, a class of lipid mainly located in the outer monolayer of the erythrocyte membrane, should have induced echinocytosis instead of the mostly observed stomatocytosis. The explanation for this discrepancy could be based on the lipid scrambling mechanism proposed by Schrier et al. [32]. According to it, some cationic amphipaths produce a rapid scrambling of the erythrocyte bilayer with phosphatidylcholines (PC) and sphingomyelins (SM) moving inward while phosphatidylethanolamines (PE) moves outward along with phosphatidylserines (PS). Thus, the interaction of diclofenac with PC in the inner monolayer would lead to stomatocytosis, an effect that can be produced by as little as 0.6% enrichment of the cytoplasmic monolayer [32]. An alternative explanation might lie in that diclofenac interacted with proteins located in the inner monolayer of the erythrocyte membrane. On the other hand, the concentration-dependent effect of diclofenac on isolated unsealed human erythrocyte membrane (IUM) was explored by fluorescence spectroscopy at two different depths: at the hydrophilic/hydrophobic level and in the deep hydrophobic core of the membrane lipid bilayer. The incorporation of diclofenac induced a mild increase of the generalized polarization, result that can be rationalized as a moderate decrease in the molecular dynamics and/or water content at the glycerol backbone level of the polar head groups, whereas the moderate decrease of the anisotropy up to 0.3 mM implies a moderate disorder of the acyl chain packing order at the bilayer hydrophobic core.

Interestingly, the corresponding results of both probes embedded in DMPC LUV clearly show that in liquid crystalline phase at 37 °C diclofenac exerts a rigidifying effect on DMPC bilayer. The difference in DPH anisotropy behavior in both systems could be ascribed to the presence of lipid–protein interfaces in IUM.

Amazingly, the therapeutic range for plasma diclofenac concentrations and the relationship of plasma concentration to clinical response and toxicity have not been clearly established. Nevertheless, the experimental findings are certainly of interest as they indicate that a diclofenac concentration as low as 10 μM affects the human erythrocyte shape. It must be considered that alteration of the normal biconcave shape of red blood cells increases their resistance to entry into capillaries [33], which could contribute to decreased blood flow, loss of oxygen, and tissue damage through microvascular occlusion [34].

Acknowledgements

The authors thank Fernando Neira for his technical assistance and to DAAD for the Ph.D. scholarship (to M.M.). This work was supported by grants from the FONDECYT (1060990) and the CONICYT-BMBF (065-4-2007).

References

- [1] W.E. Smalley, W.A. Ray, J.R. Daughterty, M.R. Griffin, Nonsteroidal anti-inflammatory drugs and the incidence of hospitalization for peptic ulcer disease in elderly persons, *Am. J. Epidemiol.* 141 (1995) 539–545.
- [2] W. Tomisato, K. Tanaka, T. Katsu, H. Kakuta, K. Sasaki, S. Tsutsumi, T. Hoshino, M. Aburaya, D. Li, T. Tsuchiya, K. Suzuki, K. Yokomizo, T. Mizushima, Membrane permeabilization by non-steroidal anti-inflammatory drugs, *Biochem. Biophys. Res. Commun.* 323 (2004) 1032–1039.
- [3] A. Burke, E. Smyth, G.A. FitzGerald, Analgesic–antipyretic agents: pharmacotherapy of gout, in: L.L. Brunton (Ed.), Goodman & Gilman's The Pharmacological Basis of Therapeutics, 11th ed., McGraw-Hill, USA, 2006, ch. 26.
- [4] L. Biagini Lopes, M.V. Scarpa, N.L. Pereira, L. Cardoso de Oliveira, A. Gomes Oliveira, Interaction of sodium diclofenac with freeze-dried soya phosphatidylcholine and unilamellar liposomes, *Braz. J. Pharm. Sci.* 42 (2006) 497–504.
- [5] A. Inoue, S. Muranaka, H. Fujita, T. Kanno, H. Tamai, K. Utsumi, Molecular mechanism of diclofenac-induced apoptosis of promyelocytic leukemia: dependency on reactive oxygen species, akt, bid, cytochrome c, and caspase pathway, *Free Radic. Biol. Med.* 37 (2004) 1290–1299.
- [6] Martindale, in: J.E.F. Reynolds (Ed.), The Extra Pharmacopoeia, 30th ed., The Pharmaceutical Press, London, 1993, pp. 10–11.
- [7] L.M. Lichtenberger, Where is the evidence that cyclooxygenase inhibition is the primary cause of nonsteroidal anti-inflammatory drug (NSAID)-induced gastrointestinal injury? Topical injury revisited, *Biochem. Pharmacol.* 61 (2001) 631–637.
- [8] M.N. Giraud, C. Motta, J.J. Romero, G. Bommelaer, L.M. Lichtenberger, Interaction of indomethacin and naproxen with gastric surface-active phospholipids: a possible mechanism for the gastric toxicity of non-steroidal anti-inflammatory drugs (NSAIDs), *Biochem. Pharmacol.* 57 (1999) 247–254.
- [9] H. Aki, M. Yamamoto, Biothermodynamic characterization of erythrocyte hemolysis induced by phenothiazine derivatives and anti-inflammatory drugs, *Biochem. Pharmacol.* 39 (1990) 396–398.
- [10] Y. Masubuchi, S. Nakayama, T. Horie, Role of mitochondrial permeability transition in diclofenac-induced hepatocyte injury in rats, *Hepatology* 35 (1997) 544–551.
- [11] J.Y. Chen, W.H. Huestis, Role of membrane lipid distribution in chlorpromazine-induced shape change of human erythrocytes, *Biochim. Biophys. Acta* 1323 (1997) 299–309.
- [12] J.M. Boon, B.D. Smith, Chemical control of phospholipid distribution across bilayer membranes, *Med. Res. Rev.* 22 (2000) 251–281.
- [13] P.F. Devaux, A. Zachowsky, Maintenance and consequences of membrane phospholipids asymmetry, *Chem. Phys. Lipids* 73 (1994) 107–120.
- [14] M. Suwalsky, S. Mennickent, B. Norris, F. Villena, F. Cuevas, C.P. Sotomayor, The antiepileptic drug diphenylhydantoin affects the structure of the human erythrocyte membrane, *Z. Naturforsch.* 59c (2004) 427–431.
- [15] M. Suwalsky, S. Mennickent, B. Norris, F. Villena, C.P. Sotomayor, Effects of the antiepileptic drug carbamazepine on human erythrocytes, *Toxicol. In Vitro* 20 (2006) 1363–1369.
- [16] M. Suwalsky, F. Villena, C.P. Sotomayor, S. Bolognin, P. Zatta, Human cells and cell membrane molecular models are affected in vitro by chlorpromazine, *Biophys. Chem.* 135 (2008) 7–13.
- [17] R. Koyanova, M. Caffrey, Phases and phase transitions of the phosphatidylcholines, *Biochim. Biophys. Acta* 1376 (1998) 91–145.
- [18] M. Suwalsky, C. Rivera, F. Villena, C.P. Sotomayor, M. Jemiola-Rzeminska, K. Strzalka, Arsenite interactions with phospholipids bilayers as molecular models for the human erythrocyte membrane, *Biophys. Chem.* 127 (2007) 28–35.
- [19] T. Parasassi, E. Gratton, Membrane lipid domains and dynamics as detected by laurdan fluorescence, *J. Fluoresc.* 5 (1995) 59–69.

- [20] J.T. Dodge, C. Mitchell, D.J. Hanahan, The preparation and chemical characterization of haemoglobin-free ghosts of human erythrocytes, *Arch. Biochem. Biophys.* 100 (1963) 119–130.
- [21] J.R. Lakowicz, *Principles of Fluorescence Spectroscopy*, 2nd ed. Kluwer/Plenum, New York, 1999.
- [22] T. Parasassi, G. De Stasio, A. D'Ubaldo, E. Gratton, Phase fluctuation in phospholipid membranes revealed by laurdan fluorescence, *Biophys. J.* 57 (1990) 1179–1186.
- [23] M. Suwalsky, Phospholipid bilayers, in: J.C. Salamone (Ed.), *Polymeric Materials Encyclopedia*, vol. 7, CRC, Boca Raton, FL, 1996, pp. 5073–5078.
- [24] S.A. Uyemura, A.C. Santos, F.E. Mingatto, M.C. Jordani, C. Curti, Diclofenac sodium and mefenamic acid: potent inducers of the membrane permeability transition in renal cortex mitochondria, *Arch. Biochem. Biophys.* 342 (1997) 231–235.
- [25] F.E. Mingatto, A.C. Santos, S.A. Uyemura, M.C. Jordani, C. Curti, In vitro interaction of nonsteroidal anti-inflammatory drugs on oxidative phosphorylation of rat kidney mitochondria: respiration and ATP synthesis, *Arch. Biochem. Biophys.* 334 (1996) 303–308.
- [26] H. Ferreira, M. Lúcio, J.L.F.C. Lima, A. Cordeiro-da-Silva, J. Tavares, S. Reis, Effect of anti-inflammatory drugs on splenocyte membrane fluidity, *Anal. Biochem.* 339 (2005) 144–149.
- [27] W.P. Siu, P.B.L. Pun, C. Latchoumycandane, U.A. Boelsterli, Bax-mediated mitochondrial outer membrane permeabilization (MOMP), distinct from the mitochondrial permeability transition, is a key mechanism in diclofenac-induced hepatocyte injury: multiple protective roles of cyclosporine A, *Toxicol. Appl. Pharmacol.* 227 (2008) 451–461.
- [28] L.-Y. Liu, X.-W. Fei, Z.-M. Li, Z.-H. Zhang, Y.-A. Mei, Diclofenac, a nonsteroidal anti-inflammatory drug, activates the transient outward K^+ current in rat granule cells, *Neuropharmacol.* 48 (2005) 918–926.
- [29] H. Orhan, G. Sahin, In vitro effects of NSAIDs and paracetamol on oxidative stress-related parameters of human erythrocytes, *Exp. Toxicol. Pathol.* 53 (2001) 133–140.
- [30] S.M.B. Souza, O.N. Oliveira Jr., M.V. Scarpa, A.G. Oliveira, Study of the diclofenac/phospholipid interactions with liposomes and monolayers, *Colloids Surf. B* 36 (2004) 13–17.
- [31] L.B. Lopes, M.V. Scarpa, G.V.J. Silva, D.C. Rodrigues, C.V. Santillini, A.G. Oliveira, Studies on the encapsulation of diclofenac in small unilamellar liposomes of soya phosphatidylcholine, *Colloids Surf. B* 39 (2004) 151–158.
- [32] S.L. Schrier, A. Zachowski, P.F. Devaux, Mechanisms of amphipath-induced stomatocytosis in human erythrocytes, *Blood* 79 (1992) 782–786.
- [33] S. Svetina, D. Kuzman, R.E. Waugh, P. Zibert, B. Zeks, The cooperative role of membrane skeleton and bilayer in the mechanical behaviour of red blood cells, *Bioelectrochem.* 62 (2004) 107–113.
- [34] S.L. Winski, D.E. Carter, Arsenate toxicity in human erythrocytes: characterization of morphological changes and determination of the mechanism of damage, *J. Toxicol. Env. Health, Part A* 53 (1998) 345–355.