

## MAGNETITE AND MAGNETOTAXIS IN ALGAE

F. F. TORRES DE ARAUJO,\* M. A. PIRES,\* R. B. FRANKEL,<sup>†</sup> AND C. E. M. BICUDO<sup>‡</sup>\*Departamento de Física, Universidade Federal do Ceará, Fortaleza, Ceará, Brazil; <sup>†</sup>Francis Bitter National Magnet Laboratory, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139; and <sup>‡</sup>Instituto de Botânica, São Paulo, Brazil

**ABSTRACT** Magnetotactic algae of the genus *Anisonema* (Euglenophyceae) have been isolated from a coastal mangrove swamp in northeastern Brazil. The magnetotactic response is based on a permanent magnetic dipole moment per cell  $\sim 7 \cdot 10^{-10}$  emu. Each cell contains many magnetite ( $\text{Fe}_3\text{O}_4$ ) particles organized in chains.

## INTRODUCTION

Magnetite ( $\text{Fe}_3\text{O}_4$ ) has been reported as a biomineralization product in bacteria (1, 2) and in many multicellular organisms (3). In magnetotactic bacteria, magnetite particles are deposited intracellularly in membrane vesicles (magnetosomes) (4) and have dimensions ranging from 400 to 1,200 Å and morphologies that are species specific

(5). The particles are often arranged in a chain that imparts a magnetic dipole moment to the cell that is responsible for the magnetotactic response (6). In chitons, magnetite particles of irregular shape are deposited extracellularly in an organic matrix on the tooth denticles of the radula (7). In the other eucaryotic organisms, magnetite particles have only been identified magnetically in whole organisms and after extraction (8). Thus their in vivo arrangement is unknown, although they are thought to be responsible for magnetic effects on behavior (1, 3, 9). Here

Correspondence should be addressed to R. B. Frankel.




FIGURE 1 (a) Light micrograph of magnetotactic algae oriented in a magnetic field at the edge of a water drop. Cells are  $\sim 20 \mu\text{m}$  long,  $12 \mu\text{m}$  wide and  $4 \mu\text{m}$  thick. (b) Light micrograph of an individual alga.




FIGURE 2 (a) Electron micrograph of *A. platysomum* (bar length: 5  $\mu\text{m}$ ). (b) Electron micrograph of periphery of cell showing chains of particles (bar length: 2.5  $\mu\text{m}$ ). (c) Electron micrograph of particle chain (bar length: 0.5  $\mu\text{m}$ ).




FIGURE 3 Electron micrograph showing relative orientations of particles and chains at edge of cell.

we report that eucaryotic cells, euglenoid, motile, magnetotactic algae of the genus *Anisonema*, contain many  $\text{Fe}_3\text{O}_4$  particles organized in chains.

## METHODS

The algae were collected in brackish water sediments from a coastal mangrove swamp near Fortaleza, Brazil. Their magnetotactic response was observed microscopically in drops of water and sediment with small bar magnets or Helmholtz coils placed on or near the microscope stage as previously described for magnetotactic bacteria (10). Cells eventually migrated along magnetic field lines either parallel or antiparallel to the field direction, that is, they were either North-seeking or South-seeking, respectively. Killed cells rotated and oriented along magnetic field lines. Cells were flat and ovate (Fig. 1) with length and width of  $\sim 20$  and  $\sim 12$   $\mu\text{m}$ , respectively. There were two flagella per cell, one trailing. On the basis of light microscopic examination the organism was identified as *Anisonema platysomum* Skuja (11).

Algae were isolated from the sediments by using their magnetotactic response to collect them at the edge of a drop of water and sediment on a microscope slide, where they were picked up with a micropipette and deposited on a carbon-coated electron microscope grid. They were then fixed in situ by incubation in 1% glutaraldehyde in  $\text{K}_2\text{HPO}_4$  buffer for several minutes. The grids were examined in a 100 keV VG HB5 STEM electron microscope (VG microscopes, East Grinstead, U.K.) equipped

TABLE I  
LATTICE SPACINGS FROM ELECTRON  
DIFFRACTION IN ALGAL PARTICLES

Algal particles d-spacings	$\text{Fe}_3\text{O}_4$ d-spacings	Lattice planes hkl
$\text{\AA}$	$\text{\AA}^*$	
$4.8 \pm 0.5$	4.8	111
4.2		
2.9	2.967	220
2.6	2.532	311
2.1	2.099	400
1.8	1.715	422
1.6	1.616	511
1.5	1.485	440
1.2	1.281	533

\*ASTM Card 19-629.




FIGURE 4 Values of the  $180^\circ$  rotation time for a killed alga suspended in water after reversal of the magnetic field plotted as a function of magnetic field strength. The line is a theoretical fit of Eq. 1 to the data with  $M = 5 \cdot 10^{-10}$  emu and  $R = 8$   $\mu\text{m}$ .

with energy dispersive x-ray analysis (EDX) and electron diffraction capabilities (Figs. 2 and 3). The size of the algae prevented penetration of the electron beam at the centers of the cells (Fig. 2a). However, numerous electron-opaque particles arranged in chains oriented more or less parallel to the long axis of the cell could be seen at the electron-transparent periphery of each cell (Figs. 2b and 3). Individual particles were roughly arrowhead or tooth shaped (Fig. 2c) with lengths of 800–1,800  $\text{\AA}$  (1,400  $\text{\AA}$  average) and widths of 400–500  $\text{\AA}$  (480- $\text{\AA}$  average).

## RESULTS

EDX spectra were recorded with the electron beam focused on and off the particles. Comparison of the spectra showed that the particles were highly enriched in iron compared to the cell wall and cytoplasm. No Fe x rays were observed beyond the apparent cell boundary, indicating that the particles were intracellular. Electron diffraction patterns were recorded for forty individual particles. The patterns indicated that each particle was a single crystal with orientation that varied from particle to particle along the chain. Analysis of the diffraction patterns gave interplanar spacings that are consistent with the known interplanar spacings of  $\text{Fe}_3\text{O}_4$  (Table I).

Magnetotactic bacteria, cocci and spirilla, were also found in the same sediments with the algae. These organisms, typically 1–3  $\mu\text{m}$ , contained single or double chains of cuboidal-shaped magnetite particles with linear dimensions 400–600  $\text{\AA}$  similar to those reported previously (1, 2). Arrow-head shaped particles similar to those in the algae have previously been found in bacteria from sediments in New Zealand (12) and New England (13).

Thus magnetotactic algae of the genus *Anisonema* contain numerous chains of  $\text{Fe}_3\text{O}_4$  particles that appear to be located in or near the cell wall. The individual particles are within the single-magnetic-domain size range for  $\text{Fe}_3\text{O}_4$ . Hence, each chain is a permanent, magnetic dipole (14). If the moments of all the chains are oriented parallel to each other, a cell would have a magnetic dipole moment equal to the sum of the moments of all its particles. An estimate of the magnetic dipole moment of whole algae was obtained from measurements of the  $180^\circ$  rotation time

of killed cells suspended in water, after reversal of the direction of the magnetic field produced by a pair of Helmholtz coils. Using the coefficient of viscous drag for a flat disk, the rotation time  $\tau$  is related to the magnetic moment  $M$  by the equation (15)

$$\tau = \frac{32R^3\eta}{3MH} \ln\left(\frac{2MH}{kT}\right), \quad (1)$$

where  $R$  is the radius of the cell,  $\eta$  is the viscosity of water,  $H$  is the magnetic field, and  $kT$  is Boltzmann's constant multiplied by temperature. Experimental sets of values of  $\tau$  vs.  $H$  for eleven algal cells were each fit (Fig. 4) with Eq. 1, yielding an average permanent, magnetic moment per cell  $M = 6.7 \cdot 10^{-10}$  emu. This is  $\sim 1,000$  times the magnetic moment of a typical magnetotactic bacterium. The saturation magnetization in magnetite is  $480 \text{ emu cm}^{-3}$ . Therefore the magnetic moment of an average-sized particle ( $1,400 \times 480 \times 480 \text{ \AA}$ ) is  $1.5 \cdot 10^{-13}$  emu. Hence, each algal cell must contain on the order of  $3 \cdot 10^3$  magnetite particles, with the particles occupying  $\sim 0.2\%$  of the cell volume.

Although the motility of the algae in a magnetic field is more complex than magnetotactic bacteria, the magnetotactic response mechanism of the algae appears to be similar to that in magnetotactic bacteria, i.e., passive orientation of the cell by the torque exerted by the magnetic field on its permanent, magnetic dipole (14). The fact that the algal magnetic moment is three orders of magnitude larger than typical bacterial moments means that algae and bacteria have similar ratios of magnetic torque to viscous drag, that is, they have similar recovery times after deorientation events (16).

The biological significance of magnetotaxis in these organisms is unknown at present, but since Fortaleza is within  $4^\circ$  of the magnetic equator, both North-seeking and South-seeking algal cells will be directed horizontally. The genus *Anisonema* is typically heterotrophic and the sediments are a rich source of nutrients. Thus magnetotaxis in these cells might serve to keep them in the sediments and away from high oxygen tension, high light intensity, and high temperatures at the water surface.

We thank A. Garratt-Reed for electron microscopy, R. P. Blakemore for technical suggestions and comments, F. A. Germano for computer analysis, and J. Danon, D. M. S. Esquivel, H. Lins de Barros, and S. Mann for discussions.

R. B. Frankel was partially supported by the Office of Naval Research. The Francis Bitter National Magnet Laboratory is supported by the National Science Foundation.

Received for publication 11 December 1985 and in final form 24 March 1986.

## REFERENCES

1. Frankel, R. B., R. P. Blakemore, and R. S. Wolfe. 1979. Magnetite in freshwater magnetotactic bacteria. *Science (Wash. DC)*. 203:1355-1356.
2. Towe, K. M., and T. T. Moench. 1981. Electron-optical characterization of bacterial magnetite. *Earth Planet. Sci. Lett.* 52:213-220.
3. Lowenstam, H. A., and J. L. Kirschvink. 1985. Iron biomineralization: a geobiological perspective. In *Magnetite Biomineralization and Magnetoreception in Organisms*. J. L. Kirschvink, D. S. Jones, and B. J. MacFadden, editors. Plenum Publishing Corp., New York. 3-13.
4. Balkwill, D. L., D. Maratea, and R. P. Blakemore. 1980. Ultrastructure of a magnetotactic spirillum. *J. Bacteriol.* 141:1399-1408.
5. Mann, S., R. B. Frankel, and R. P. Blakemore. 1984. Structure, morphology and crystal growth of bacterial magnetite. *Nature (Lond.)*. 310:405-407.
6. Frankel, R. B., and R. P. Blakemore. 1980. Navigational compass in magnetic bacteria. *J. Magn. Magn. Mater.* 15-18:1562-1564.
7. Towe, K. M., and H. A. Lowenstam. 1967. Ultrastructure and development of iron mineralization in the radular teeth of *C. stelleri* (Mollusca). *J. Ultrastruct. Res.* 17:1-13.
8. Walker, M. M., J. L. Kirschvink, S. B. R. Chang, and A. E. Dizon. 1984. A candidate magnetic sense organ in the yellowfin tuna, *T. albacores*. *Science (Wash. DC)*. 224:751-753.
9. Lins de Barros, H. G. P., D. M. S. Esquivel, J. Danon, and L. P. H. Oliveira. 1981. Magnetotactic algae. *Ann. Acad. Bras. Cienc.* 54:258-259.
10. Blakemore, R. P., and R. B. Frankel. 1981. Magnetic navigation in bacteria. *Sci. Am.* 245(6):58-65.
11. Skuja, J. 1939. Beitrag zur Algenflora Lettlands II. *Acta Horti. Bot. Univ. Latv.* 11-12:41-169.
12. Blakemore, R. P., R. B. Frankel, and A. J. Kalmijn. 1981. South-seeking magnetotactic bacteria in the southern hemisphere. *Nature (Lond.)*. 286:384-385.
13. Blakemore, R. P., N. A. Blakemore, D. A. Bazylinski, and T. T. Moench. 1985. The magnetotactic bacteria. *Bergey's Manual of Systematic Bacteriology*. Vol. 3. In press.
14. Frankel, R. B. 1984. Magnetic guidance of organisms. *Annu. Rev. Biophys. Bioeng.* 14:85-103.
15. Esquivel, D. M. S., H. G. P. Lins de Barros, M. Fariña, P. H. A. Aragão, and J. Danon. 1983. Microorganisms magnetotactiques de la region de Rio de Janeiro. *Biol. Cell.* 47:227-234.
16. Rosenblatt, C., R. B. Frankel, and R. P. Blakemore. 1985. A birefringence relaxation determination of rotational diffusion of magnetotactic bacteria. *Biophys. J.* 47:323-325.