

# RESONANCE RAMAN STUDY OF THE PINK MEMBRANE PHOTOCHEMICALLY PREPARED FROM THE DEIONIZED BLUE MEMBRANE OF *H. HALOBIUM*

C. PANDE,\* R. H. CALLENDER,\* C.-H. CHANG,‡ AND T. G. EBREY‡

\*Physics Department, City College of the City University of New York, New York 10031; and

‡Department of Physiology and Biophysics, University of Illinois at Urbana-Champaign, Urbana, Illinois

**ABSTRACT** We report here the Resonance Raman spectrum of a 'pink' membrane ( $\lambda_{\max} \sim 495$  nm) photochemically generated from the deionized 'blue' membrane (Chang et al., 1985). Comparison of the Raman spectrum of the pink membrane with that of the model compounds, as well as the chromophore extraction data, indicate that the chromophore in the pink membrane is in the 9-*cis* configuration. The Schiff base peak at  $\sim 1,652$   $\text{cm}^{-1}$  shifts to  $\sim 1,622$   $\text{cm}^{-1}$  upon deuteration of the pink membrane, showing that the chromophore is bound to the bacterio-opsin by a protonated Schiff base linkage. The location of the Schiff base peak, as well as the 30  $\text{cm}^{-1}$  shift that it undergoes upon deuteration, are quite different from the corresponding values for the native bacteriorhodopsin, suggesting differences in the local environment for the Schiff base in these pigments.

## INTRODUCTION

Bacteriorhodopsin, the only protein present in the purple membrane of the extreme halophile *Halobacterium halobium*, is a retinal containing protein. The retinal molecule is bound to the protein, in molar equivalence, by a Schiff base linkage to the  $\epsilon$ -amino group of Lys<sub>216</sub>. Light absorbed by this chromoprotein is utilized by the bacterium to create a proton gradient across its cell membrane. This protonmotive force is then utilized to synthesize ATP, which is subsequently used to drive endergonic cellular metabolic processes (see reviews by Stoekenius and Bogomolni, 1982; Ebrey, 1982).

Light-adapted bacteriorhodopsin, bR<sub>568</sub>, is a metastable species containing retinal in the all-*trans* configuration. This species decays thermally to the more stable dark-adapted form, bR<sub>560</sub>, which contains retinal in both the 13-*cis* and the all-*trans* isomeric forms in roughly equal amounts. It has been found recently that a new species, the blue membrane ( $\lambda_{\max} \sim 603$  nm), is formed from bR<sub>560</sub> upon complete removal of all cations by any of a variety of techniques (Kimura et al., 1984; Chen et al., 1984; Chang et al., 1985). The blue membrane, like bR<sub>560</sub>, also consists of a heterogeneous pigment population, with the chromophore in both the 13-*cis* and the all-*trans* isomeric forms, as revealed by chromophore extraction as well as resonance Raman studies (Smith et al., 1985; Pande et al., 1985). The transition from bR<sub>560</sub> to the blue membrane is revers-

ible. The reversibility is rather nonspecific for the ion type, but the amount of ions required to cause the change follows the order:  $\text{M}^+ > \text{M}^{2+} > \text{M}^{3+}$ . The bR<sub>560</sub>, regenerated from the blue membrane by the addition of ions, is virtually indistinguishable from the native bR<sub>560</sub>, as judged by the absorption as well as the resonance Raman spectroscopy (Pande et al., 1985).

Recently, while studying the photochemistry of the blue membrane, we found that its absorption maximum shifts from  $\sim 603$  nm to  $\sim 495$  nm upon irradiation with light above 640 nm, and the membrane appears pink in color (Chang et al., in preparation). This species is fairly stable in the dark, but converts back to the blue membrane upon illumination with short wavelength light ( $\lambda < 500$  nm). In the presence of cations, even though the pink membrane is stable, illumination with short wavelength light results in the formation of the native purple membrane, not the blue membrane.

These results are very interesting, because they show that cations not only regulate the absorption properties of bR, but also determine its photochemical reactions (i.e., the photoproduct of the blue membrane is different from that of the purple membrane). In order to understand the above results and to gain more structural insight, we have used resonance Raman spectroscopy. This technique allows the determination of the isomeric state of the chromophore as well as the nature of the chromophore-protein interactions. Results from our resonance Raman

measurements, in agreement with the chromophore extraction data, show that the chromophore exists predominantly in the 9-*cis* isomeric form in the pink membrane. The chromophore appears to be bound to the protein via a protonated Schiff base linkage. The Schiff base peak position, as well as its shift upon deuteration, are quite different from those observed in bR, suggesting that the Schiff base environment is different in the pink membrane and in its ultimate precursor, the native bacteriorhodopsin.

## MATERIALS AND METHODS

The protonated Schiff base of 9-*cis* retinal with N-butylamine was prepared as described previously (Pande et al., 1981). The reaction product was confirmed by its absorption maximum around 440 nm, compared to a considerably blue shifted  $\lambda_{\text{max}}$  of the unprotonated Schiff base ( $\sim 360$  nm).

Pink membrane was prepared from the papain-treated blue membrane (Chang et al., 1985) by irradiation with deep red light ( $\lambda > 640$  nm) at  $\sim 4^\circ\text{C}$  (pH 4.8) using a glass cut-off filter (CS 2-58) in front of a 400 W slide projector. After  $\sim 215$  min of irradiation,  $\sim 80\%$  of the blue membrane was converted to the pink membrane with a  $\lambda_{\text{max}}$  around 495 nm.

Deuterated form of the pink membrane was prepared by first concentrating the pink membrane by centrifugation at  $4^\circ\text{C}$  and 20,000 g for  $\sim 4$  h, followed by suspension in  $\text{D}_2\text{O}$ . This procedure was repeated two times with the final pellet dissolved in minimum amount of  $\text{D}_2\text{O}$ . Extreme care was taken during this process to avoid contamination from extraneous ions.

The isomeric composition of the chromophores was determined by the retinal oxime extraction method as described earlier (Pande et al., 1985).

Resonance Raman spectra of the pink and the blue membrane were measured at 80 K using apparatus previously described (Pande et al., 1981). At this temperature and the illumination conditions, no spectral changes could be observed even upon prolonged irradiation with the exciting light. This would suggest that the pink membrane does not slowly photoconvert to other species under these conditions. We have confirmed this by low temperature absorption spectroscopy, which shows that 476 nm light does not lead to any photoproducts of the pink membrane at 80 K.

The pink membrane sample was excited with  $\sim 5$  mW of 476 nm light from an  $\text{Ar}^+$  laser (model 165; Spectra-Physics, Inc., Mountain View, CA). Under these conditions the blue membrane contribution to the pink membrane spectrum would be negligible (due to the lower resonance enhancement at this wavelength). The blue membrane spectrum was obtained with  $\sim 5$  mW of the 488-nm light from the same laser, in order to obtain a somewhat better spectrum, than was possible with the 476-nm excitation. The protonated Schiff base spectrum was measured in a flow system. The temperature of the sample was maintained around  $10^\circ\text{C}$  and the sample flow velocity was  $\sim 1,500$  cm/s. The sample was excited with 476 nm light ( $\sim 2$  mW) as above. Under these conditions, the photoconversion probability in the flow experiment is less than a few percent (Pande et al., 1981) and the Raman spectrum is essentially due to the 9-*cis* protonated Schiff base.

Raman spectra were obtained with a Spex Triplemate spectrometer (model 1877; Spex Industries, Inc., Edison, NJ) with a 1,200 gr/mm grating. The detector system in this set-up consists of a cooled solid state detector (model 1420-2; EG&G Princeton Applied Research Corp., Princeton, NJ) coupled to a controller unit (model 1218; EG&G Princeton Applied Research). This multichannel Raman system, controlled by an LSI-11 computer, can collect data simultaneously in about 700 pixels. This allows a "viewing window" of  $\sim 900$   $\text{cm}^{-1}$ , with a spectral resolution of 8  $\text{cm}^{-1}$ , under our experimental conditions. The wavenumber scale of the spectra was calibrated against the Raman lines of toluene.

## RESULTS AND DISCUSSION

Upon irradiation with deep red light ( $\lambda > 640$  nm), the blue membrane ( $\lambda_{\text{max}} \sim 603$  nm) partially converts to the pink membrane ( $\lambda_{\text{max}} \sim 495$  nm). The kinetics of this conversion is very slow as can be seen in Fig. 1, which demonstrates the effect of red light on a sample of the papain-treated blue membrane. The blue membrane does not completely convert to the pink species, even after 215 min of irradiation. Further irradiation did not have any effect on this pigment composition, suggesting that the system had reached a photochemical equilibrium. Since the residual unconverted species has the absorption maximum at  $\sim 600$  nm and it shows an isosbestic point with the pink membrane spectrum, we believe that it represents the unconverted blue membrane. Our inability to convert this  $\sim 600$  nm absorbing species completely to the 495 nm absorbing species is probably due to a very much higher quantum efficiency of the back reaction, compared to the forward one. The concentration of this  $\sim 600$  nm absorbing species in these preparations was estimated to be  $20 \pm 5\%$ . Under identical irradiation conditions, the native (unproteolyzed) deionized blue membrane reaches a photochemical equilibrium which contains greater amounts of the long wavelength absorbing species. Thus, in order to avoid the complications arising in the interpretation of the spectra, we chose to study the pink membrane obtained from the papain-treated blue membrane.

The resonance Raman spectrum of the pink membrane is shown in Fig. 2 *a*. For comparison, we have also included the resonance Raman spectra of the blue membrane that was used to generate the pink membrane (Fig. 2 *b*), and the protonated Schiff base of 9-*cis* retinal (Fig. 2 *c*). One striking feature of these figures is the clear enhancement of




FIGURE 1 Absorption changes in the papain-treated blue membrane (pH 4.8) following irradiation for various times with deep red light ( $\lambda > 640$  nm) at  $\sim 4^\circ\text{C}$ , using a glass cut-off filter (CS 2-58) in front of a 400 W slide projector. The sample concentration was  $\sim 10$   $\mu\text{M}$ . Spectra represent measurements made following irradiation times of 5, 17, 42, 82, 142 and 217 min, respectively. The absorption at  $\sim 495$  nm increases, at the expense of the  $\sim 600$ -nm peak, with the irradiation time up to almost 217 min. The system appears to reach a photostationary state by this time, since further irradiation did not have any effect on the pigment composition.




FIGURE 2 Resonance Raman spectra of: (a) Pink membrane sample, obtained from papain-treated blue membrane (see text), was excited with  $\sim 5$  mW of 476-nm line from an  $\text{Ar}^+$  laser at 80 K; (b) Papain-treated blue membrane was excited with  $\sim 5$  mW of 488-nm light from an  $\text{Ar}^+$  laser, also at 80 K. Slightly higher wavelength was chosen here to improve the spectral quality (see text); (c) Protonated Schiff base of 9-*cis* retinal at  $\sim 10^\circ\text{C}$  using the flow technique (see text). The sample was excited with  $\sim 2$  mW of 476-nm light.

the retinal vibrational modes in the 900 to  $1,300\text{ cm}^{-1}$  region, relative to the ethylenic mode, when the chromophore is present in the protein matrix, compared to when it is in solution. The salient features of these spectra, as well as their implications, are discussed in detail in the sections that follow.

### Ethylenic Region ( $\text{C}=\text{C}$ )

The dominant ethylenic band in the pink membrane spectrum is broad, and is centered around  $1,542\text{ cm}^{-1}$ . The absorption maximum of the pink membrane, based on the observed correlation between the absorption maximum and the ethylenic peak position in the retinal based pigments (Aton et al., 1977), is expected to be at  $\sim 508\text{ nm}$ , which is in reasonable agreement with the observed value of  $\sim 495\text{ nm}$  (Fig. 1). The  $\sim 50\text{ cm}^{-1}$  width of the ethylenic peak suggests that the pink membrane may represent a heterogeneous pigment population, rather than a pure species. For example, the width (fwhm) of this band is  $\sim 25\text{ cm}^{-1}$  for the dark-adapted bR, which contains an equimolar mixture of the 13-*cis* and the all-*trans* isomeric pigments, compared to the  $\sim 14\text{ cm}^{-1}$  band-width of this mode for the light-adapted bR, which contains only the all-*trans* pigment form. The slight hump near  $\sim 1,565\text{ cm}^{-1}$

could not be due to the residual blue membrane in the sample, since the dominant ethylenic peak of the blue membrane lies at  $\sim 1,529\text{ cm}^{-1}$  (Fig. 2 b). Moreover, the contribution of any blue membrane in the sample to the Raman spectrum of the pink membrane, at this excitation wavelength, would be much smaller than its actual concentration, due to the smaller resonance enhancement factors for the blue species ( $\lambda \sim 603\text{ nm}$ ). The contribution from the free retinals can also be ruled out because their ethylenic mode is expected to be at  $1,580\text{ cm}^{-1}$  or higher (Aton et al., 1977). According to the well known correlation between the  $\lambda_{\text{max}}$  and the predominant  $\text{C}=\text{C}$  stretching mode in the retinal systems, noted earlier, the absorption maximum of a species having the main ethylenic band at  $1,565\text{ cm}^{-1}$  is expected to be around  $435\text{ nm}$ . However, no such species is apparent in the absorption spectrum (Fig. 1), which suggests that if this, indeed, results from the contribution from such a species, its concentration either has to be small, or it is produced in small quantities as a result of a rapid photostationary equilibrium upon irradiation at 80 K. It should be pointed out that a species with a similar absorption maximum ( $\sim 450\text{ nm}$ ) and containing predominantly a 9-*cis* chromophore was observed when the pH of the pink membrane, photochemically generated from the acid-blue membrane, was raised above 6 (Fischer et al., 1981). Since the acid-blue membrane seems operationally indistinguishable from the deionized-blue membrane, the presence of a small amount of this  $\sim 450\text{ nm}$  absorbing species in the corresponding deionized pink membrane at pH 4.8 is not inconceivable. Another possibility for the broad ethylenic peak is that two or more  $\text{C}=\text{C}$  modes have unusual Raman intensity. Retinal has five possible  $\text{C}=\text{C}$  vibrational modes, only few of these are intense enough to be normally seen in the resonance Raman spectra of the pigments (see e.g., Callender and Honig, 1977, for a discussion). Thus, the observed shoulder at  $\sim 1,565\text{ cm}^{-1}$  could very well result from one of these modes becoming intense upon the formation of the pink membrane. Another, rather intense, band appears at  $1,594\text{ cm}^{-1}$ ; a similar band at  $\sim 1,600\text{ cm}^{-1}$  is also seen in the Raman spectrum of the 9-*cis* visual pigment, isorhodopsin (see e.g., Oseroff and Callender, 1974; Mathies et al., 1977). A relatively strong band at  $\sim 1,603\text{ cm}^{-1}$  is also shown by the 9-*cis* protonated Schiff base of retinal (Fig. 2 c), while it is absent in the all-*trans* and the 13-*cis* isomeric forms (Pande et al., 1981).

### Schiff Base Region ( $\text{C}=\text{N}$ )

The Schiff base peak in the pink membrane spectrum is fairly well defined, and is located at  $1,652\text{ cm}^{-1}$  (Fig. 2 a). Deuteration of the sample (data not shown) results in a  $30\text{ cm}^{-1}$  shift in this peak to  $1,622\text{ cm}^{-1}$ . For comparison, the Schiff base peak for the blue membrane and the native bR lie at  $\sim 1,639\text{ cm}^{-1}$  and  $1,641\text{ cm}^{-1}$ , respectively, (Fig. 2 b; Pande et al., 1985), whereas the corresponding values for the protonated Schiff bases of retinals all lie at  $\sim 1,661$

$\text{cm}^{-1}$  (Fig. 2 c; Pande et al., 1981). Furthermore, the shifts in the Schiff base mode upon deuteration are  $\sim 17 \text{ cm}^{-1}$  for bR and  $\sim 28 \text{ cm}^{-1}$  for the protonated Schiff bases of retinal. The Schiff base peak position, as well as its shift upon deuteration, appear to be very sensitive to the Schiff base environment (Schiffmiller et al., 1985). Although the blue membrane spectrum has not been measured in  $\text{D}_2\text{O}$  due to its extreme lability, comparison of the available data suggests that the Schiff base environment of the native bR changes somewhat upon formation of the blue membrane, but changes even more when the blue membrane converts to the pink species. Besides causing a shift in the Schiff base position, deuteration of the sample also results in some other changes in the Raman spectrum, which we do not understand at the present time.

### Finger-Print Region ( $1,100\text{--}1,350 \text{ cm}^{-1}$ )

The finger-print region of the pink membrane spectrum shows marked differences from that of the blue membrane (compare Figs. 2 a and 2 b). This region of the spectrum is very sensitive to the chromophore structure, which suggests that the conversion of the blue to the pink membrane is associated with significant changes in the chromophore structure. The peak at  $1,144 \text{ cm}^{-1}$  in the pink membrane spectrum is absent in the spectra of both the blue and the purple membranes (Pande et al., 1985). Furthermore, the absence of this band in the 13-*cis* and the all-*trans* protonated Schiff bases of retinal (Pande et al., 1981), and its presence at  $1,142 \text{ cm}^{-1}$  in the 9-*cis* form (Fig. 2 c), suggests that this band is characteristic of the 9-*cis* chromophore. These data suggest that the pink membrane contains the chromophore in the 9-*cis* isomeric form. It is interesting to note that a similar band appears at  $\sim 1,153 \text{ cm}^{-1}$  in bovine isorhodopsin (Oseroff and Callender, 1974; Mathies et al., 1977), which is also a 9-*cis* pigment.

Our chromophore extraction data reveals that the pink membrane contains predominantly ( $\sim 77\%$ ) a 9-*cis* isomeric pigment, in agreement with our Raman data. Two minor components, 15% all-*trans*, and 8% 13-*cis*, were also observed in the extracted mixture. This isomeric composition is similar to that of the pink membrane obtained by irradiation of the acid-blue membrane (Maeda et al., 1980; 1981; Fischer et al., 1981). The all-*trans* and 13-*cis* isomers are in a proportion, 65:35, very close to what we have determined for the blue membrane (Pande et al., 1985) and these isomers represent  $\sim 23\%$  of the total retinals present in our pink membrane sample. Since this is close to the amount of the  $\sim 600 \text{ nm}$  absorbing species we estimate from Fig. 1, we tentatively assume that the *trans* and the 13-*cis* isomers represent unconverted blue membrane. Thus, the 495 nm species probably contains only 9-*cis* retinal.

While bovine isorhodopsin can be regenerated from the 9-*cis* retinal and opsin, no pigment is formed with the 9-*cis* retinal using bacterio-opsin (Schreckenbach et al., 1978). Our ability to photochemically generate a 9-*cis* pigment

from the deionized bR shows that ions play a pivotal role in governing the photochemistry of these pigments, presumably exerting their influence on the energetics of the protein structural dynamics.

It is interesting to note that a 9-*cis* pigment with an absorption maximum similar to that of the pink membrane reported here, can also be generated photochemically from the acid-blue membrane (Maeda et al., 1980; Fischer et al., 1981). A similarity in the photochemistry of the deionized- and the acid-blue membranes is not surprising, since they have similar isomeric composition, as well as absorption and Raman spectra (Kimura et al., 1984; Chang et al., 1985; Smith et al., 1985; Pande et al., 1985).

A simplistic picture based on the observed results would suggest that the formation of the blue membrane, either acid or deionized, results in a change in the chromophore binding pocket of the protein, that could now accommodate (or stabilize) a structure such as the 9-*cis* isomer, compared to the more elongated all-*trans* or 13-*cis* form. Thus, the transition state leading to the 9-*cis* pigment would then be much lower in energy, resulting in a feasible pathway to the 9-*cis* pigment.

We would like to thank Drs. W. E. Scott and P. Sorter of the Hoffman La Roche Inc., for their generous gift of the 9-*cis* retinal. We also thank Burr Nelson for his technical assistance. Support of this project by the research grants from the National Institutes of Health (EYO 3142), the National Science Foundation (82-01924) and the Department of Energy (02-82 ER 12087) is gratefully acknowledged.

Received for publication 2 December 1985 and in final form 3 March 1986.

### REFERENCES

- Aton, B., A. G. Doukas, R. H. Callender, B. Becher, and T. G. Ebrey. 1977. Resonance Raman studies of the purple membrane. *Biochemistry*. 16:2995-2999.
- Callender, R., and B. Honig. 1977. Resonance Raman studies of visual pigments. *Annu. Rev. Biophys. Bioeng.* 6:33-55.
- Chang, C.-H., J. G. Chen, R. Govindjee, and T. Ebrey. 1985. Cation binding by bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA*. 82:396-400.
- Chen, J. G., C.-H. Chang, R. Govindjee, and T. Ebrey. 1984. Bacteriorhodopsin's purple color is partially due to divalent cations. *Biophys. J.* 45 (2, Pt. 2):214a. (Abstr.).
- Ebrey, T. 1982. Light Energy Transduction in *H. Halobium*. In *Membrane and Transport*. A. Martonosi, editor. Plenum Publishing Corp., New York. 323-330.
- Fischer, U. C., P. Towner, and D. Oesterhelt. 1981. Light induced isomerization, at acidic pH, initiates hydrolysis of bacteriorhodopsin to bacterio-opsin and 9-*cis* retinal. *Photochem. Photobiol.* 33:529-537.
- Kimura, Y., A. Ikegami, and W. Stoeckenius. 1984. Salt and pH-dependent changes of the purple membrane absorption spectrum. *Photochem. Photobiol.* 40:641-646.
- Maeda, A., T. Iwasa, and T. Yoshizawa. 1980. Formation of 9-*cis*- and 11-*cis*-retinal pigments from bacteriorhodopsin by irradiating purple membrane in acid. *Biochemistry*. 19:3825-3831.
- Maeda, A., T. Iwasa, and T. Yoshizawa. 1981. Photoreaction of the acidified form of bacteriorhodopsin and its 9-*cis* derivative in purple membrane at low temperatures. *Photochem. Photobiol.* 33:559-565.
- Mathies, R., T. B. Freedman, and L. Stryer. 1977. Resonance Raman studies of the conformation of retinal in rhodopsin and isorhodopsin. *J. Mol. Biol.* 109:367-372.

- Oseroff, A. R., and R. Callender. 1974. Resonance Raman spectroscopy of rhodopsin in retinal disk membranes. *Biochemistry*. 13:4243-4248.
- Pande, C., R. Callender, C.-H. Chang, and T. Ebrey. 1985. Resonance Raman spectra of the blue and the regenerated purple membranes of *H. halobium*. *Photochem. Photobiol.* 42:545-548.
- Pande, J., R. Callender, and T. Ebrey. 1981. Resonance Raman study of the primary photochemistry of bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA*. 8:7379-7382.
- Schiffmiller, R., R. H. Callender, W. H. Waddell, R. Govindjee, T. G. Ebrey, H. Kakitani, B. Honig, and K. Nakanishi. 1985. Resonance Raman studies of bacteriorhodopsin analogues. *Photochem. Photobiol.* 41:563-567.
- Schreckenbach, T., B. Walckhoff, and D. Oesterhelt. 1978. Specificity of the retinal binding site of bacteriorhodopsin: chemical and stereochemical requirements for the binding of retinol and retinal. *Biochemistry*. 17:5353-5359.
- Smith, S. O., and R. Mathies. 1985. Resonance Raman spectra of the acidified and deionized forms of bacteriorhodopsin. *Biophys. J.* 47:251-254.
- Stoeckenius W., and R. Bogomolni. 1982. Bacteriorhodopsin and related pigments of halobacteria. *Annu. Rev. Biochem.* 52:587-616.