

Here, initial investigations on the interaction of benzo[b]fluorenone (BF) with the human telomeric forming oligonucleotide HT4 using circular dichroism (CD), thermal melting and steady-state fluorescence are reported. CD studies suggest that BF exhibits binding selectivity for G-quadruplex over model double-stranded DNA structures. BF binds to hybrid-mixed type quadruplex structures in the presence of physiological concentrations of  $K^+$  ions (100mM). However, even in the absence of salt, BF can induce an anti-parallel type quadruplex conformation. Thermal melting studies on quadruplex HT4 in the presence and absence of salt show BF has little effect on the melting temperature, suggesting that BF may associate with the G-quadruplex structure through a non-intercalating mode. As BF is known to be a solvatochromic fluorophore, steady state fluorescence emission spectra were taken for BF titrated with the HT4 quadruplex formed at 100mM  $K^+$ , to explore the ability of BF to act as a potential probe of the environment of its binding site. Based on the red shift of BF emission, BF/quadruplex association appears to involve a relatively polar and hydrogen-bonded environment. This work was supported by NIH SCORE Grant S06 GM 060654.

### 1387-Pos

#### Which Comes First, The Deformation or the Binding?

**Sungchul Hohng, Sangsu Bae.**

Seoul National University, Seoul, Republic of Korea.

Which comes first, the deformation or the binding?

High specificity of protein-DNA interaction is often related with specific deformation of the binding site. B-Z transition is the most dramatic structural change induced by protein-DNA interaction, where some segment of DNA abruptly changes from the right-handed B-DNA to the left-handed Z-DNA by the help of specific proteins.

Here, we report single-molecule FRET studies on protein-induced Z-DNA formation. DNA duplexes with six CG-repeats were prepared. To monitor the conformational dynamics of the CG-repeat, we labeled Cy3 and Cy5 at each end of the CG-repeat. Surface-immobilized DNA molecules did not show any structural dynamics in normal physiological conditions. When a Z-DNA inducing protein,  $\alpha$ , was added to the buffer solution, however, fluorescence intensity increased abruptly without any accompanying FRET change. Abrupt FRET change occurred with time delay ( $\sim 10$  minute at  $25^\circ\text{C}$  on average). When the proteins were washed out, molecules didn't recover the original FRET value for more than 3 hours, but molecules without the FRET change readily recovered their original fluorescence intensity. From these result, we conclude that  $\alpha$  protein weakly interact with B-DNA, but the interaction becomes extremely strong once Z-DNA is formed.

Next, we prepared a DNA duplex with methylated cytosine in the CG repeat. With millimolar  $Ni^{2+}$  in the buffer solution, we observed the intrinsic B-Z transition dynamics, and Z-DNA stabilization by  $\alpha$  proteins. The transition time from B-DNA to Z-DNA, however, was not affected by the presence of  $\alpha$  proteins, which strongly support that  $\alpha$  protein induces Z-DNA by passively trapping Z-DNA structure transiently formed by the intrinsic B-Z transition dynamics. Even though we cannot directly observe Z-DNA, Z-DNA's are actually waiting there inside the cell to play their biological roles on time.

### 1388-Pos

#### Monitoring Structural Transitions of a DNA Holliday Junction Using an Acoustic Wave Sensor


**Achilleas Tsortos.**

FORTH, Heraklion-Crete, Greece.

Structural DNA nanotechnology deals with building DNA molecules with particular geometrical features; creating nanoscale shapes and patterns using long DNA scaffolds, as in DNA origami, or changing DNA shape in a controlled way and in response to an external stimulus, as in molecular switches. Such nano-machines can perform computation, actuation and diagnostic tasks.

We took advantage of a biomolecular structure comprising a four-way DNA (Holliday) junction which belongs to a molecular machines group capable of moving between distinct states. This particular molecular switch exists in either an open (extended) or closed (coaxially stacked) conformation and it is proposed to be used as a principle for sequence-specific nucleic acid recognition. In this study we used the quartz crystal microbalance with energy dissipation (QCM-D) to study the applicability of this structure to detect DNA hybridization on a device surface and test its potential to act as a controllable switch.

We present a novel way of monitoring in real time both oligonucleotide binding and the transition from the closed to the open state of the junction. This transition can be reversed and repeated indefinitely in a fully controllable way.



### 1389-Pos

#### Single Molecule FRET Measures Structure and Fast Dynamics of DNA and RNA Four-Way Junctions

**Claus A.M. Seidel, Stanislav Kalinin, Suren Felekyan, Alessandro Valeri, Simon Sindbert.**

Heinrich-Heine University Duesseldorf, Duesseldorf, Germany.

Using a confocal fluorescence microscope multiparameter fluorescence detection (MFD) enables us to simultaneously collect all fluorescence information such as intensity, lifetime, anisotropy in several spectral ranges) from picoseconds to seconds. MFD and fluorescence correlation spectroscopy is applied to perform single-molecule FRET studies with an ultimate level of precision in determining separations with FRET of 1% of the Förster radius [J.Phys.Chem.B 110, 6970 (2006), J.Phys.Chem.B 112, 8361 (2008)]. In addition we can unambiguously distinguish between stochastic processes and broadening due to static or dynamic heterogeneity. In this way we measured bends and kinks in dsDNA. The high accuracy allowed us the detection of sequence-dependent DNA bending by  $16^\circ$  [PNAS 105, 18773 (2008)]. Moreover we studied the Mg-dependent structural dynamics of a four-way DNA (Holliday-) junction in order to find out whether the postulated extended square structure accumulates indeed as an intermediate or whether it should be considered more as a very short lived transition state. We found a complex Mg dependent kinetics, which must be described by a four species model with only two distinct FRET and two kinetic levels. The species with the same FRET value differ in their conformational flexibility: one is quasi-static, the other is dynamic. Our FRET data are clearly inconsistent with an accumulation of a single extended square junction structure at very low Mg concentrations. Finally we compare the structure and dynamics of DNA- and RNA four-way junctions. Thereby the structure of RNA four-way junction was characterized by 24 FRET distances, which allowed us to prove the existence of 3 of the 4 possible stacking conformers. These studies show that sm FRET studies are valuable tool to complement the structural and dynamic information obtained by X-ray crystallography or NMR spectroscopy.

## Protein-Nucleic Acid Interactions II

### 1390-Pos

#### Kinetic Enhancement of NF- $\kappa$ B/DNA Dissociation by I $\kappa$ B $\alpha$

**Vera Alverdi, Byron Hetrick, Simpson Joseph, Elizabeth A. Komives.**

UCSD, La Jolla, CA, USA.

The nuclear factor kappa B (NF- $\kappa$ B) family of transcription factors is involved in inter- and intracellular signaling, cellular stress response, growth, survival, and apoptosis. Specific inhibitors of NF- $\kappa$ B transcription including I $\kappa$ B $\alpha$ , I $\kappa$ B $\beta$ , and I $\kappa$ B $\epsilon$ , block the transcriptional activity of p65 and c-Rel-containing NF- $\kappa$ B dimers. DNA binding by NF- $\kappa$ B is inhibited by the ankyrin repeat protein kappa B (I $\kappa$ B $\alpha$ ), which sequesters NF- $\kappa$ B to the cytosol. The mechanism and kinetics of DNA binding inhibition by I $\kappa$ B $\alpha$  are still unknown, but we recently demonstrated that NF- $\kappa$ B can be "stripped" off DNA by I $\kappa$ B $\alpha$ . We are investigating the effect of I $\kappa$ B $\alpha$  on the association and dissociation rates of NF- $\kappa$ B/DNA complex formation using titration measurements, stop flow fluorescence and ITC. We are using pyrene labeled DNA or I $\kappa$ B $\alpha$  or NF- $\kappa$ B to study the fluorescence changes occurring during the NF- $\kappa$ B "stripping". Our results show that I $\kappa$ B $\alpha$  increases the dissociation rate of the DNA from the NF- $\kappa$ B complex in a concentration-dependent manner and with high efficiency. We are studying also I $\kappa$ B $\beta$  which appears to stabilize the NF- $\kappa$ B/DNA interaction. This could suggest the formation of a ternary complex DNA/NF- $\kappa$ B/I $\kappa$ B $\beta$ .

### 1391-Pos

#### Prion Aptamer, Free and Bound States

**Abigail Miller, Lori Goldner.**

NIST, Gaithersburg, MD, USA.

Aptamers are short single strands of DNA or RNA that bind to proteins, peptides, and small molecules. They are likely to fold into different structures when free in solution than when they are bound to a molecular target. The free structures are difficult to determine experimentally, though they can be modeled by calculating the minimum thermodynamic states. We test the validity of the thermodynamic models of a prion aptamer using single molecule pair Forster resonance energy transfer (spFRET) as a structural reporter. The FRET states of the unbound aptamers, the hybridized aptamers and the aptamers bound to PrP peptides are characterized. The DNA aptamers to PrP has a pair of thermodynamic states of roughly the same energy at  $25^\circ\text{C}$ . Their presence in solution is characterized by comparing the single stranded aptamers to its hybridized configuration, thereby removing any internal structure of the aptamers. We demonstrate the existence of both unbound thermodynamic states as well as different interactions between the aptamer and each PrP peptide from static data measuring spFRET in solution.