

mutant (R108A) were performed to investigate the diffusion characteristics of these 'inactive' and 'active' states of the A₃-receptor. One slow moving complex was identified at the cell membrane of wild-type A₃-GFP transfected cells, with a diffusion co-efficient (0.087 μm²/s) similar to that of τ_{D3} for the XAC-X-BY630; similar complexes were identified in the mutant A₃-receptor cell lines. We have subsequently used FCS in conjunction with fluorescent agonist and antagonist A₃-receptor ligands to compare the ligand binding and diffusion properties of these different activity states of the receptor at the subcellular level.

1514-Pos

Computational Insight into the Ligand-Induced Conformational Specificity of G-Protein Coupled Receptors

Davide Provasi, Juan Carlos Mobarec, Marta Camacho Artacho, Marta Filizola.

Mount Sinai School of Medicine, New York, NY, USA.

Several observations in the G-protein coupled receptor (GPCR) literature support the existence of ligand-specific intermediate conformational states that are likely to be involved in differential activation of signaling pathways. Fluorescence spectroscopy studies provide direct evidence for ligand-specific receptor conformations of the β₂-adrenergic receptor, making this system an attractive target to test the ability of computational methodologies to predict different activated states of GPCRs. To this end, we designed a computational strategy that combines adiabatic biased molecular dynamics (ABMD) and metadynamics simulations. Firstly, ABMD is used to generate transition paths between the experimental inactive crystal structure of the β₂-adrenergic receptor and a conformation containing established features of activated states of GPCRs (modeled using the opsin crystal structures). Secondly, metadynamics is applied to study how ligands with different efficacies affect the free-energy of different metastable states identified along these putative activation pathways. The calculated free-energy profiles of the different ligand-β₂ adrenoceptor complexes help rationalize the published experimental results, including the different kinetics of catecholaminergic agonists such as epinephrine, norepinephrine, dopamine, and isoproterenol. Representative structures of the identified energy basins suggest specific residues and contacts that may help stabilize different activated states of the receptor. This information holds promise for the crystallization of different GPCR conformations.

1515-Pos

Structural and Kinetic Modeling of an Activating Helix Switch in the Rhodopsin-Transducin Interface

Peter W. Hildebrand.

Uni Berlin Charite, Berlin, Germany.

Extracellular signals prompt G protein-coupled receptors (GPCRs) to adopt an active conformation (R*) and to catalyze GDP/GTP exchange in the α-subunit of intracellular G proteins (Gαβγ). Kinetic analysis of transducin (G_tαβγ) activation shows that an intermediary R*G_tαβγGDP complex is formed which precedes GDP release and formation of the nucleotide-free R*G protein complex. Based on this reaction sequence we explore the dynamic interface between the proteins during formation of these complexes. We start from the R* conformation stabilized by a G_tα C-terminal peptide (GαCT) obtained from crystal structures of the GPCR opsin. Molecular modeling allows reconstruction of the fully elongated C-terminal α-helix of G_tα (α5) and shows how α5 can be docked to the open binding site of R*. Two modes of interaction are found. One of them - termed stable or S-interaction - matches the position of the GαCT peptide in the crystal structure and reproduces the hydrogen bridge networks between the C-terminal reverse turn of GαCT and conserved E(D)RY and NPxxY(x)_{5,6}F regions of the GPCR. The alternative fit - termed intermediary or I-interaction - is distinguished by a tilt (42°) and rotation (90°) of α5 relative to the S-interaction. It shows different α5 contacts with the NPxxY(x)_{5,6}F region and the second cytoplasmic loop of R*. From the two α5 interactions, we derive a 'helix switch' mechanism for the transition of R*G_tαβγGDP to the nucleotide-free R*G protein complex. It illustrates how α5 might act as a transmission rod to propagate the conformational change from the receptor-G protein interface to the nucleotide binding site.

1516-Pos

Agonist-Specific Effects of TM V Serine Mutations on Dopamine D2S Receptor Voltage-Sensitivity

Kristoffer Sahlholm, Daniel Marcellino, Johanna Nilsson, Kjell Fuxe, Peter Århem.

Karolinska Institutet, Stockholm, Sweden.

Voltage-sensitivity has recently been demonstrated for agonist potency and affinity at certain G protein-coupled receptors. Using an electrophysiology

assay in *Xenopus* oocytes, we have previously shown that the potency of dopamine in activating G protein-coupled potassium channels (GIRK) via the dopamine D2S receptor is reduced by depolarization from -80 to 0 mV. We recently investigated the voltage-sensitivities of a range of structurally related dopaminergic agonists at the D2S receptor.

The findings of this study led us to propose that a conformationally constrained interaction of the agonist with transmembrane segment (TM) VI of D2 is required for voltage-sensitivity. The hypothesis assumes that for the flexible phenethylamines, two hydroxyls (such as in dopamine) interacting with the conserved serines in TM V are necessary for voltage-sensitivity. Conversely, N,N-dipropyl-2-aminotetralin (DPAT) agonists do not require hydroxyls for voltage-sensitivity due to their inherently more rigid structure. To test this hypothesis, we mutated three conserved serines in TM V (S193A, S194A, and S197A) which have been shown to mediate binding to agonist hydroxyls. The voltage-sensitivity of non-hydroxylated DPAT was similar to that observed with the wild-type receptor at all of the three mutants, suggesting that the mutations did not allosterically alter the voltage-sensing properties of the receptor.

The S193A mutation drastically diminished voltage-sensitivity of dopamine, concomitantly with a marked reduction in potency. However, the S194A mutation which slightly decreased potency, did not appreciably affect the voltage-sensitivity of dopamine. At the S197A mutant, dopamine efficacy was decreased to such a degree that voltage-sensitivity could not be assessed.

In the literature, S193 has consistently been assigned a major role in dopamine binding. Our results suggest that this residue might also be important for voltage-sensitive interactions between dopamine and the D2S receptor.

1517-Pos

Influence of Membrane Composition on Function of Human Peripheral Cannabinoid Receptor CB2

Tomohiro Kimura, Alexei A. Yeliseev, Krishna Vukoti, Klaus Gawrisch. NIH/NIAAA, Bethesda, MD, USA.

The human peripheral cannabinoid receptor CB2 is abundant in tissues of immune and hematopoietic systems. CB2 belongs to the class of heptahelical G-protein coupled receptors and regulates a wide range of physiological functions through binding of endogenous and exogenous cannabinoid ligands. We studied the influence of electrical surface potential of membranes and of hydrocarbon-chain order on rates of G-protein activation by CB2. The membrane surface potential was determined by a measurement of the electrophoretic mobility of proteoliposomes, while lipid hydrocarbon-chain order was quantified by a measurement of the order parameters using ²H NMR. The receptor, expressed in *E. coli*, was purified and functionally reconstituted into lipid bilayers composed of phosphatidylcholine (PC), phosphatidylserine (PS), and cholesteryl hemisuccinate (CHS). CB2 was fully activated with the synthetic agonist CP-55,940. The rate of G-protein activation by the receptor increased about two-fold with increasing CHS content in the lipid matrix from 25 to 41 mol%. Similar effect was observed with increasing PS content. The increased activation rate correlated with a larger negative ζ-potential caused by the negatively charged headgroups. The increased order of lipid acyl chains due to interactions with the cholesteryl backbone of CHS had no significant effect on G-protein activation rates, as confirmed by addition of cholesterol instead of CHS. The results indicate the importance of anionic lipids for efficient coupling between the CB2 receptor and G-proteins.

1518-Pos

A Polybasic Region in the C-terminus of M3 Muscarinic Acetylcholine Receptors Mediates an Interaction with Gq Heterotrimers

Kou Qin, Sudha Kuravi, Nevin Lambert.

Medical College of Georgia, Augusta, GA, USA.

G protein-coupled receptors (GPCRs) form stable complexes with heterotrimeric G proteins when the former are activated and when the latter are not bound to guanine nucleotides. In addition to these active-state ternary (ligand-receptor-G protein) complexes some GPCRs have been suggested to form preassembled or precoupled complexes with G proteins prior to activation. We have previously reported that immobile M3 muscarinic receptors (M3Rs) decrease the mobility of heterotrimers that contain Gαq, consistent with an M3R-Gq complex. This interaction is unaffected by receptor ligands in intact cells, and is specific for M3Rs and Gq, as immobile M4Rs do not decrease the mobility of Gq heterotrimers, and immobile M3Rs do not decrease the mobility of GoA heterotrimers. In order to determine the structural basis of this interaction, we constructed a series of CFP-labeled M3R/M4R chimeras and tested their ability to decrease the mobility of venus-labeled Gq (Gq-V) using fluorescence recovery after photobleaching (FRAP). A chimera consisting of the M3R with the c-terminus of the M4R (M3M4ct) did not decrease Gq-V mobility. A polybasic region