

2141-Plat**Troponin Regulatory Function and Dynamics Revealed by H/D Exchange-Mass Spectrometry**

Devanand Kowlessur, Larry S. Tobacman.

University of Illinois at Chicago, Chicago, IL, USA.

Troponin is the thin filament protein that confers tight, Ca^{2+} -dependent control over muscle contraction. The mechanism of this regulation was investigated by detailed mapping of the dynamic properties of cardiac troponin, using amide hydrogen exchange-mass spectrometry, in the presence of either saturation or non-saturation of the regulatory Ca^{2+} binding site in the NH_2 -domain of subunit TnC. Troponin was found to be highly dynamic, with 60% of amides exchanging H for D within seconds of exposure to D_2O . In contrast, portions of the TnT-TnI coiled-coil exhibited high protection from exchange, more than six hours, identifying the most stable portion of the trimeric troponin complex. Regulatory site Ca^{2+} binding altered dynamic properties (i.e., H/D exchange protection) locally, near the binding site and in the TnI switch helix that attaches to the Ca^{2+} -saturated TnC NH_2 -domain. More notably, Ca^{2+} also altered the dynamic properties of other parts of troponin: the TnI inhibitory peptide region that binds to actin, the TnT-TnI coiled-coil, and the TnC COOH-domain that contains the regulatory Ca^{2+} sites in many invertebrate as opposed to vertebrate troponins. Mapping of these affected regions onto troponin's highly extended structure indicates contacts important in conformational change: in the low Ca^{2+} state the TnI region that effects inhibition bends back and interacts with the end of the TnT-TnI coiled-coil, as previously suggested by intermediate resolution X-ray data of skeletal muscle troponin. Thus, troponin-mediated Ca^{2+} sensitive regulation of muscle contraction consists of Ca^{2+} -triggered switching between alternative sets of intra-troponin interactions.

2142-Plat**Magnesium Stabilizes the Closed Conformation of the C-Domain of Troponin C**

Sarah S. Learman, Rishabh Khandelwal, Franklin Fuchs, Zenon Grabarek.

Boston Biomedical Research Institute, Watertown, MA, USA.

Activation of the thin filament in striated muscles requires both the binding of Ca^{2+} to the N-domain of troponin C (TnC) and the binding of myosin cross-bridges to actin, which has been shown to alter the C-domain conformation. Here we have evaluated the structural and functional consequences of divalent cation exchange in skeletal and cardiac TnC (sTnC and cTnC). We have used intrinsic tyrosine fluorescence, circular dichroism (CD), and the fluorescent nonspecific hydrophobic probe bis-ANS to monitor changes in domain conformation in response to Ca^{2+} and Mg^{2+} binding in the sTnC, cTnC, and in a cTnC mutant in which the invariant Glu residue at the 12th position of the calcium binding loops III and IV were substituted with Asp (cTnCDD). Ca^{2+} binding causes an increase in Tyr fluorescence and α -helical content in sTnC and cTnC, but not in cTnCDD. Ca^{2+} induced C-domain opening characteristic of sTnC and cTnC was also greatly reduced in cTnCDD, as measured by bis-ANS fluorescence. Thus the Asp to Glu substitutions appear to prevent the C-domain from opening. Bis-ANS Ca^{2+} titrations also showed that high Ca^{2+} concentrations may be sufficient to open the N-domain of cTnC, which was reported to remain in the closed conformation in the Ca^{2+} -bound state. Lastly, bis-ANS Mg^{2+} titrations indicate that Mg^{2+} does not cause domain opening in either cTnC or cTnCDD. The closed conformation of the Mg^{2+} -bound C-domain of TnC implies a different mechanism of interaction with TnI than that in the presence of Ca^{2+} and suggests that the Mg^{2+} - Ca^{2+} exchange in TnC may contribute to the thin filament activation of muscle contraction. This conclusion is consistent with our observation that physiological concentrations of Mg^{2+} significantly lower the Ca^{2+} -sensitivity of reconstituted cardiac thin filaments.

2143-Plat**Phosphomimetic Substitutions in One or Both Ser43/45 Residues of Cardiac Troponin I Produces Comparable Changes in Contractile Performance**

Tamara Stevenson, Sarah Kampert, Justin Steggerda, Gail Romanchuk, Margaret Westfall.

University of Michigan, Ann Arbor, MI, USA.

Cardiac troponin I (cTnI) is phosphorylated on three clusters of residues in response to protein kinase C (PKC) activation. Previously, studies on the cTnSer43/45 cluster showed phosphomimetic Asp substitution reduced peak shortening and accelerated re-lengthening in adult cardiac myocytes. The goal of the present study is to determine whether one or both Ser residues contribute to the functional response observed with cTnSer43/45Asp. We studied adult rat cardiac myocytes 2 and 4 days after viral-mediated gene transfer of

cTnIFLAG, cTnSer43Asp or cTnSer45Asp (+FLAG). Western analysis indicated similar levels of cTnI replacement developed for all groups, and extensive replacement with cTnIFLAG ($71 \pm 9\%$, $n=6$), and FLAG-tagged epitopes of cTnIS43D ($72 \pm 3\%$, $n=8$) and cTnIS45D ($70 \pm 5\%$, $n=8$) within 4 days. Further analysis showed no significant change in cTnI stoichiometry and confocal analysis confirmed a sarcomeric incorporation pattern for each mutant. In functional studies, shortening amplitude decreased significantly in chronically paced myocytes expressing non-tagged Ser43Asp and/or Ser45Asp compared to controls (Control = $0.149 \pm 0.008 \mu\text{m}$, $n=36$; cTnSer43/45Asp = $0.110 \pm 0.006 \mu\text{m}$; $n=32$ *; cTnSer43Asp = 0.095 ± 0.007 , $n=44$ *; cTnSer45Asp = 0.108 ± 0.007 , $n=50$; * $p < 0.05$ vs control) 4 days after gene transfer. An accelerated re-lengthening accompanied this reduced shortening (Time to 75% relaxation = $\text{TTR}_{75\%}$ (ms): Control = 79 ± 4 ; cTnSer43/45Asp = 62 ± 4 *; cTnSer43Asp = 63 ± 4 *, cTnSer45Asp = 65 ± 3 *; * $p < 0.05$ vs control). Interestingly, each single mutant also accelerated the time to peak shortening (TTP (ms): Control = 83 ± 5 ; cTnSer43Asp = 68 ± 3 *; cTnSer45Asp = 67 ± 2 *; * $p < 0.05$ vs control) while cTnSer43/45Asp did not (84 ± 5). These initial results provide evidence that each Ser residue in the Ser43/45 cluster is capable of altering cTnI function in response to phosphorylation by PKC, yet phosphorylation of both residues does not produce an additive response.

2144-Plat**Effect of Troponin Ca^{2+} Binding Properties on the Kinetics of Myofibril Force Initiation and Relaxation**Nicoletta Piroddi¹, Karen L. Kreutziger², Beatrice Scellini¹, Scott Lundy², Cecilia Ferrantini¹, Chiara Tesi¹, Michael Regnier², Corrado Poggesi¹.¹Università di Firenze, Firenze, Italy, ²University of Washington, Seattle, WA, USA.

We have engineered the Ca^{2+} binding properties of troponin C (TnC) to study the role of increased (I60Q sTnC) and decreased (M80Q sTnC^{F27W}) Ca^{2+} dissociation rate (k_{off}) on activation and relaxation of skeletal muscle. Previously we reported that myofibril force development kinetics (k_{ACT}) are not influenced by decreasing k_{off} from Tn, but are slowed by an increase in k_{off} (Kreutziger et al. 2008 JPhysiol. 586:3683-3700) at low $[\text{P}_i]$ (5 μM). The time to initiation of force ($k_{\text{A}}\text{lag}$) following a rapid (~10ms) switch from pCa 9.0 to pCa 3.5 provides information about thin filament activation rate and our preliminary data suggest this rate may also be sensitive to k_{off} . In rabbit psoas myofibrils (15°C) $k_{\text{A}}\text{lag}$ (~20 ms for native or WT sTnC) is almost eliminated for M80Q sTnC^{F27W} and increased by I60Q sTnC (~40-50 ms). Additionally, though k_{ACT} is similar for force increases from either full or partial activation to full activation, $k_{\text{A}}\text{lag}$ disappears when starting from partial activation. We have also reported that fast and slow phase rates of relaxation are not affected by k_{off} , but that duration of the slow phase is affected in skeletal myofibrils. Here we report that lag prior to initiation of the slow phase ($k_{\text{R}}\text{lag}$) may be also influenced by k_{off} . Opposite to $k_{\text{A}}\text{lag}$, $k_{\text{R}}\text{lag}$ (~20 ms for WT or native sTnC) was increased (~40-50 ms) by decreased k_{off} (M80Q sTnC^{F27W}) and almost eliminated by increased k_{off} (I60Q sTnC). These experiments demonstrate a potential approach to study thin filament activation/deactivation kinetics without the need for fluorescent probes attached to thin filament proteins that can affect their function. Supported by Telethon GGP07133, MIUR (CP, CT), NIH-HL65497 (MR).

2145-Plat**Nebulin Alters Crossbridge Cycling Kinetics and Increases Thin Filament Activation - a Novel Mechanism for Increasing Tension and Reducing Tension Cost**Coen Ottenheijm¹, Murali Chandra², R. Mamidi², Steven Ford², Carlos Hidalgo³, Christian Witt⁴, Ger Stienen¹, Siegfried Labeit⁴, Alan Beggs⁵, Henk Granzier³.¹VU University medical center, Amsterdam, Netherlands, ²Washington StateUniversity, Pullman, WA, USA, ³University of Arizona, Tucson, AZ, USA,⁴University Hospital Mannheim, Mannheim, Germany, ⁵Harvard Medical

School, Boston, MA, USA.

Nebulin is a giant filamentous F-actin binding protein that binds along the thin filament of the skeletal muscle sarcomere. Although nebulin is usually viewed as a structural protein, here we investigated whether nebulin plays a role in muscle contraction by using skinned muscle fiber bundles from a nebulin knockout (NEB KO) mouse model. We measured force-pCa and force-ATPase relations, as well as the rate of tension redevelopment (k_{tr}) in tibialis cranialis fibers. To rule out any alterations in troponin (Tn) isoform expression and/or status of Tn phosphorylation, we studied fibers that had been reconstituted with fast skeletal muscle recombinant Tn. We also performed a detailed analysis of myosin heavy chain, myosin light chain (MLC) and MLC2