

In this study we have addressed the assembly of lipid bilayers in arrays and the stability of established membranes in different scaffold geometries.

To establish planar lipid membranes across large scale partition aperture arrays, we created a disposable single-use horizontal chamber design that supports combined optical-electrical measurements. Lipid bilayers could easily and efficiently be established across CO₂ laser micro structured 8 × 8 aperture partition arrays with average aperture diameters of 301 ± 5 μm.

To demonstrate the functionality of the lipid bilayers established across the 8 × 8 arrays, controllable reconstitution of the biotechnological and physiological relevant peptides valinomycin and gramicidin A, together with the membrane proteins α-Hemolysin and FomA were carried out. The results showed that the design supports low current (high sensitivity) recordings of membrane peptides and proteins by incorporating gramicidin A, α-Hemolysin and FomA into the established lipid bilayers. Finally, we tested the scalability of the assembly of lipid bilayers by creating rectangular 24 × 24 and hexagonal 24 × 27 lipid membrane arrays respectively. The two different geometries of the micro structured aperture arrays seem to support stable and functional membrane arrays, however, with somewhat different electrical properties. We propose that the presented design may be suitable for further developments of sensitive biosensor assays.

2534-Pos

Comparison of the Effects of Cholesterol or 3β-Hydroxy-5-Oxo-5,6-Secosterol-6-ol on the Thermotropic and Structural Properties of Mixtures of Phosphatidylethanolamine and Phosphatidylcholine

Ellen J. Wachtel¹, Diana Bach¹, Richard M. Epand², Rachel F. Epand², Israel R. Miller¹.

¹Weizmann Institute of Science, Rehovot, Israel, ²McMaster University, Hamilton, ON, Canada.

The oxidation of cholesterol with ozone produces 3β-hydroxy-5-oxo-5,6-secocholestan-6-ol. This oxysterol has been implicated in a number of pathological conditions *in vivo* including atherosclerotic plaque formation and amyloidogenesis. We have shown previously that this oxysterol strongly modifies the physical properties of model membranes composed of different phosphatidylethanolamines or phosphatidylserine. In the present work we have extended our studies to ternary mixtures composed of phosphatidylethanolamine and phosphatidylcholine with sterols, either 3β-hydroxy-5-oxo-5,6-secocholestan-6-ol or cholesterol. We use differential scanning calorimetry and small angle X-ray diffraction to characterize the phase behavior of mixtures of dipalmitoleoylphosphatidylethanolamine (diPoPE) and dipalmitoleoylphosphatidylcholine (diPoPC) or 1-palmitoyl-2-oleoyl-phosphatidylethanolamine (POPE) and 1-palmitoyl-2-oleoyl-phosphatidylcholine (POPC). We compare the effect of the two sterols on the temperature of the transition of the ternary system from the liquid crystalline to the hexagonal phase (T_H) and on the curvature of the resulting cylindrical micelles. Addition of low concentrations of diPoPC increases T_H while adding cholesterol to this mixture significantly lowers T_H. The effect of 3β-hydroxy-5-oxo-5,6-secocholestan-6-ol is much weaker than that of cholesterol. With regard to the curvature of the cylindrical micelles, the addition of diPoPC and 3β-hydroxy-5-oxo-5,6-secocholestan-6-ol have opposing effects, while cholesterol does not effect the curvature at all. Low concentrations of POPC in POPE cause an increase in T_H and decrease the curvature of the cylindrical micelles, while cholesterol has the opposite effect.

2535-Pos

Self-Assembly Simulations of Membranes Containing Phospholipid Oxidation Products

Himanshu Khandelia, Ole G. Mouritsen.

University of Southern Denmark, Odense, Denmark.

Products of phospholipid oxidation (OXPLs) are involved with the genesis or pathology of several diseases. OXPLs can modify the physical properties of biological membranes, thereby possibly altering several biological processes near membranes including signaling pathways. We have used atomistic and coarse grained simulations to investigate the properties of OXPL-containing lipid bilayers. We ran self assembly simulations of mixtures of palmitoyl-oleoyl-phosphatidylcholine (POPC) with two different OXPLs: PazePC, which is anionic, and PoxnoPC, which is zwitterionic. The total sampling time exceeds 1 millisecond. Despite having shortened and polar acyl chains, the two OXPLs POPC self assemble into stable lipid bilayers with POPC. The bilayers can accommodate at least 25% OXPL, although such bilayers have a lower area compressibility modulus. As an example of the modification of a membrane-associated biological process, we show that KALP-23 peptides partition differently in POPC-OXPL and POPC bilayers. The peptides adopt a transmembrane orientation more easily when OXPLs are present in the bilayers.

2536-Pos

Study of the Cholesterol Umbrella Effect in DPPC and DOPC Bilayers by Molecular Dynamics Simulation

Jian Dai, Mohammad Alwarawrah, Juyang Huang.

Texas Tech University, Lubbock, TX, USA.

The instability of cholesterol clusters in dipalmitoylphosphatidylcholine (DPPC) and dioleoylphosphatidylcholine (DOPC) lipid bilayers was investigated via atomistic Molecular Dynamics (MD) simulation. Cholesterol clusters in phosphatidylcholine (PC) bilayers are found to be very unstable and to readily disperse into cholesterol monomers. The instability may result from the difficulty for the system to prevent water exposure to cholesterol's aggregated hydrophobic bodies in a cluster. The system responds to artificially arranged cholesterol clusters in several interesting manners: (i) Cholesterol clusters quickly form a "frustum" shape to reduce water penetration through cholesterol headgroups; (ii) Many clusters bury themselves deeper into the bilayer interior, causing local bilayer deformation; (iii) Cholesterol fluctuates rapidly, both laterally and vertically to the bilayer plane, in order to escape from clusters. These fluctuations result in the disintegration of clusters, and in one incidence, a highly unusual flip-flop event of a cholesterol across the DOPC bilayer occurs. Our results show that cholesterol has a strong tendency to avoid forming clusters in lipid bilayers and that the fundamental cholesterol-cholesterol interaction is unfavorable. Furthermore, the radial distribution functions of cholesterol hydroxyl oxygen to various headgroup atoms of PC reveal that the PC headgroups surrounding cholesterol have a clear tendency to reorient and extend toward cholesterol. The range of this "Umbrella Effect" can reach up to 2-3 nm, larger than previously reported.

2537-Pos

Flip-Flop Motions of Lipid Molecules in Mixed Bilayer Systems

Fumiko Ogushi, Reiko Ishitsuka, Toshihide Kobayashi, Yuji Sugita.

RIKEN advanced science institute, Wako, Japan.

The cell membrane is composed of a wide variety of lipid molecules, cholesterol, and membrane proteins. Lipid molecules in the membrane have several time scales of motions ranging from femtosecond to seconds. The flip-flop motion, in which lipid molecules move from one leaflet to the other, is known to be one of the slowest: it typically occurs within several tens of seconds, or much longer. Recently, experimental studies revealed that cholesterol (CHOL), diacylglycerols (DAG), and ceramides (CER) show fast flip-flop motions in some membranes. However, the molecular mechanisms underlying the motions remain elusive.

In this work, we performed coarse-grained molecular dynamics simulations, using MALTINI force field parameters. We examined flip-flop motions of CHOL, DAG, and CER in phospholipid bilayer systems, composing of DAPC(di-20:4), SAPC(18:0-20:4), and POPC(16:0-18:1). In the simulations using DAPC membranes, we observed flip-flop motions of CHOL, DAG, and CER within a microsecond. The flip-flop rate of CHOL was the highest, whereas that of DAG was lower than CHOL. CER flipped only once during the simulation. This tendency of flip-flop motions is strongly correlated with the relative positions of the lipids to the bilayer membranes: CHOL stays almost at the center of the membrane, whereas the head group of CER is located at the water/membrane interface and interacted with solvent molecules strongly.

The flip-flop motions of lipids were also affected with the membrane environment. Within 1-microsecond simulations, CHOL flipped 257 times in DAPC, 196 times in SAPC, and 5 times in POPC. Thus, the flip-flop rate is strongly correlated with the number of double bonds in the acyl chains of bilayer phospholipids, suggesting the importance of the membrane fluidity. These simulation results qualitatively agree with existing experimental data and shed light on the molecular mechanisms underlying the dynamics of biomembranes.

2538-Pos

The Effect of Cholesterol on Membrane Chain-Chain Packing

Ming-Tao Lee¹, Wei-Chin Hung².

¹National Synchrotron Radiation Research Center, Hsinchu, Taiwan,

²R.O.C. Military Academy, Fengshan, Taiwan.

Membrane structure is very important issue for cell. Membrane is not only the "wall" for protecting cell but also the interface for exchanging signal, ions and molecules. Many evidences show that membrane protein will fold to functional structure by associating with suitable membrane structure. Lipid chain-chain packing is one of important structures and will affect membrane thickness, lipid lateral diffusion and membrane domain formation. We will use grazing incident X-ray diffraction to probe lipid chain-chain packing. The 12keV X-ray light source in BL13A beam line of NSRRC and home-made humidity-temperature controlled chamber will be applied in the measurements. Cholesterol will