

expose their anionic lipid content during chilling, we propose that PLA2-IIA is triggered during this process, and that the activity of the enzyme on the fluid domains induces an overall increase in rigidity in the platelet plasma membrane due to an increase in the proportion of ordered lipids. We present a combined FTIR and Laurdan fluorescence spectroscopy study on the thermotropic phase behavior of human platelets, and show that during cold-induced activation membrane rigidity increases. We present results on how the activity of sPLA2-IIA is regulated by the thermotropic phase behavior of the platelet plasma membrane after platelets have been artificially activated through sonication. We show that PLA2-IIA activity increases the rigidity of these activated platelet plasma membranes.

2556-Pos

Neutron Scattering and MD Simulation Study of DOPC and DOPC/cholesterol Bilayers

Eduardo Jardon-Valedez¹, J. Alfredo Freites¹, Ella Mihailescu², David Worcester², Stephen H. White¹, Douglas J. Tobias¹.

¹UCI-Irvine, Irvine, CA, USA, ²NIST Center for Neutron Research, Gaithersburg, MD, USA.

Cholesterol represents about 35% of the lipid content in the plasma membrane. Relevant cellular processes that have been associated with cholesterol-enriched microdomains include, among others, signaling, trafficking, cytoskeleton organization, apoptosis, and cell adhesion. In this work we studied the effects of cholesterol on model bilayers of DOPC using a combination of molecular dynamics simulations and neutron scattering experiments. Taking advantage of the higher scattering length of deuterium relative to hydrogen, the location of labeled components can be determined by difference-structure analysis of the scattering length density profiles. By labeling the terminal methyl groups of DOPC and a cholesterol ring, we determined their respective mass distributions across the membrane in DOPC and DOPC/cholesterol bilayers. The neutron data reveal the terminal methyl groups in a pure DOPC bilayer visit the head group-water interface as observed by the prominent wings in the mass distribution. A narrower terminal methyl group distribution of DOPC in the presence of cholesterol suggests an enhancement of the order of the lipid acyl chains. Cholesterol heads, on the other hand, point toward the aqueous interface, as expected. Order parameters S_{CD} calculated from the MD simulations confirm that DOPC is more ordered in presence of cholesterol. The structural analysis based on MD simulation in concert with neutron diffraction experiments provides valuable data for advancing understanding of role of cholesterol in altering transbilayer communication in cholesterol-enriched microdomains found in biological membranes.

Membrane Receptors & Signal Transduction II

2557-Pos

Distribution and Dynamics of RBL IgE Receptors (FcεRI) Quantitatively Observed on Planar Ligand-Presenting Surfaces

Kathrin Spendier, Amanda Carroll-Portillo, Keith Lidke, Bridget S. Wilson, Jerilyn A. Timlin, V.M. Kenkre, James L. Thomas.

University of New Mexico, Albuquerque, NM, USA.

There is considerable interest in the signaling mechanisms of immunoreceptors, especially when triggered with membrane-bound ligands. When T cells, B cells, or mast cells bind to monovalent ligands on fluid supported lipid bilayers, receptor clustering, signaling, and receptor redistribution into immunological synapses follow. We have quantitatively studied the kinetics of redistribution of IgE receptors (FcεRI) on RBL-2H3 mast cell surfaces. To separate the kinetics of receptor redistribution from cell spreading, the initial cell-substrate contact time was precisely defined (± 50 ms) by micropipette cell manipulation. Using total internal reflection fluorescence, the distribution and dynamics of receptor clusters were imaged. We find strong quantitative evidence that initial receptor aggregation occurs at cellular protrusions, which are visible bright spots/regions on cells contacting ligand-free surfaces. The initial size of these regions is independent of the substrate and the presence or absence of ligand, and they were found to be randomly distributed over the interfacial contact area. Using a finite-element diffusion model, we found that the initial rate of accumulation of receptors at the protrusions is consistent with diffusion-limited trapping with $D \sim 10^{-1} \mu\text{m}^2/\text{s}$. At longer timescales, individual clusters on ligand-bearing membranes were observed to move with both a diffusive and a directed component of motion; clusters eventually coalesced near the center of the contact region (~ 1 minute). The dynamics of the early cluster motion is similar to the dynamics of membrane fluctuations of cells on ligand-free fluid membranes. Thus, the same cellular machinery may be responsible for both processes.

Sandia is a multi-program laboratory operated by Sandia Corporation, a Lockheed Martin Company, for the United States Department of Energy under Contract DE-ACO4-94AL85000.

2558-Pos

Multi-Color Quantum Dot Tracking to Correlate FcεRI Aggregate Size with Mobility

Patrick J. Cutler, Bridget S. Wilson, Janet M. Oliver, Keith A. Lidke, Diane S. Lidke.

University of New Mexico, Albuquerque, NM, USA.

The high affinity IgE receptor, FcεRI, is the primary immune receptor expressed on mast cells and basophils. These receptors bind circulating IgE with high affinity and are activated when multivalent allergen crosslinks IgE-bound receptors, initiating a complex signaling cascade that leads to the release of inflammatory mediators by degranulation. It is also well documented that FcεRI crosslinking leads to receptor immobilization. However, the exact mechanism of crosslink-induced activation is not completely understood.

We have used single quantum dot (QD) tracking to characterize the mobility of QD-IgE-FcεRI in the resting and activated state. We have previously shown that crosslink-induced immobilization is dependent on the actin cytoskeleton as well as the antigen dose. Furthermore, we have determined that small, mobile clusters are signaling competent and that immobilization is the trigger for internalization rather than signal initiation. In these experiments, a combination of QD-IgE and dark IgE were crosslinked with multivalent antigen; therefore, the cluster size could not be directly correlated with mobility. We have now developed an experimental strategy and algorithmic approach to differentiate between monomers, dimers, trimers, and tetramers. Using a four-color beam splitter, we can simultaneously track FcεRI complexes labeled with four spectrally distinct QDs (525, 585, 655, and 705) at rates up to 30 frames/s. From these data, we use QD localization and channel overlay accuracies to identify monomers, dimers, trimers, and tetramers and determine the diffusion coefficient for each type of aggregate. We demonstrate this approach with both simulated and experimental data.

2559-Pos

Fluorogen Activating Peptides for Single Molecule Localization-Based Superresolution and Single Particle Tracking of FcεRI Subunits

Samantha L. Schwartz.

University of New Mexico, Albuquerque NM, NM, USA.

Fluorogen activating peptides (FAPs) are genetically expressible tags that increase the fluorescence excitation cross-section of dye binding partners by up to four orders of magnitude. The binding of FAPs with corresponding fluorogens is characterized by nanomolar affinity, corresponding to bound lifetimes of up to ten seconds. FAPs exhibit resistance to photobleaching, with single peptides repeatedly binding and activating new dye molecules. The resulting long observation lifetimes for single peptides make the FAP system a convenient, expressible probe for single particle tracking on live cells. The intermittency due to the equilibrium of dye binding and unbinding can be used for localization-based superresolution.

We apply this technology to study the localization and dynamics of the high affinity IgE receptor, FcεRI, the primary multi-subunit receptor on mast cells and basophils. Circulating, allergen-specific IgE molecules bind to FcεRI receptors via the α -subunit. Receptor crosslinking by cognate multivalent allergen initiates the immune response by activation of the immunoreceptor tyrosine-based activation motifs on the β - and γ -subunits. We have previously characterized the dynamics of the FcεRI by tracking of quantum dot (QD)-labeled IgE bound to FcεRI α . To compare the behavior of the α - and γ -subunits, we have generated an FAP-tagged FcεRI γ -subunit that is expressed on the cell surface of rat basophilic leukemia cells and binds an exogenous fluorogen based on the malachite green dye. We use this probe to determine dynamics of the γ -subunit during resting and activated states and to generate superresolution images of FcεRI γ distribution. We use QD-IgE, in conjunction with the FAP-FcεRI γ , for two-color single particle tracking to observe the relative spatial and temporal dynamics of the α - and γ -subunits.

2560-Pos

Receptor Cluster Size Affects Signaling in Breast Epithelial Cancer Cells

Qian Xu, Jay T. Groves.

UC Berkeley, Berkeley, CA, USA.

Our purpose is to study a potentially novel mechanical regulatory mechanism in the EphA2 signaling pathway. EphA2 is a receptor tyrosine kinase that is known to be up-regulated in 40% of human breast cancers and plays an active role in metastasis. Activation of EphA2 occurs after binding to its ligand, presented on an opposing cell membrane. Monomer binding is followed by