

Our results indicate that LPA-membrane interactions depend on initial surface pressure, phospholipid headgroup and degree of acyl chain saturation, presence and amount of cholesterol, aqueous media conditions, and aggregation state of LPA. These data suggest that, in addition to its function as a ligand for specific GPCR, LPA interacts directly with the target membrane, constituting a role for this phospholipid as a physical regulatory molecule for LPA cellular signaling pathways.

3590-Pos

Molecular Hydration Investigated using Extended Membrane Surfaces

Adriana L. Rogozea, Matthew J. Justice, Horia I. Petrache.

Indiana University Purdue University Indianapolis, Indianapolis, IN, USA.

In biological cells and in solutions, biomolecules are in constant competition for water. The availability of water is critical to a wide array of processes such as protein folding, molecular recognition, and cell signaling. To understand these mechanisms, we distinguish between the water molecules residing in the vicinity of biological macromolecules and the most distant water molecules in the bath. What happens on the surface of macromolecules? How to investigate the role and the properties of interfacial water? The problem is that the spatial extent of hydration layers is very small, on the order of nanometers or less. How to produce an experimentally measurable signal? Our approach to investigate the many remaining mysteries of molecular hydration is by using the naturally extended water surfaces within multilamellar lipid structures. We show how addition of salts, sugars, and most buffers make the interlamellar (D) spacing of synthetic phospholipids membranes to increase. In contrast to this behavior, low concentrations of highly hygroscopic molecules such as PEG (polyethylene glycol) and DMSO (dimethyl sulfide) are found to decrease the D-spacing. A very interesting case is that of the small molecular weight PEG 400 with unusual effects on the interfacial hydration, in such a way that at some concentration threshold the PEG molecules overcome a mixing barrier and become included in the forbidding interlamellar water space. This behavior is due to competitions between the strong exclusions forces from extended hydration surfaces and entropy. We are currently investigating how biological relevant molecules such as "Factor V" (BSA Bovine Serum Albumin) are modifying cellular osmoregulation and its effects on membranes stress. Knowing how molecules interact at the lipid-water interface could prove beneficial in the drug design of anesthetics, cryoprotectants of mammalian cells, and in general, of molecular stressor affecting biological cells.

Signaling & Membrane Transformations

3591-Pos

Determination of Threshold Forces for Tether Formation in Vesicles

Daniel Stark, Thomas Killian, Robert Raphael.

Rice University, Houston, TX, USA.

Membrane tether experiments, which involve applying point forces to membranes to form thin tubes of the membrane, provide a powerful method to drastically alter membrane curvature. To study the formation of membrane tethers, we have designed and built a magnetic force transducer (MFT) using microfabrication techniques. MFTs have traditionally suffered from an inability to precisely calibrate the force. Quantitative determination of the tether formation force using theoretical models of membrane mechanics, may be done very accurately if the applied force is known with sub-piconewton accuracy. Our initial results indicate our ability to control the amplitude of the force up to 10 pN within ± 0.2 pN over a constant length of 50 microns. We have used this device to determine tether formation forces from POPC giant unilamellar vesicles and find that formation forces range from 3 to 10 pN. When tethers are repeatedly pulled from the same vesicle, the formation force is constant, suggesting that individual GUVs may have differences in their mechanical properties. A particular advantage of our device is that a wide range of dynamic force profiles can be applied via a computer-controlled interface, enabling studies of responses of membranes to dynamic force application at kilohertz frequencies.

3592-Pos

Quantifying Pathogen Recognition of the Cell Membrane: Simple Sugars Simulate the Functions of Complex Glycans

Xiaoyang Zhu¹, Athena Guo².

¹University of Texas, Austin, TX, USA, ²MicroSurfaces, Inc., Austin, TX, USA.

The first step leading to the breaching of the cell membrane and infection of healthy cells is often the multivalent recognition and adhesion of glycan binding proteins (GBPs) on pathogens and glycans on host cell membranes. Such

multivalent interaction depends critically on the mobility and density of signaling molecules on the membrane surface. While glycan microarrays have been used in exploring multivalent interactions, the lack of mobility and the difficulty in controlling surface density both limit their quantitative applications. Here we apply a fluidic glycan microarray, with glycan density varying for orders of magnitude, to profile cell surface interaction using a model system, the adhesion of *Escherichia coli* (*E. coli*) to mannose. We show the quantitative determination of monovalent and multivalent adhesion channels; the latter can be inhibited by nanoparticles presenting a high density of mannosyl groups. These results reveal a new *E. coli* adhesion mechanism: the switching in the FimH adhesion protein avidity from monovalent to multivalent as the density of mobile mannosyl groups increases; such avidity switching enhances binding affinity and triggers multiple fimbriae anchoring. Affinity enhancement towards FimH has only been observed before for oligo-mannose due to the turn on of secondary interactions outside the mannose binding pocket. We suggest that the new mechanism revealed by the fluidic microarray is of general significance to cell surface interactions: the dynamic clustering of simple sugar groups (homogeneous or heterogeneous) on the fluidic membrane surface may simulate the functions of complex glycan molecules.

3593-Pos

Dynamical Basis of the Enhancement of the Enzymatic Activity of Factor VIIa by Tissue Factor

Y Zenmei Ohkubo, Emad Tajkhorshid.

Department of Biochemistry, Center for Biophysics and Computational Biology, and Beckman Institute, University of Illinois at Urbana-Champaign, Urbana, IL, USA.

Activation of coagulation factors is regulated by their binding to and complex formation on the surface of anionic membrane resulting in an increase of the enzymatic activity by several orders of magnitude. Complex formation of tissue factor (TF) and factor VIIa (FVIIa) on anionic membranes constitutes one of the key steps in the coagulation cascade. The mechanisms for the enhancement of the enzymatic activity of FVIIa by TF is not fully understood, primarily due to the lack of atomic models for the membrane-bound form of the TF:FVIIa complex.

We report a first membrane-bound model of the TF:FVIIa complex resulting from large-scale molecular dynamics simulations. The dynamics of FVIIa and soluble TF (sTF) was investigated in solution and on the membrane, both in their isolated and complex forms. Our model of membrane-bound GLA domain was used to construct the membrane-bound models of the TF:FVIIa complex and monomers. All-atom simulations were performed for tens of nanoseconds to investigate the protein dynamics after establishing optimal protein-protein/protein-lipid interactions.

The results reveal that sTF restricts the motion of FVIIa, thus optimally orienting its catalytic triad for the interaction with its substrate factor X (FX). Several direct interactions between the membrane lipids and the side chains of sTF, including the regions of K159-K166 and D180-N184 that form the exosite of the substrate FX, are observed. Interestingly, sTF sustains two distinct orientations against the membrane and different conformations of the K159-K166 loop, in the isolated and complex forms. These findings suggest that TF orients itself on the membrane surface through the interactions both to the membrane and FVIIa, independent of its trans-membrane anchoring helix and that the exosite on TF is available for FX binding only after TF:FVIIa complex is formed.

3594-Pos

Regulation of Phospholipase C Beta - Rac1 Cytoskeletal Pathways by Gamma Synuclein

Urszula P. Golebiewska^{1,2}, Jason-Flor Sisante¹, Jaak Raudsepp^{1,3}, Gyongyi Mihalyne¹, Suzanne Scarlata¹.

¹Stony Brook University, Stony Brook, NY, USA, ²Queensborough Community College, Bayside, NY, USA, ³Northport High School, Northport, NY, USA.

The breast cancer specific gene protein 1 also known as γ -synuclein is undetectable in normal or benign breast lesions, but it is highly expressed in infiltrating breast cancer. The precise role of γ -synuclein in malignancies is not well known. We investigated whether γ -synuclein might affect activity of phospholipase C β 2 (PLC β 2). PLC β 2 is also absent in normal breast tissue, but it is highly expressed in breast tumors where it is correlated with the progression and migration of the tumor. Expression of PLC β 2 is highly correlated with expression of γ -synuclein. We found that γ -synuclein binds PLC β 2 in vitro with high affinity, $K_d = 23 \pm 3$ nM. PLC β 2 is activated by heterotrimeric G protein and by members of Rho family of GTPases (in particular Rac1), which are