

British Journal of Pharmacology

**Instructions to Authors
Nomenclature Guidelines**

**Index
Volumes 111–113
1994**

BRITISH JOURNAL OF PHARMACOLOGY

The *British Journal of Pharmacology* welcomes contributions in all fields of experimental pharmacology including neuroscience, biochemical, cellular and molecular pharmacology. The Board of Editors represents a wide range of expertise and ensures that well-presented work is published as promptly as possible, consistent with maintaining the overall quality of the journal.

Edited for the British Pharmacological Society by

A.T. Birmingham
(Chairman)

R.W. Horton **W.A. Large**
(Secretaries)

Editorial Board

J.A. Angus *Melbourne, Australia*
M.L.J. Ashford *Cambridge*
G.W. Bennett *Nottingham*
W.C. Bowman *Glasgow*
N.G. Bowery *London*
Alison F. Brading *Oxford*
S.D. Brain *London*
K.D. Butler *Horsham*
M. Caulfield *London*
R. Chess-Williams *Sheffield*
M.K. Church *Southampton*
T. Cocks *Melbourne, Australia*
S.J. Coker *Liverpool*
R.A. Coleman *Ware*
Helen M. Cox *London*
A.J. Cross *London*
V. Crunelli *Cardiff*
T.C. Cunnane *Oxford*
F. Cunningham *London*
A. Dray *London*
J.R. Docherty *Dublin*
J.M. Edwardson *Cambridge*
P.C. Emson *Cambridge*
W. Feniuk *Cambridge*
J.R. Fozard *Basle, Switzerland*
Allison D. Fryer *Baltimore, USA*

Sheila M. Gardiner *Nottingham*
C.J. Garland *Bristol*
L.G. Garland *Beckenham*
A. Gibson *London*
R. W. Gristwood *Cambridge*
D.W.P. Hay *Philadelphia, USA*
P.G. Hellewell *London*
P.E. Hicks *Edinburgh*
S.J. Hill *Nottingham*
S.M.O. Hourani *Guildford*
J.C. Hunter *Cambridge*
C.C. Jordan *Ware*
D.A. Kendall *Nottingham*
P. Leff *Loughborough*
R. McMillan *Macclesfield*
J. Maclagan *London*
C.A. Maggi *Florence, Italy*
Janice M. Marshall *Birmingham*
G. Martin *Beckenham*
W. Martin *Glasgow*
A. Mathie *London*
D.N. Middlemiss *Harlow*
P.K. Moore *London*
R.J. Naylor *Bradford*
C.D. Nicholson *Oss, The Netherlands*

C.P. Page *London*
A.N. Payne *Beckenham*
F.L. Pearce *London*
J.D. Pearson *London*
A.G. Renwick *Southampton*
M.H.T. Roberts *Cardiff*
C. Robinson *London*
G.J. Sanger *Harlow*
M.A. Simmonds *London*
J.M. Sneddon *Sunderland*
P. Sneddon *Glasgow*
K. Starke *Freiburg, Germany*
R.J. Summers *Melbourne, Australia*
P.V. Taberner *Bristol*
J. Tamargo *Madrid, Spain*
C. Thiemerann *London*
M.D. Tricklebank *Harlow*
M.B. Tyers *Ware*
S.P. Watson *Oxford*
K.J. Watling *Boston, USA*
A.H. Weston *Manchester*
B.J.R. Whittle *Beckenham*
Eileen Winslow *Riom, France*
B. Woodward *Bath*
E.H.F. Wong *California, USA*

Corresponding Editors

P.R. Adams *Stony Brook, U.S.A.*
C. Bell *Melbourne, Australia*
F.E. Bloom *La Jolla, U.S.A.*
A.L.A. Boura *Newcastle, Australia*
N.J. Dun *Toledo, U.S.A.*
R.F. Furchtgott *New York, U.S.A.*
T. Godfraind *Brussels, Belgium*
S.Z. Langer *Paris, France*

R.J. Miller *Chicago, U.S.A.*
R.C. Murphy *Denver, U.S.A.*
E. Muscholl *Mainz, Germany*
R.A. North *Portland, U.S.A.*
M. Otsuka *Tokyo, Japan*
M.J. Rand *Melbourne, Australia*
S. Rosell *Södertälje, Sweden*
P. Seeman *Toronto, Canada*

L. Szekeres *Szeged, Hungary*
B. Uvnas *Stockholm, Sweden*
P.A. Van Zwieten *Amsterdam, Netherlands*
V.M. Varagić *Belgrade, Yugoslavia*
G. Velo *Verona, Italy*
Wang Zhen Gang *Beijing, China*
M.B.H. Youdim *Haifa, Israel*

Submission of manuscripts: Manuscripts (two copies) should be sent to The Editorial Office, British Journal of Pharmacology, St. George's Hospital Medical School, Cranmer Terrace, London SW17 0RE.

Authors should consult the Instructions to Authors and the Nomenclature Guidelines for Authors in Vol. 114, 245–255. These Instructions and Guidelines also appear with the journal Index for Volumes 111–113, 1994. A checklist of the essential requirements is summarised in each issue of the journal, or as the last page of the issue.

Whilst every effort is made by the publishers and editorial committee to see that no inaccurate or misleading data, opinion or statement appears in this Journal, they and the *British Pharmacological Society* wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Accordingly, the *British Pharmacological Society*, the publishers and the editorial committee and their respective employees, officers and agents accept no liability whatsoever for the consequences of any such inaccurate or misleading data, opinion or statement.

INSTRUCTIONS TO AUTHORS

With effect from 1 January 1995

The *British Journal of Pharmacology* welcomes contributions in all fields of experimental pharmacology, including neuroscience, biochemical, cellular and molecular pharmacology, for publication as full papers or as high priority Special Reports.

Papers should normally be based on new results obtained experimentally and should constitute a significant contribution to pharmacological knowledge. Papers which reassess pharmacological concepts based on earlier results will also be considered as will purely theoretical papers. Papers dealing only with descriptions of methods are acceptable if new principles are involved.

Contributions that have already been published, or accepted or are under consideration for publication, with essentially the same content will not be considered. This restriction does not apply to results published as abstracts of communications, letters to editors, or as contributions to symposia, provided that the submission adds significantly to the information available in the previously published contribution.

Papers are only accepted if accompanied by a Declaration which must be signed by all Authors. This Declaration concerns the originality of the submitted paper and assigns the copyright of all papers accepted for publication to Stockton Press Ltd. on behalf of the British Pharmacological Society. See pages 6 and 7 for details.

The Journal will not consider papers which describe experiments on animals which do not fall clearly within the current laws governing animal experimentation in the United Kingdom. Authors must make it clear that the procedures they use were as humane as possible and the doses (initial and subsequent) of anaesthetics and analgesics should be clearly stated; the method of assessing anaesthesia, particularly after the administration of skeletal muscle relaxants (neuromuscular blocking drugs), must be well defined. The Society has an Ethics Committee which can be consulted by authors through the Secretaries to the Editorial Board.

When investigations on normal human subjects are reported, evidence of approval by a local Ethics Committee must be given. Papers concerned with clinical trials or investigations of the effects of drugs on patients are not appropriate for this Journal.

Authors are strongly urged to keep their manuscripts as short as they reasonably can. An effective way is to reduce the Discussion and the number of figures to a minimum and to avoid repetition of information that has already been published. Authors should remember that a reader may be influenced by literary style and will appreciate simple but accurate prose.

It is important to note that failure to comply with 'Instructions to Authors' may lead to considerable editorial delays.

FULL PAPERS

Manuscripts must be typed on one side of A4 paper. Words at the end of lines should not be divided because they may become incorrectly hyphenated. Handwritten characters or symbols (e.g. Greek letters) should be spelled out in full in the margin. Papers in recent issues of the *British Journal of Pharmacology* should be consulted for the general layout of the paper and also for details. The following subsections are used:

1. Title page
2. Summary
3. Introduction
4. Methods

5. Results
6. Discussion and conclusions
7. Acknowledgements
8. List of references
9. Tables

10. Figures and captions

The type must not be smaller than 12 pitch or 10 point. Each section must be typed in **double spacing** with margins of not less than 2.5 cm all round and each page should be numbered. The original and one copy of the typescript should be supplied.

Title page

The title should normally contain **no more than 150 characters** and should not consist of a sentence (statement or conclusion) or be interrogative. A **short running title** containing not more than 50 characters and spaces is also required. The title page should include the names of authors and their appropriate addresses. It should be made clear which address relates to which author. Authors' present addresses differing from those at which the work was carried out should be given as footnotes on the title page and references at the appropriate place in the author list by superscript numbers. A footnote may also be used to indicate the author to whom correspondence should be sent. The use of footnotes for any other reason is not allowed. If the address to which proofs should be sent is not that of the first mentioned author, clear instructions should be given in a covering note and not on the title page. The title page should be paginated as page 1 of the paper.

Summary

The summary will be printed at the beginning of the paper. It should not exceed 5% of the length of the paper and should contain a brief account of the problem, the methods, results and the conclusions. It should be arranged in **numbered and concise paragraphs**. Up to ten **keywords** or phrases of two to three words (including names and terms used in the title) should be displayed at the end of the summary. Keywords will be used to compile the annual index. The quality of the index will thus be determined by the appropriateness of the keywords. These may be selected by reference to the most recent Index of the Journal. Avoid unhelpful or unqualified terms such as 'rat', 'drug' etc.

Introduction

The introduction should give a short and clear account of the background of the problem and the rationale of the investigation. Only previous work that has a direct bearing on the present problem should be cited.

Methods

The methods must be described in sufficient detail to allow the experiment to be interpreted and repeated by the reader. However, detailed repetition of methods which have been adequately described previously should be avoided and references given, although a brief outline is often helpful.

Drugs should be listed in a separate paragraph. Their names should be 'approved names' as published previously in British Approved Names, 1990 (HMSO). If a drug has no 'approved name' its chemical name must be used and the rules set out in the current *Handbook for Chemical Society*

Authors (London, Chemical Society) observed, or its structural formula given. Cumbersome chemical names should be suitably abbreviated for later reference in the paper.

The doses of drugs should be given as unit weight per body weight, e.g. mmol kg⁻¹ or mg kg⁻¹; concentrations should be given in terms of molarity, e.g. nM or μ M.

Reference should be made to any statistical analyses that have been performed on the results in order, for example, to determine the significance of differences between results obtained under different conditions.

Results

The description of the experimental results should be succinct but, nevertheless, in sufficient detail to allow the experiments to be repeated by others. Typical single experiments may be presented with a clear statement that n number of similar experiments had similar results. Where appropriate, however, the mean results with confidence limits or with standard errors of the means and the number of observations should be given. Statistical tests of significance should be performed where appropriate. The results of such tests should be stated as the numerical value of the probability (P) that is calculated, with any necessary clarification (e.g. one-tail or two-tail test).

Every effort should be made to avoid unnecessary repetition of data in the text, tables and figures. Conclusions and theoretical considerations should not be elaborated in this section.

Discussion

The purpose of the discussion is to present a brief and pertinent interpretation of the results against the background of existing knowledge. Any assumptions on which conclusions are based must be stated clearly. A mere recapitulation of the results is not acceptable. A review-like treatment, which reduces the impact on the reader, should also be avoided. The main conclusion should be conveyed in a final paragraph.

Acknowledgements

Acknowledgements should be brief but should include reference to sources of support. Sources of drugs not widely available commercially should be acknowledged.

References

In the text, references to other work should take the form: (Bolton & Kitamura, 1983) or, 'Bolton & Kitamura (1983) showed that . . .'. If there are more than two authors, the first author's name should be given followed by *et al.* (Bülbring *et al.*, 1981).

References to 'unpublished observations' or 'personal communications' should be mentioned in the text only, and not included in the list of references. Papers which have been submitted and accepted for publication, should be included in the list of references with the names of the periodicals and 'in press'. A photocopy should normally be submitted with the manuscript. If this is not possible, authors should indicate whether the work cited is an abstract or a full paper. Papers in preparation or which have been submitted but not yet finally accepted for publication must not be included in the list of references.

The reference list at the end of the manuscript must be arranged alphabetically according to the surname of the first author. When the surnames of authors are identical, the alphabetical order of their initials takes precedence over the year of publication. The AUTHORS' names are followed by the year of publication in brackets. If more than one paper by the same authors in one year are cited, a, b, c, etc. are placed after the year of publication, both in the text and in

the list of references. The title of the article is given in full, followed by the *abbreviated title of the periodical*, volume number and first and last page numbers. The abbreviations used for periodicals are those of the most recent edition of the International List of Periodical Title Word Abbreviations. References to articles in books should consist of names of authors, year of publication, title of article followed by the *title of the book*, the editors, volume numbers, if any, and page numbers, the place of publication and the names of the publishers. For example:

BOLTON, T.B. & KITAMURA, K. (1983). Evidence that ionic channels associated with the muscarinic receptor of smooth muscle may admit calcium. *Br. J. Pharmacol.*, **78**, 405–416.
BRADING, A.F. (1981). Ionic distribution and mechanisms of transmembrane ion movements in smooth muscle. In *Smooth Muscle: An Assessment of Current Knowledge*, ed. Bülbring, E., Brading, A.F., Jones, A.W. & Tomita, T. pp. 65–92. London: Edward Arnold.

Tables

Each table should be given on a separate page, paginated as part of the paper. Tables should be numbered consecutively with arabic numerals and the number should be followed by a brief descriptive caption, occupying not more than two lines, at the head of the table. The proportions of the text area should be borne in mind when designing the layout of tables. For the sake of clarity, tables should not have more than 120 characters to a line, with spaces between columns counted as four characters. The absolute maximum is 180 characters to a line. Each column should have a heading and the units of measurement should be given in parentheses in the heading. Except in special circumstances, tables should be self-explanatory; the necessary descriptions should be at the bottom of the table.

Figures

To avoid unnecessary Figures, particularly those requiring half-tone reproduction, only critical points of the text should be illustrated. If coloured Figures are desired, the Authors should discuss their requirements with the Secretaries, preferably before submission.

Please note that unsatisfactory Figures will be returned to the Author for revision. The Journal reserves the right to reject a manuscript if the Figures are unacceptable.

Submission Requirements

- (a) The Authors' names and the Figure number must be indicated lightly *in pencil* on the back of each Figure; if necessary, use an adhesive label to avoid damage to the Figure.
- (b) Each copy of the manuscript must be accompanied by one set of labelled Figures (i.e. complete with lettering and numbering, arrows, etc.). An original set and one high quality photocopy will suffice.
- (c) Another original set of Figures identical in size but without any letters or numbers must also be supplied for the use of the Publisher. Arrows and event marks on experimental records may be retained, provided they are larger than 3 mm wide. The Publisher will choose the correct style of typeface of an appropriate size to suit the final size of the Figure on the printed page.
- (d) No submitted Figure should exceed 210 × 297 mm (A4).
- (e) Each Figure must be accompanied by a legend; each legend should be typed on a *separate* sheet of paper and paginated as part of the manuscript. Legends should explain the Figures in sufficient detail that, whenever possible, they can be understood without reference to the text.

Line width (axes)	Line width (graphs)	Symbol size	Figure will reduce to this percentage of the original size
—	—	Δ□○	100 (No reduction)
—	—	Δ□○	80
—	—	Δ□○	70
—	—	Δ□○	60
—	—	Δ□○	50
—	—	Δ□○	40

Line Figures

It is best to submit an original drawing (black ink on heavy white paper or faint blue graph paper) which has been prepared to conform with the style and convention of the Journal, because redrawing is expensive. The original drawing should be lettered in pencil and should be larger (up to two times as large) than the intended size in the Journal.

It is important that the printed symbols and lines should retain their clarity. To achieve this the symbols and lines in original drawings should be sharply defined and of an even density and breadth. When graphs are generated by computer, lines must not show noticeable stepping. Heavier (broader) lines should be used for curves than for the axes of graphs. The table above illustrates line widths and symbol sizes to be used on a figure and the appropriate reductions in the final printed form.

Symbols should be chosen from the following set

○ ● □ ■ △ ▲ ▽ ▾ ◊ ◆ + ×

The preferred order to shading of histogram columns is: open (clear), closed (solid), cross-hatched, heavily stippled and other (if required).

The explanation of the symbols and column headings should be given in the Figure legend and not as a key in the Figure itself.

Line Figures should normally have only left and bottom axes; box-style Figures and those using 3-dimensions are not acceptable.

Where the Figure is a composite of more than one graph, experimental record, etc., particular care is needed to minimise the spaces between each part, without overcrowding the entire Figure.

Figure 1 illustrates a simple properly-drawn graph in its original form (a) and in its reduced form (b) as it would appear in the Journal.

Photographs and photomicrographs

These should be submitted, twice as large as their intended published size, as good quality prints of high contrast especially where traces and records are illustrated. The originals must not contain arrows, lettering or numbering; these must be accurately located on a duplicate print (or photocopy). When submitting half-tone illustrations for publication authors should remember that it is not possible to reproduce Figures to a finer quality than the original photographs/photomicrographs provided. Critical areas

should be marked on a second copy or on an overlay, so that the Printer can choose the correct exposure. Maximum trim areas should be marked on a second copy of the photograph/photomicrograph or on a tracing overlay, i.e. authors should show any parts of the photographs that could be excluded from the finished half-tone illustration. A calibration bar must be provided on the photomicrograph to ensure that, if the Printer reduces the plate, the scale is reduced in the correct proportion.

Proofs

Two sets of page proofs will be supplied, one of which should be retained by the authors. The other should be corrected immediately and returned to the Publisher. Corrections should be kept to a minimum.

SPECIAL REPORTS

The purpose of *Special Reports*, which have superseded 'Short Communications', is to provide rapid publication for new and important results which the Editorial Board considers are likely to be of special pharmacological significance. *Special Reports* will have publication priority over all other material and so authors are asked to consider carefully the status of their work before submission.

In order to speed publication there is normally no revision allowed beyond very minor typographical or grammatical corrections. If significant revision is required, the Board may either invite rapid re-submission or, more probably, propose that it be re-written as a Full Paper and be re-submitted for consideration. In order to reduce delays, proofs of *Special Reports* will be sent to authors but **essential corrections must reach the Publisher within 48 hours of receipt**. Authors should ensure that their submitted material conforms exactly to the following requirements.

Special Reports should normally occupy no more than two printed pages of the Journal; two illustrations (Figures or Tables, with legends) are permitted. As a guideline, with type face of 12 pitch and double-line spacing, a page of A4 paper could contain about 400 words. The absolute maximum length of the *Special Report* is 1700 words. For each Figure or Table, please deduct 200 words. The manuscript should comprise a Title page, a Summary consisting of a single short paragraph, followed by keywords (maximum of 10), Introduction, Methods, Results, Discussion and References (maximum of 10). In all other respects, the requirements are the same as for Full Papers.




Figure 1 (a) Artwork as drawn. (b) Artwork reduced to 60 per cent of its original size for publication in the Journal.

STATEMENT AND COPYRIGHT AGREEMENT

The following Statement, Declaration and Copyright Agreement should be read carefully by Authors who should then send a copy of their Declaration with their manuscript following the example given in this section.

Statement

1. Submission of a manuscript will be taken to imply that the Authors have obtained permission to publish from (a) their employers or institution, if they have a contractual or moral obligation to do so, and (b) those whose unpub-

lished work, including papers accepted for publication (i.e. in press), has been cited or those whom the Authors wish to acknowledge as having improved the content or presentation of the manuscript.

2. The Authors must declare that the manuscript contents are original and that they have not already been published or accepted for publication, either in whole or in part, in any form other than as an abstract or other preliminary publication in an unrefereed article. Furthermore, the Authors must verify that no part of the manuscript is under consideration for publication elsewhere and it will not be submitted elsewhere if accepted by the *British Journal of Pharmacology* and not before a decision has been reached by the Editorial Board.

Declaration

I/We assign to Stockton Press, on behalf of the British Pharmacological Society, the copyright of my/our manuscript, currently entitled

‘.....’,
for publication in the *British Journal of Pharmacology*

Furthermore I/We have read, understood and accepted the terms and conditions as set out in Statement and Copyright, Instructions to Authors *Br. J. Pharmacol.* 1995, 114, 245–251.

Name

Signed

Name

Signed

Name

Signed

Date

Copyright

1. The Authors must agree that, when the above manuscript has been accepted for publication in this Journal, the worldwide copyright shall pass to the Stockton Press Ltd. on behalf of the British Pharmacological Society, on the understanding that the assignment of copyright will not affect subsisting Patent Rights arrangements pertaining to it. The Authors also accept that, when accepted, the contents will not be published subsequently in the same or similar form in any language without the consent of the Publisher or Editorial Board of the Journal.

This Agreement shall not compromise the Authors' rights to reproduce their own work (see 3 below). For its part, the British Pharmacological Society will protect the interest of Authors in the matter of copyright.

2. The Authors must declare that, where excerpts from copyrighted works have been included, the Authors have obtained written permission from the Copyright owners and have credited the sources in the manuscript. They must also warrant that the article contains no libellous or unlawful statements and does not infringe the rights of others.
3. The Authors will be entitled to publish any part of the paper in connection with any other work by them, provided adequate acknowledgement is given.
4. If it is appropriate, the Authors' employer may sign this Declaration. It is understood that proprietary rights, with the exceptions of Copyright and Patent Rights are reserved by the signee.
5. If an Author is a U.S. Government employee and the work was done in that capacity, the assignment applies only to the extent allowed by U.S. law. If an Author is an employee of the British Government, HMSO will grant a non-exclusive licence to publish the paper in the Journal, provided British Crown Copyright and user rights (including Patent Rights) are reserved.
6. If for good reason a co-author is unable to sign this Declaration, the other Author or co-authors may sign on his or her behalf, provided that this is clearly stated and on the understanding that they will make every effort to inform the person concerned of the terms of the agreement.

When submitting a manuscript for editorial consideration, Authors should confirm their acceptance of these terms by signing a Declaration to that effect. The recommended wording is given in the example. No paper will be accepted for publication without such a Declaration being signed by each Author (see paragraph 6 above). If the manuscript is not accepted for publication, the assignment will be null and void.

ABBREVIATIONS AND SYMBOLS**Physico-chemical quantities**

The *British Journal of Pharmacology* uses the SI symbols for units. The following prefixed for multiples of units should be used:

<i>Multiplier</i>	<i>Prefix</i>	<i>Symbol</i>
10^{-1}	deci	d
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p
10^{-15}	femto	f
10^{-18}	atto	a

<i>Multiplier</i>	<i>Prefix</i>	<i>Symbol</i>
10^3	kilo	k
10^6	mega	M
10^9	giga	G
10^{12}	tera	T

Thus, micron = μ m; ångstrom = 0.1 nm. Mixed prefixes are not permissible, thus m μ g should be ng. The symbols d (10^{-1}) and c (10^{-2}) should be restricted to those occasions on which there is a strongly felt need for them (e.g. cm).

Use of the solidus

The solidus should be avoided as far as possible and the negative index substituted, e.g. mg kg $^{-1}$ rather than mg/kg; pmol mm $^{-2}$ min $^{-1}$ rather than pmol/mm 2 /min.

SYMBOLS

Symbols denoting physical quantities are usually printed as italic capitals (indicated by single underline in typescript). A dash over the symbol indicates a mean value; a dot over the symbol indicates a time derivative. Suffixes may be used to indicate 'where' and 'what'. They are printed as inferiors on the line. Multiple suffixes should be avoided if a simpler symbol adequately defined is unambiguous, but if necessary should be separated by commas e.g. P_{A, CO_2} denotes partial pressure of CO₂ alveolar air.

CHEMICAL AND BIOLOGICAL ABBREVIATIONS

Authors should also consult *Nomenclature Guidelines for Authors* contained in this issue of the Journal. The abbreviations listed may be used without definition *except* those for chemicals, drugs and enzymes which must be written in full at first mention in the title, summary and again in the text. At first mention they should be followed by the abbreviation in brackets. Subsequently, the abbreviation alone may be used.

The list of abbreviations for chemical, drug and enzyme names is clearly not comprehensive and includes only a few commonly used examples.

Use abbreviations sparingly as extensive use can make the text hard to follow.

Physico-chemical quantities

Quantity	Preferred unit	Symbol
Amount (of substance)	mole	mol
Capacitance	farad	F
Concentration	moles per litre	M or mol l ⁻¹
Current	ampere	A
Electrical conductance	siemens	S
Electromotive force	volt	V
Flow (blood or other liquid)	litres per second (or min)	1 s ⁻¹ or 1 min ⁻¹
Flow (air or other gas)	litres per second (or min)	1 s ⁻¹ or 1 min ⁻¹
Force	newton	N
Frequency of regular event	hertz	Hz
Length	metre	m
Mass	gram	g
Power	watt	W
Pressure (or partial pressure)	pascal*	Pa
Radioactivity	becquerel or curie	Bq (60 d.p.m.) or Ci (3.7 × 10 ¹⁰ Bq)
Resistance (electrical)	ohm	Ω
Temperature	degree celsius	°C
Time	second (preferred)	s
	minute	min
	hour	h
Volume (blood or other liquid)	litre	l
Volume (air or other gas)	litre	l
Work	joule	J

* mm of mercury (mmHg) are allowed if conventional, and if mercury manometer is used for calibration.

Chemical and biological abbreviations

acetylcholine	ACh	dextro-(absolute configuration)	D-
acetylcholinesterase	AChE	dextro-(optical rotation)	(+)-
adenosine 3':5'-cyclic monophosphate	cyclic AMP	diameter	diam.
adenosine 5'-phosphate	AMP	diameter, inside	i.d.
adenosine triphosphatase	ATPase	diameter, outside	o.d.
γ-aminobutyric acid	GABA	diffusion coefficient	D
analysis of variants	F	3,4-dihydroxyphenylalanine	DOPA
adrenaline	Ad	3,4-dihydroxyphenylethylamine	dopamine
analytical standard of reagent purity	A.R.	direct current	d.c.
anhydrous	anhyd.	disintegration per minute	d.p.m.
approximate(ly)	approx.	dissociation constant	K _D
approximately equals	≈	dissociation constant, negative logarithm of	pK
aqueous	aq.	distilled	dist.
arg-vasopressin	AVP	dry ice	solid CO ₂
boiling point	b.p.	edition	edn
bovine serum albumin	BSA	editor(s)	ed.
cardiovascular system	CVS	effective concentration	EC ₅₀
catechol-O-methyl transferase	COMT	effective dose, median	ED ₅₀
central nervous system	CNS	electrocardiogram	ECG
cerebrospinal fluid	CSF	electrocorticogram	ECoG
chi-squared (statistics)	χ ²	electroconvulsive therapy	ECT
clearance	c	electroencephalogram	EEG
coenzyme A	CoA	electromyogram	EMG
concentrated	conc.	electron spin resonance	e.s.r.
correlation coefficient	r	endothelial-derived relaxing factor	EDRF
cubic	cu.	epithelial-derived relaxing factor	EpDRF
degree of freedom (statistics)	d.f.	equilibrium constant	K
deoxyribonucleic acid	DNA	equivalent (general use)	equiv.
deoxyribonuclease	DNase	erythrocyte	r.b.c.
		erythrocyte sedimentation rate	ESR
		ethylenediaminetetraacetic acid	EDTA
		excitatory postsynaptic potential	e.p.s.p.
		experiment	expt
		experimental	exptl

fatty acids, nonesterified	NEFA	page/pages	p./pp.
figure(s) (with reference number)	Figure(s)	para-	p-
figure (diagram)	figure	paragraph	para. or ¶
gas-liquid chromatography	g.l.c.	parts per million	p.p.m.
glomerular filtration rate	GFR	per cent	%
haemoglobin	Hb	platelet activating factor	PAF
half-life	$t_{\frac{1}{2}}$	posterior	post.
high-frequency	h.f.	probability (significance level in a statistical test)	P
high performance liquid chromatography	h.p.l.c.		
human serum albumin	HSA	radioimmunoassay	RIA
hydrogen-ion concentration	[H ⁺]	rectus (configuration by the sequence rule)	R
hydrogen-ion activity, negative logarithm of (hydrogen-ion exponent)	pH	red blood corpuscle	RBC
6-hydroxydopamine	6-OHDA	relative band speed to front (chromatography)	R_F
N-[2-Hydroxyethyl]piperazine-N'- [2-ethanesulphonic acid]	HEPES	relative molecular mass	M_r
5-hydroxyindoleacetic acid	5-HIAA	relative retention time (gas chromatography)	t_r
5-hydroxytryptamine	5-HT	renal plasma flow	RPF
immunoglobulins	IgA, IgD, IgE, IgG, IgM	resistance (respiratory)	R
inhibitor constant	K_i	respiratory conductance	Sgaw
inhibitory concentration	IC ₅₀	revolutions per minute	r.p.m.
inhibitory postsynaptic potential	i.p.s.p.	ribonucleic acid	RNA
insoluble	insol.		
international unit	iu	section	§
intra-arterial	i.a.	sedimentation coefficient	s
intracellular fluid	ICF	(ultracentrifugation)	
intradermal	i.d.	sinister (configuration by the sequence rule)	S
intramuscular	i.m.	soluble	sol.
intraperitoneal	i.p.	solution	soln.
intracerebroventricular	i.c.v.	Spearman rank coefficient	r_s
intravenous	i.v.	standard deviation	s.d.
isotope (atomic mass)	¹³¹ I	(of observed sample)	
e.g. iodine-131		standard error (of estimate mean value)	s.e.mean
isotopically substituted compounds e.g.	[¹⁴ C]-ethanol	standard error (of sampling)	s.e.
laevo-(absolute configuration)	L-	standard temperature and pressure	STP
laevo-(optical rotation)	(-)-	subcutaneous	s.c.
lethal dose, median	LD ₅₀	sum (statistical):	
leukotriene	LT	of hypothetical population	Σ
logarithm to base e	log _e or ln	of observed sample	S or Σ
logarithm to base 10	log ₁₀	temperature	temp.
maximum	max.	thin layer chromatography	t.l.c.
mean arterial pressure	MAP	time, clock – 24 h clock used	t
mean value of (statistics)	\bar{x}	e.g. 18 h 30 min	
melting point	m.p.	time constant	τ
meta	m-	2-amino-2-hydroxymethyl- propan-1,3-diol	Tris
Michaelis constant	K_M	ultraviolet	u.v.
minimum	min.	unit	u
mobility (electrophoresis)	m	vacuum	vac.
monoamine oxidase	MAO	valency	e.g. Fe ²⁺ ; Fe(II) protoporphyrin
noradrenaline	NA		
nuclear magnetic resonance	n.m.r.		
number	no. or No.	volume by volume	v/v
number of observations (statistics)	n	wavelength	λ
ortho	o-	weight	wt.
packed cell volume	PCV	weight by volume	w/v

NOMENCLATURE GUIDELINES FOR AUTHORS

With effect from 1 January 1995

The Nomenclature Working Party (NWP) of the Editorial Board of the *British Journal of Pharmacology* has consulted many acknowledged experts in an effort to clarify and standardize receptor and other nomenclature systems for use by Editors until a complete set of recommendations of the IUPHAR Committee on Receptor Nomenclature and Drug Classification (NC-IUPHAR) are published.

NWP is unanimous in its view that, with rare exceptions, the Journal should use spellings, names and abbreviations that have been chosen by international bodies convened for the purpose.

For receptor nomenclature, with few exceptions, the Journal generally follows the guidelines laid down in the current *Trends in Pharmacological Sciences* Receptor and Ion Channel Nomenclature Supplement and the reports of NC-IUPHAR published in *Pharmacological Reviews*.

1 Definition of receptors and subtypes

Receptors and their subtypes are defined in terms of the relative potencies of agonists and selectivities of antagonists in functional studies, by the binding of such ligands and to structural information, where available.

2 Format of receptor names

It was agreed that, until NC-IUPHAR provides full recommendations:

- (a) Editors will permit with reluctance new nomenclature systems in papers accepted for publication if and only if there are compelling reasons to introduce a new terminology (or modify an accepted one). The criteria upon which the new receptor type or subtype are defined must be given, together with adequate explanations of the relationship between the previous nomenclature (fully referenced) and the proposed one.
- (b) Only well-established and universally accepted subtype names (e.g. muscarinic and nicotinic acetylcholine receptors; α - and β -adrenoceptors) will be acceptable without any reference to the originator of these terms. In cases of controversy concerning further subdivision of the subtype, full referencing must be given.
- (c) When receptors are expressed from DNA/RNA that has been introduced into cells and these receptors display a similar pharmacological profile to the native receptors, they should be denoted by use of lower case, e.g. m_1 for expressed receptor and M_1 for native receptor. The stoichiometry of the expressed receptor should be indicated, where appropriate, e.g. for an immature muscle nicotinic acetylcholine receptor, it might be $(\alpha 1)_2 \beta 1 \gamma \delta$.
- (d) Receptor subtypes should be designated by means of a subscript numeral or capital letter. Some double subscripts (i.e. numerical plus letter) are acceptable.

3 Types of receptor

The NWP accepts that there are additional receptors to those described below which can be considered as well

established. In many cases, however, their existence has been confirmed only in cloning studies and it is as yet unclear how they relate to similar subdivisions proposed on the grounds of differences in agonist and antagonist potencies in various tissues.

- (a) *Acetylcholine receptors* The two principal subfamilies are muscarinic and nicotinic acetylcholine receptors (the abbreviations muscarinic AChRs and nicotinic AChRs are acceptable, but not mAChR or nAChR, which may be confused with muscle and neuronal subtypes).
- (b) *Muscarinic acetylcholine receptors* The principle subtypes are M_1 , M_2 , M_3 and M_4 .
- (c) *Nicotinic acetylcholine receptors* The principal subgroups are muscle and neuronal nicotinic acetylcholine receptors.
- (d) *Adenosine receptors* Known also as P_1 purinoceptors (see purinoceptors, 3u). (See Fredholm, B.B., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 145–156.)
- (e) *Adrenoceptors* The principal subtypes are α_1 -, α_2 -, β_1 -, β_2 - and β_3 -adrenoceptors. Additional subtypes must be fully referenced. (See Bylund, D.B., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 121–136.)
- (f) *Angiotensin receptors* The principle subtypes are AT_1 and AT_2 .
- (g) *Bombesin receptors* Proposed subtypes such as BB_1 , BB_2 may be used but must be fully referenced.
- (h) *Bradykinin receptors* The principle subtypes are B_1 and B_2 receptors.
- (i) *Calcitonin gene-related peptide (CGRP) receptors* Proposed CGRP receptor subtypes must be fully referenced.
- (j) *Cannabinoid receptors* These should follow the style CB_1 , CB_2 and be fully referenced.
- (k) *Cholecystokinin (CCK) receptors* The principle subtypes are CCK_A and CCK_B receptors.
- (l) *Dopamine receptors* D_1 and D_2 dopamine receptors are recognised. Other subtypes must be fully referenced.
- (m) *Endothelin receptors* The principle subtypes are ET_A and ET_B receptors. (See Masaki, T., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 137–142.)
- (n) *Excitatory amino acid receptors* Three ionotropic subtypes are recognised and named: (1) NMDA receptors; (2) AMPA receptors, and (3) kainate receptors. A second class is the metabotropic (mGlu) receptor family. Further subtypes of excitatory amino acid receptors must be fully referenced.
- (o) *γ -Aminobutyric acid (GABA) receptors* The principal subtypes are $GABA_A$ and $GABA_B$ receptors. Regulatory sites on the $GABA_A$ receptor should be referenced.
- (p) *Histamine receptors* The principle subtypes are H_1 , H_2 and H_3 receptors.
- (q) *5-Hydroxytryptamine (5-HT) receptors* The principle subtypes are $5-HT_1$, $5-HT_2$, $5-HT_3$ and $5-HT_4$. Further

subdivisions require full referencing. (See Hoyer, D., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 157–203).

(p) *Leukotriene receptors* When first mentioned, the style leukotriene (LT) receptor should be used, thereafter LT receptor. Receptors should be designated according to the leukotriene that selectively or preferentially binds to them. All leukotriene receptor subtypes should be fully referenced.

(q) *Neuropeptide Y (NPY) receptors* Proposed subtypes should be fully referenced.

(r) *Opioid receptors* The principal subtypes are μ -, δ - and κ -opioid receptors. Other proposed subtypes should be fully referenced.

(s) *Oxytocin receptors* (see Vasopressin and oxytocin receptors).

(t) *Prostanoid receptors* The principal types are DP, EP, FP, IP and TP receptors. When first mentioned, the style prostanoid (XP) receptor should be used, thereafter XP receptor (where X denotes the type). Proposed subtypes should be referred to as XP_n , (e.g. EP₁, EP₂) and referenced. (See Coleman, R.A., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 205–229).

(u) *Purinoceptors* The principal subtypes are P₁ and P₂ receptors. Subdivision of P₁ into A₁, A₂ and A₃ subtypes and of P₂ into P_{2X} and P_{2Y} are permitted. Other subtypes e.g. P_{2T}, P_{2Z} should be fully referenced. (See Fredholm, B.B., *et al.*, (1994) *Pharmacol. Rev.*, **46**, 143–156.)

(v) *Somatostatin (SS) receptors* Proposed subtypes should be fully referenced.

(w) *Tachykinin receptors* The term tachykinin is preferred to neurokinin. The principle subtypes are NK₁, NK₂ and NK₃ receptors.

(x) *Vasoactive intestinal peptide (VIP) receptors* Proposed subtypes should be fully referenced.

(y) *Vasopressin and oxytocin receptors* The principle subtypes are V_{1A}, V_{1B}, V₂ and OT receptors.

4 Naming of ion Channels

The four main ion channels are named according to the abbreviation of the ion normally carrying the current (i.e. K channels, Na channels, Ca channels and Cl channels). In naming channels the charge on the ion is understood and need not be stated. Further subtypes, e.g. K_{ATP} must be fully referenced. For Ca channels, see Spedding, M. & Paoletti, P. (1992) *Pharmacol. Rev.*, **44**, 363–376.

5 Naming of nerve fibres

Many nerve fibres are now known to release more than one transmitter, and future work may show that this is in fact the general rule. In that case, the concept of the same transmitter being released either at different developmental stages or under various experimental conditions would no longer hold, and single adjectives that imply this (e.g. cholinergic, noradrenergic) would become inappropriate when applied to nerve fibres, as distinct from transmitter functions. For the present, those nerve fibres that are known to function by releasing more than one identified transmitter may be described accordingly; for example, noradrenergic-purinergic, cholinergic-peptidergic (in alphabetical order, the order implying no priority of function). N.B. The suffix 'ergic' should continue to be applied only to nerve fibres and to the transmission event, in accordance with Dale's intentions. For example, 'cholinergic' indicates that the nerve fibre, or the transmission, functions under particular conditions through the release of a

choline-like substance. The suffix should not be used loosely to mean 'pertaining to'. Hence, for example, the expression 'cholinergic receptor' (rather than acetylcholine receptor) is an inappropriate use of the term. Transmission events involving nitric oxide may be referred to as *nitrenergic*. However, *nitrenergic* may be used to describe axons only when there is sufficient evidence that nitric oxide is released from them as a neurotransmitter.

(a) *Catecholamine releasing nerve fibres* The adjective to be applied to nerve fibres that release dopamine as a transmitter is dopaminergic (not DAergic).

Nerve fibres that are known to function by releasing noradrenaline are to be described as noradrenergic. The term adrenergic should now be reserved for nerve fibres known to release adrenaline. Where the identity of the catecholamine is uncertain, catecholaminergic should be used.

(b) *Some other adjectives describing nerve fibre function* NANC is an acceptable abbreviation of non-adrenergic, non-cholinergic for peripheral efferent nerve fibres when the identity of the transmitter(s) is unknown other than the fact that neither (nor)-adrenaline nor acetylcholine is involved. It should be defined when introduced. NANCergic, e-NANC (or NANC-e) and i-NANC (or NANC-i) are not acceptable terms.

Glutamatergic, not glutaminergic, should be used to describe nerve fibres releasing glutamate. In referring to peptide-releasing nerve fibres (e.g. those that may release substance P or vasoactive intestinal peptide) the nomenclature to be used is peptidergic (X), e.g. peptidergic (SP), peptidergic (VIP), not SPergic, VIPergic.

The terms 5-hydroxytryptamine (5-HT) and 5-hydroxytryptaminergic (i.e. nerves releasing 5-hydroxytryptamine) are preferred to those of serotonin and serotoninergic. The term 5-HTergic is not acceptable, except to avoid frequent repetition of 5-hydroxytryptaminergic.

Likewise, the terms purinergic (ATP) and purinergic (adenosine) are preferred.

6 Terms used to describe agonist and antagonist action

The following terms can be used without full definition. Where appropriate, other terms may be used but must be accompanied by a full definition.

Terms used to describe affinity and potency

(a) *EC₅₀* The concentration of an agonist that produces 50% of the maximal response for that agonist *in vitro*. The agonist may be stimulatory or inhibitory. When EC₅₀ values are determined in the presence of other agonists or antagonists the concentration of the latter should be stated. Related terms, e.g. EC₂₅, are acceptable if accompanied by a full definition.

(b) *IC₅₀* This term may be used in the following ways: (i) The concentration of antagonist that reduces the response to a sub-maximal concentration of agonist by 50%; the concentration of agonist should be stated. (ii) The concentration of competing agonist or antagonist that inhibits the binding of a radioligand by 50%; the concentration of radioligand should be stated.

(c) *ED₅₀* This term may be used in the following ways: (i) The dose of an agonist or antagonist that produces 50% of the maximal possible effect of that agonist or antagonist *in vitro*. (ii) The dose of drug that produces the effect under investigation in 50% of the population.

- (d) K The dissociation equilibrium constant (mol l^{-1}) for ligand-receptor interactions. The reciprocal is called the affinity constant or association equilibrium constant. When necessary for clarity subscripts (letters or numerals, or a combination of both) may be added but these must be clearly explained when first used.
- (e) n_H The Hill coefficient.
- (f) pA_2 The negative logarithm to base 10 of the concentration of an antagonist that makes it necessary to double the concentration of agonist needed to elicit a given submaximal response. Note that the definition is empirical and does not pre-suppose the mechanism of antagonism. The pA_2 value can be determined from a Schild plot with unconstrained slopes, but only provides an estimate of the pK_B if the antagonism has been shown to meet all of the criteria of competition.
- (g) pD_2 The negative logarithm to base 10 of the EC_{50} .
- (h) pIC_{50} The negative logarithm to base 10 of the IC_{50} .
- (i) pK The negative logarithm to base 10 of K (with or without subscripts as appropriate: see 6(d)).

Terms used to describe the mode of antagonism

- (a) *Competitive antagonism* Used to describe antagonists that bind reversibly to the agonist binding site.
- (b) *Irreversible competitive antagonism* Used to describe antagonists that bind irreversibly to the agonist binding site.
- (c) *Non-competitive* Used to describe antagonists that bind reversibly to a distinct (allotopic) site.
- (d) *Irreversible non-competitive antagonism* Used to describe antagonists that bind irreversibly to a distinct (allotopic) site.

7 Enzymes

The IUBMB Enzyme Commission (EC) number and full name (Enzyme Nomenclature 1992, Academic Press, San Diego) must be quoted when first mentioned in text. Subsequently the accepted trivial name is used. Trivial names may be used in the title.

8 Other nomenclature requirements

- (a) *Racemates* Authors must state unambiguously in the Methods section of papers which isomers were used, e.g. (+)- or (-)-propranolol, and must bring to the attention of the reader the composite character of drugs that are mixtures of stereoisomers. Further-

more, the implications of the composite nature of such drugs studied for the interpretation of the data measured and the conclusions drawn must be made explicit. Note that the terms d- or l- for dextro- and laevo-rotatory are now obsolete, and the prefixes (+)- or (-)- respectively should be used. Capital D and L refer to the absolute configurations and of course remain acceptable when appropriate.

- (b) *Purines* This term should not be used as a synonym for purine nucleotides or nucleosides.
- (c) *Eicosanoids* The system of nomenclature to be used for eicosanoids is that published in *Methods in Enzymology*, (1990), 187, 1-9. This scheme incorporates recent changes in the style of abbreviation of hydroperoxy-, epoxy- and oxo-unsaturated fatty acids e.g. 12(S)-hydroperoxyeicosatetraenoic acid which was formerly abbreviated as 12(S)-HPETE now becomes 12(S)-HpETE. In manuscripts, the first use of the full chemical name of any eicosanoid should indicate double bond geometry when this is known.
- (d) *Peptide nomenclature* The preferred style is capital letters to designate the first letter of the word. Otherwise, upper and lower case letters should be used (e.g. Enk-IR, enkephalin-like immunoreactivity). When numbers are used these should be placed after a hyphen on the same line as the abbreviation, e.g. ET-1.
- (e) *Cell lines* Cell type, sources and originating species need to be defined.
- (f) *Molecular biology* Abbreviations pertaining to molecular biological techniques need to be defined or presented in such a way that they can be recognised by the non-specialist e.g. the oligonucleotide sequence, TAGC.
- (g) *Tension* Tension is force and should be calibrated in newtons (1 newton = 1 kg m s^{-1}) or in kg weight, g weight, or mg weight etc. It should not be calibrated in units of mass (e.g. kg). (See Miller D.J. in *Trends Pharmacol. Sci.*, 1988, 9, 124-5).
- (h) *Ions* When referring to ions, the charge should be indicated. e.g. Na^+ , Ca^{2+} , $2\text{Na}^+/\text{Ca}^{2+}$ exchange, etc.
- (i) *Inhibitors of nitric oxide synthase* The most commonly used and currently accepted abbreviation for N^{C} -nitro-L-arginine and N^{G} -nitro-L-arginine methyl ester are L-NOARG and L-NAME. Unless alternative international agreement is reached, these will be the abbreviations accepted by the journal.

INDEX TO VOLUMES 111, 112 AND 113

AUTHOR INDEX

A

Aasen, A.J. *see* Onali, P., 113, 775
 Abassi, Z.A., Golomb, E., Agbaria, R., Roller, P.P., Tate, J. and Keiser, H.R. Hydrolysis of iodine labelled urodilatin and ANP by recombinant neutral endopeptidase EC 3.4.24.11, 113, 204
 Abboe-Offei, B. *see* Moreland, S., 112, 704
 Abdul-Hussein, N. *see* Einstein, R., 111, 199
 Acar, D. *see* Rump, A.F.E., 112, 757
 Adachi-Akahane, S. *see* Okuyama, R., 113, 451
 Adams, D.J. *see* Cuevas, J., 111, 663
 Adeagbo, A.S.O., Tabrizchi, R. and Triggle, C.R. The effects of perfusion rate and N^G -nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
 Adeagbo, A.S.O. *see* Pak, J.-M., 112, 1209
 Adgey, A.A.J. *see* Herity, N.A., 112, 423
 Adner, M. *see* Ekelund, U., 112, 887
 Adzet, T. *see* Alegret, M., 112, 551
 Agbaria, R. *see* Abassi, Z.A., 113, 204
 Aghai, Z. *see* Bax, W.A., 113, 1471
 Agvald, P. *see* Persson, M.G., 111, 825
 Ahluwalia, A., Maggi, C.A., Santicioli, P., Lecci, A. and Giuliani, S. Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017
 Ahmad, S. *see* Laniyonu, A., 113, 403
 Akaike, N. *see* Nishikawa, M., 112, 1160
 Akaike, N. *see* Uneyama, C., 112, 349
 Akaike, N. *see* Uneyama, H., 111, 245
 Akarasereenont, P., Mitchell, J.A., Appleton, I., Thiemermann, C. and Vane, J.R. Involvement of tyrosine kinase in the induction of cyclooxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
 Akarasereenont, P. *see* Mitchell, J.A., 113, 1008
 Akbar, M.T. *see* Shoaib, M., 111, 1073
 Aldasoro, M. *see* Martínez, M.C., 113, 419
 Alegret, M., Ferrando, R., Vázquez, M., Adzet, T., Merlos, M. and Laguna, J.C. Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551
 Alemany, R. *see* Olmos, G., 111, 997
 Alemayehu, A., Lock, K.R., Coatney, R.W. and Chou, C.C. L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205
 Alexander, B. *see* Browse, D.J., 113, 987
 Alexander, S.P.H., Curtis, A.R., Kendall, D.A. and Hill, S.J. A₁ adenosine receptor inhibition of cyclic AMP formation and radioligand binding in the guinea-pig cerebral cortex, 113, 1501
 Alexander, S.P.H., Losinski, A., Kendall, D.A. and Hill, S.J. A comparison of A₂ adenosine receptor-induced cyclic AMP generation in cerebral cortex and relaxation of pre-contracted aorta, 111, 185
 Alexander, S.P.H. *see* Cartmell, J., 111, 364
 Alexander, S.P.H. *see* Hernández, F., 113, 216
 Alexander, S.P.H. *see* Iredale, P.A., 111, 1252
 Allawi, H.S., Wallace, P., Pitcher, A., Gaffen, Z., Bland-Ward, P.A. and Moore, P.K. Effect of 7-nitro indazole on neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
 Allcock, G.H. *see* Warner, T.D., 112, 207
 Alldus, C. *see* Buchan, K.W., 112, 1251
 Allen, G.T. *see* Tesfamariam, B., 112, 55
 Allen, J.D. *see* Herity, N.A., 112, 423
 Allen, S.P. *see* Dashwood, M.R., 112, 386
 Altavilla, D. *see* Squadrato, F., 113, 912
 Altavilla, D. *see* Squadrato, F., 113, 1153
 Amenta, F. *see* Lakhani, P.P., 111, 1296

Amin, F. *see* Makki, T., 111, 555
 Amirchett-Rao, S. *see* Wright, I.K., 112, 1146
 Ammendolia, L. *see* Squadrato, F., 113, 912
 Ammendolia, L. *see* Squadrato, F., 113, 1153
 Anagnostopoulos, T. *see* Belachgar, F., 112, 465
 Anderson, G.P. *see* Wessler, I., 113, 1221
 Anderson, R., Sheehan, M.J. and Strong, P. Characterization of the adenosine receptors mediating hypothermia in the conscious mouse, 113, 1386
 Andersson, K.-E. *see* Ishizuka, O., 113, 111
 Andriopoulou, P. *see* Pipili-Synetos, E., 111, 894
 Angus, J.A. *see* Chin, J.P.F., 112, 566
 Ansay, M. *see* Delaunois, A., 113, 973
 Antunes, E. *see* Grassi-Kassisse, D.M., 113, 675
 Antunes, E. *see* Lopes-Martins, R.A.B., 113, 81
 Anwyl, R. *see* Manahan-Vaughan, D., 112, 1083
 Aoki, H., Kobayashi, S., Nishimura, J. and Kanaide, H. Sensitivity of G-protein involved in endothelin-1-induced Ca^{2+} influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
 Aoki, H., Nishimura, J., Kobayashi, S. and Kanaide, H. Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
 Appleton, I. *see* Akarasereenont, P., 113, 1522
 Appleton, I. *see* Tomlinson, A., 113, 693
 Arakida, Y. *see* Watanabe, M., 111, 123
 Argenbright, L.W. *see* Norman, K.E., 111, 117
 Arias-Montafio, J.A., Berger, V. and Young, J.M. Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
 Arimura, A., Asanuma, F., Kurosawa, A. and Harada, M. Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339
 Artaiz, I. *see* Zazpe, A., 113, 968
 Arvola, P. *see* Kähönen, M., 112, 925
 Arvola, P. *see* Vuorinen, P., 112, 71
 Asai, T. *see* Mukai, J., 111, 389
 Asano, M. *see* Yokota, Y., 113, 324
 Asano, T. *see* Nakazato, Y., 111, 935
 Asano, Y. *see* Muto, S., 111, 173
 Asanuma, F. *see* Arimura, A., 111, 339
 Ashcroft, F.M. *see* Smith, P.A., 112, 143
 Ashford, M.L.J. *see* Lee, K., 111, 632
 Ashford, M.L.J. *see* Sellers, A.J., 113, 659
 Askew, S. *see* Salas, E., 112, 1071
 Astolfi, M., Treggiari, S., Giachetti, A., Meini, S., Maggi, C.A. and Manzini, S. Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570
 Atcheson, R., Lambert, D.G., Hirst, R.A. and Rowbotham, D.J. Studies on the mechanism of [³H]-noradrenaline release from SH-SY5Y cells: the role of Ca^{2+} and cyclic AMP, 111, 787
 Atkinson, J., Tatchum-Talom, R. and Capdeville-Atkinson, C. Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat, 111, 1184
 Atkinson, J. *see* Makki, T., 111, 555
 Atkinson, J. *see* Thorin, E., 111, 351
 Atkinson, J. *see* Thorin, E., 113, 363
 Atwal, P. *see* Hunter, J.C., 112, 1002
 Aupetit-Faisant, B. *see* Moreau, R., 112, 649
 Austin, C.E., Foreman, J.C. and Scadding, G.K. Reduction by Hoe 140, the B₂ kinin receptor antagonist, of antigen-induced nasal blockage, 111, 969
 Aylett, A.J. *see* Balmforth, A.J., 112, 277
 Azzurri, M. *see* Passani, M.B., 112, 695

B

Bacman, S., Borda, E., Denduchis, B., Lustig, L. and Sterin-Borda, L. Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271

Baggett, N. *see* Shaunak, S., 113, 151

Bahl, A.K., Dale, M.M. and Foreman, J.C. The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809

Bailey, C.J. and Mynett, K.J. Insulin requirement for the antihyperglycaemic effect of metformin, 111, 793

Bailey, C.J., Mynett, K.J. and Page, T. Importance of the intestine as a site of metformin-stimulated glucose utilization, 112, 671

Bailey, S.J. and Hourani, S.M.O. Differential effects of suramin on P₂-purinoreceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219

Bailey, T.J. *see* Regan, J.W., 112, 377

Baines, C.P., Szwarc, R.S. and Ball, H.A. Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334

Baird, A.W. *see* Brayden, D.J., 113, 593

Baldwin, H.A., Williams, J.L., Snarey, M., Ferreira, T., Cross, A.J. and Green, A.R. Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188

Balfour, D.J.K. *see* Shoaib, M., 111, 1073

Ball, H.A. *see* Baines, C.P., 113, 334

Ball, H.A. *see* Yard, N.J., 113, 1001

Ball, S.G. *see* Balmforth, A.J., 112, 277

Ballarín, M., Carceller, A. and Guitart, X. Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

Ballieux, R.E. *see* Benschop, R.J., 113, 1311

Balmforth, A.J., Bryson, S.E., Aylett, A.J., Warburton, P., Ball, S.G., Pun, K.-T., Middlemiss, D. and Drew, G.M. Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Bara, A.T. *see* Hecker, M., 111, 238

Bara, A.T. *see* Pörsti, I., 111, 652

Baran, H. *see* Löscher, W., 112, 97

Barber, A., Bartoszyk, G.D., Bender, H.M., Gottschlich, R., Greiner, H.E., Harting, J., Maufer, F., Minck, K.-O., Murray, R.D., Simon, M. and Seyfried, C.A. A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 61753, 113, 1317

Barber, A., Bartoszyk, G.D., Greiner, H.E., Maufer, F., Murray, R.D., Seyfried, C.A., Simon, M., Gottschlich, R., Harting, J. and Lues, I. Central and peripheral actions of the novel κ-opioid receptor agonist, EMD 60400, 111, 843

Barbier, A.J.M. and Lefebvre, R.A. Influence of S-nitrosothiols and nitrate tolerance in the rat gastric fundus, 111, 1280

Barbieri, A. *see* Lucchelli, A., 112, 763

Barclay, P.L. *see* Wong, P.S.K., 112, 881

Barelli, H., Fox-Threlkeld, J.E.T., Dive, V., Daniel, E.E., Vincent, J.P. and Checler, F. Role of endopeptidase 3.4.24.16 in the catabolism of neuropeptides, *in vivo*, in the vascularly perfused dog ileum, 112, 127

Barnes, J.M., Dev, K.K. and Henley, J.M. Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339

Barnes, P.J. *see* Belvisi, M.G., 111, 103

Barnes, P.J. *see* Mitchell, J.A., 113, 1008

Barnes, P.J. *see* Ramnarine, S.I., 113, 1183

Barnes, P.J. *see* Ward, J.K., 111, 1095

Barnette, M. *see* Grous, M., 111, 259

Barnouin, M.C. *see* Dubuc, I., 112, 352

Bartfai, T. *see* Kahl, U., 111, 1129

Bartho, L. *see* Maggi, C.A., 112, 161

Bartolini, A. *see* Malmberg-Aiello, P., 111, 1269

Bartoszyk, G.D. *see* Barber, A., 111, 843

Bartoszyk, G.D. *see* Barber, A., 113, 1317

Barturen, F. *see* García-Vallejo, P., 111, 1140

Bast, A. *see* Jansen, F.P., 113, 355

Battistini, B., O'Donnell, L.J.D., Warner, T.D., Fournier, A., Farthing, M.J.G. and Vane, J.R. Characterization of endothelin (ET) receptors in the isolated gall bladder of the guinea-pig: evidence for an additional ET receptor subtype, 112, 1244

Battistini, B., Warner, T.D., Fournier, A. and Vane, J.R. Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

Battram, C.H. *see* Raeburn, D., 113, 1423

Bauersachs, J., Hecker, M. and Busse, R. Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548

Baumann, P.A. *see* Waldmeier, P.C., 113, 1515

Bäumert, H.G. *see* McLaren, G.J., 111, 913

Bäumert, H.G. *see* Windscheif, U., 113, 1015

Bäumert, H.G. *see* Ziganshin, A.U., 111, 923

Bax, W.A., Aghai, Z., van Tricht, C.L.J., Wassenaar, C. and Saxena, P.R. Different endothelin receptors involved in endothelin-1- and sarafotoxin-S6B-induced contractions of the human isolated coronary artery, 113, 1471

Baxter, G.S., Murphy, O.E. and Blackburn, T.P. Further characterization of 5-hydroxytryptamine receptors (putative 5-HT_{2B}) in rat stomach fundus longitudinal muscle, 112, 323

Baxter, G.S. *see* Wardle, K.A., 112, 789

Baxter, K. *see* Church, J., 113, 499

Baydoun, A.R., Bogle, R.G., Pearson, J.D. and Mann, G.E. Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487

Béa, M.-L., Ghaleh, B., Giudicelli, J.-F. and Berdeaux, A. Lack of importance of NO in β-adrenoceptor-mediated relaxation of large epicardial canine coronary arteries, 111, 981

Beattie, D.T. and Connor, H.E. The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

Becherucci, C. *see* Palmi, M., 112, 449

Beck, J.K. *see* Kojda, G., 112, 533

Belachgar, F., Hulin, P., Anagnostopoulos, T. and Planelles, G. Triflocin, a novel inhibitor for the Na-HCO₃ symport in the proximal tubule, 112, 465

Belai, A. *see* Kasakov, L., 112, 403

Bélichard, P. *see* Pruneau, D., 111, 1029

Bell, B.A. *see* Stapleton, S.R., 111, 57

Bell, N.J. *see* Trezise, D.J., 113, 463

Bell-Quilley, C.P. *see* DelliPizzi, A., 111, 1

Belvisi, M.G., Patacchini, R., Barnes, P.J. and Maggi, C.A. Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103

Belvisi, M.G. *see* Mitchell, J.A., 113, 1008

Belvisi, M.G. *see* Ward, J.K., 111, 1095

Bender, H.M. *see* Barber, A., 113, 1317

Benetos, A. *see* Bouaziz, H., 113, 717

Benjamin, I.S. *see* Browse, D.J., 113, 987

Bennett, J. *see* Delamere, F., 111, 983

Bennett, M.R. *see* Nicolarakis, P.J., 111, 521

Bennett, T. *see* Gardiner, S.M., 111, 589

Bennett, T. *see* Gardiner, S.M., 112, 477

Bennett, T. *see* Gardiner, S.M., 112, 823

Bennett, T. *see* Gardiner, S.M., 113, 129

Bennett, T. *see* Kemp, P.A., 111, 325

Bennett, T. *see* Waller, J., 112, 1057

Benschop, R.J., Nijkamp, F.P., Ballieux, R.E. and Heijnen, C.J. The effects of β-adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311

Benwell, M.E.M. *see* Shoaib, M., 111, 1073

Berdeaux, A. *see* Béa, M.-L., 111, 981

Berg, T. *see* Bjørnstad-Østensen, A., 113, 1567

Berger, V. *see* Arias-Montafio, J.A., 111, 598

Bergeron, R. *see* Monnet, F.P., 112, 709

Bergsma, D.J. *see* Schambye, H.T., 113, 331

Bernardi, G. *see* Mercuri, N.B., 113, 831

Bernasconi, S. *see* Tranchina, S., 111, 575

Berry, C.N., Girard, D., Lochot, S. and Lecoffre, C. Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

Berry, M.D., Scarr, E., Zhu, M.-Y., Paterson, I.A. and Juorio, A.V. The

effects of administration of monoamine oxidase-B inhibitors on rat striatal neurone responses to dopamine, 113, 1159

Berryman, K. *see* Davenport, A.P., 111, 4

Berthiaume, N. *see* Claing, A., 112, 1202

Bertrand, C. *see* Ricciardolo, F.L.M., 113, 508

Beukers, M.W. *see* Crack, B.E., 113, 1432

Biessels, G. *see* Kappelle, A.C., 111, 887

Bishop, A.E. *see* Desai, K.M., 113, 1197

Bisset, G.W. Obituary, 111, 649

Bittiger, H. *see* Waldmeier, P.C., 113, 1515

Bjørnstad-Østensen, A. and Berg, T. The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567

Bjur, R.A. *see* Shinozuka, K., 113, 1203

Bkaily, G. *see* Claing, A., 112, 1202

Black, J.W. *see* Welsh, N.J., 112, 93

Blackburn, T.P. *see* Baxter, G.S., 112, 323

Blackburn, T.P. *see* Kennett, G.A., 111, 797

Blake, D.R. *see* Cruwys, S.C., 113, 940

Blake, D.R. *see* Walsh, D.A., 112, 435

Blanc, T. *see* Richard, V., 113, 869

Bland-Ward, P.A. *see* Allawi, H.S., 113, 282

Blier, P. and Bouchard, C. Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

Blin, N., Nahmias, C., Drumare, M.F. and Strosberg, A.D. Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911

Bo, X., Fischer, B., Maillard, M., Jacobson, K.A. and Burnstock, G. Comparative studies on the affinities of ATP derivatives for P_{2X} -purinoceptors in rat urinary bladder, 112, 1151

Boarder, M.R. *see* Purkiss, J.R., 111, 723

Boarder, M.R. *see* Purkiss, J.R., 111, 1041

Boden, P. and Woodruff, G.N. Presence of NK₃-sensitive neurones in different proportions in the medial habenula of guinea-pig, rat and gerbil, 112, 717

Boden, P.R., Pinnock, R.D., Pritchard, M.C. and Woodruff, G.N. Evaluation of a series of novel CCK_B antagonists using a functional assay in the rat central nervous system, 112, 666

Boden, P.R. and Woodruff, G.N. Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429

Bodinier, M.C. *see* Galizzi, J.P., 113, 389

Boeckxstaens, G.E. *see* De Man, J.G., 112, 341

Boeijinga, P.H. *see* Leishman, D.J., 111, 318

Bogardus, A.M. *see* Regan, J.W., 112, 377

Bogle, R.G., Whitley, G.St.J., Soo, S.-C., Johnstone, A.P. and Vallance, P. Effect of anti-fungal imidazoles on mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

Bogle, R.G. *see* Baydoun, A.R., 112, 487

Böhm, D. *see* Böhm, M., 112, 137

Böhm, M., Weinhold, C., Schwinger, R.H.G., Müller-Ehmsen, J., Böhm, D., Reichenspurner, H., Reichart, B. and Erdmann, E. Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137

Böhm, M. *see* Hauck, R.W., 113, 1043

Boksa, P. *see* Clarke, P.B.S., 111, 397

Bolser, D.C., DeGennaro, F.C., O'Reilly, S., Chapman, R.W., Kreutner, W., Egan, R.W. and Hey, J.A. Peripheral and central sites of action of GABA_B agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344

Bolton, T.B. *see* Muraki, K., 111, 1154

Bonci, A. *see* Mercuri, N.B., 113, 831

Bonafous, C., Scatton, B. and Buéno, L. Benzodiazepine-induced intestinal motor disturbances in rats: mediation by ω_2 (BZ₂) sites on capsaicin-sensitive afferent neurones, 113, 268

Bor, P. *see* Szilvassy, Z., 112, 999

Borda, E. *see* Bacman, S., 111, 271

Borgeat, P. *see* McDonald, P.P., 111, 852

Borgeat, P. *see* Marleau, S., 112, 654

Borghì, A. *see* Gallico, L., 112, 795

Bornstein, J.C. *see* Johnson, P.J., 111, 1240

Bornstein, J.C. *see* Yuan, S.Y., 112, 1095

Bouaziz, H., Joulin, Y., Safar, M. and Benetos, A. Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717

Bouchard, C. *see* Blier, P., 113, 485

Boulanger, C.M., Morrison, K.J. and Varhoutte, P.M. Mediation by M₃-muscarinic receptors of both endothelium-dependent contraction and relaxation to acetylcholine in the aorta of the spontaneously hypertensive rat, 112, 519

Boulton, A.A. *see* Zhu, M.-Y., 112, 23

Bousquet, P. *see* Monassier, L., 111, 1347

Bova, S., Rossi, G., Luciani, S., Debetto, P., Pessina, A.C. and Cargnelli, G. Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067

Bowdish, D.M. *see* Low, A.M., 113, 1263

Bowery, B., Rothwell, L.A. and Seabrook, G.R. Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

Bowery, N.G. *see* Facklam, M., 111, 648

Bowery, N.G. *see* Malcangio, M., 113, 635

Bowery, N.G. *see* Malcangio, M., 113, 1561

Bowie, D. and Smart, T.G. Species-dependent functional properties of non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with mammalian and avian brain mRNA, 111, 803

Bowser-Riley, F. *see* Shuker, M.A., 111, 951

Boyer, J.L., Zohn, I.E., Jacobson, K.A. and Harden, T.K. Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614

Bozza, P.T. *see* Pires, A.L.A., 113, 994

Braga da Motta, J.I., Cunha, F.Q., Vargaftig, B.B. and Ferreira, S.H. Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111

Brain, S.D. *see* Hall, J.M., 113, 522

Brain, S.D. *see* Hughes, S.R., 111, 425

Brandt, H.D. *see* Du Plooy, W.J., 111, 73

Braquet, P. *see* McDonald, P.P., 111, 852

Braquet, P. *see* Pires, A.L.A., 113, 994

Bravenboer, B. *see* Kappelle, A.C., 111, 887

Brayden, D.J. and Baird, A.W. A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593

Breittmayer, J.P. *see* Vigne, P., 112, 775

Briand, V., Riva, L. and Galzin, A.M. Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195

Bridal, T.R. *see* de Lorenzi, F.G., 113, 527

Brien, J.F. *see* Hussain, A.S., 113, 1508

Brignola, G., Calignano, A. and Di Rosa, M. Modulation of morphine antinociception in the mouse by endogenous nitric oxide, 113, 1372

Brinjak, R. *see* Karoum, F., 113, 1391

Briscoe, J.A.K. *see* Tammela, T.L.J., 113, 195

Bristow, L.J. *see* Tricklebank, M.D., 113, 729

Broadley, K.J. *see* Martin, S.W., 112, 595

Brouwer, F. *see* Coppes, R.P., 113, 342

Brown, C.A. *see* Chan, S.L.F., 112, 1065

Brown, M.J. *see* Tan, K.K.C., 111, 703

Brown, N. *see* Seabrook, G.R., 112, 728

Brown, N.A., Kemp, J.A. and Seabrook, G.R. Block of human voltage-sensitive Na⁺ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600

Browse, D.J., Mathie, R.T., Benjamin, I.S. and Alexander, B. The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987

Brunn, G. *see* Wessler, I., 113, 1221

Brunner, F. *see* Kühberger, E., 113, 1289

Bryson, S.E. *see* Balmforth, A.J., 112, 277

Buchan, K.W., Alldus, C., Christodoulou, C., Clark, K.L., Dykes, C.W., Sumner, M.J., Wallace, D.M., White, D.G. and Watts, I.S. Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251

Bucher, B. *see* Ouedraogo, S., 111, 625

Bucher, B. *see* Ouedraogo, S., 112, 867

Buéno, L. *see* Bonnafous, C., 113, 268

Buéno, L. *see* Gué, M., 111, 930

Bult, H. *see* De Meyer, G.R.Y., 112, 471

Bültmann, R., Kurz, A.K. and Starke, K. α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151

Bültmann, R., Trendelenburg, M. and Starke, K. Blockade of P_{2X} purinoceptors by trypan blue in rat vas deferens, 113, 349

Bültmann, R. and Starke, K. Blockade by 4,4'-diisothiocyanato-stilbene-2,2'-disulphonate (DIDS) of P_{2X} -purinoceptors in rat vas deferens, 112, 690

Bunce, K.T. *see* Gale, J.D., 111, 332

Bunce, K.T. *see* Tam, F.S.-F., 113, 143

Burns, F., Stevens, P.A. and Payne, N.J. The identification of apparently novel cyclic AMP and cyclic GMP phosphodiesterase activities in guinea-pig tracheal smooth muscle, 113, 3

Burnstock, G. *see* Bo, X., 112, 1151

Burnstock, G. *see* Kasakov, L., 112, 403

Burnstock, G. *see* Maynard, K.I., 112, 123

Burnstock, G. *see* Rubino, A., 112, 515

Burnstock, G. *see* Soediono, P., 113, 681

Burnstock, G. *see* Windscheif, U., 113, 1015

Burnstock, G. *see* Ziganshin, A.U., 111, 923

Burnstock, G. *see* Ziganshin, A.U., 113, 669

Burroughs, D.A. *see* Herd, C.M., 112, 292

Burton, J. *see* Lopes-Martins, R.A.B., 113, 81

Busse, R. *see* Bauersachs, J., 113, 1548

Busse, R. *see* Hecker, M., 111, 238

Busse, R. *see* Pörsti, I., 111, 652

Butler, A.R. *see* Salas, E., 112, 1071

Butt, S.K. *see* Field, J.L., 113, 607

Butterworth, A.R. *see* Rabbani, M., 111, 89

Buus, N.H., VanBavel, E. and Mulvany, M.J. Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579

Buxton, I.L.O. *see* Eckman, D.M., 111, 1053

C

Cage, P.A. *see* Humphries, R.G., 113, 1057

Cailmail, S. *see* Moreau, R., 112, 649

Calabresi, P. *see* Mercuri, N.B., 113, 831

Calapai, G., Mazzaglia, G., Cilia, M., Zingarelli, B., Squadrito, F. and Caputi, A.P. Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328

Calapai, G. *see* Squadrito, F., 113, 912

Calignano, A. *see* Brignola, G., 113, 1372

Calixto, J.B. *see* Schlemper, V., 111, 83

Calvo, S. *see* Kuijpers, G.A.J., 113, 471

Cambarrat, C. *see* Thollon, C., 112, 37

Campbell, T.J. *see* Martin, D.K., 111, 873

Campo, G.M. *see* Squadrito, F., 113, 912

Campo, G.M. *see* Squadrito, F., 113, 1153

Camus, J. *see* Waelbroeck, M., 112, 505

Canale, P. *see* Squadrito, F., 113, 912

Canale, P. *see* Squadrito, F., 113, 1153

Candura, S.M. *see* Tonini, M., 113, 1

Canet, E. *see* Féletalou, M., 112, 683

Cannell, M.B. *see* Grantham, C.J., 111, 483

Cano, J. *see* Santiago, M., 111, 837

Cantí, C., Martí, E., Marsal, J. and Solsona, C. Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Capasso, A. *see* Pieretti, S., 113, 1416

Capasso, F., Mascolo, N., Izzo, A.A. and Gaginella, T.S. Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of *N*^G-nitro-L-arginine methyl ester, 113, 1127

Capasso, F. *see* Izzo, A.A., 113, 228

Capdeville-Atkinson, C. *see* Atkinson, J., 111, 1184

Capdeville-Atkinson, C. *see* Thorin, E., 113, 363

Čapek, R. *see* Clarke, P.B.S., 111, 397

Caputi, A.P. *see* Calapai, G., 111, 1328

Caputi, A.P. *see* Squadrito, F., 113, 912

Caputi, A.P. *see* Squadrito, F., 113, 1153

Carayon, A. *see* Su, J., 113, 395

Carceller, A. *see* Ballarín, M., 113, 425

Cardell, L.O. and Edvinsson, L. Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H_3 receptors in the trachea, 111, 445

Carnegi, G. *see* Bova, S., 111, 1067

Cartmell, J., Kemp, J.A., Alexander, S.P.H., Shinozaki, H. and Kendall, D.A. Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364

Carvalho, A.P. *see* Malva, J.O., 113, 1439

Carvalho, C.M. *see* Malva, J.O., 113, 1439

Castoldi, A.F. *see* Tonini, M., 113, 1

Castro-Faria-Neto, H.C. *see* Pires, A.L.A., 113, 994

Cavari, S., Fiorelli, G. and Vannucchi, S. Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254

Cavero, I. *see* Floch, A., 111, 759

Cazaubon, C. *see* Roccon, A., 111, 145

Cecioni, I. *see* Modesti, P.A., 112, 81

Cenarruzabeitia, E. *see* Ramírez, M.J., 111, 419

Cenarruzabeitia, E. *see* Ramírez, M.J., 113, 77

Cereda, E. *see* Sagrada, A., 113, 711

Ceserani, R. *see* Gallico, L., 112, 795

Challiss, R.A.J., Mistry, R., Gray, D.W. and Nahorski, S.R. Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231

Chamelian, M., Lécrivain, A., Robichaud, A. and du Souich, P. Effect of dehydration and hyperosmolal hydration on lignocaine and metabolites disposition in conscious rabbits, 113, 317

Chamienia, A.L. and Johns, E.J. The renal functional responses to 5-HT_{1A} receptor agonist, flesinoxan, in anaesthetized, normotensive rat, 112, 214

Chan, K.-m. *see* Qian, Y.-m., 113, 369

Chan, S.L.F., Brown, C.A., Scarpello, K.E. and Morgan, N.G. The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065

Chang, C.C. *see* Chiou, L.C., 112, 257

Chang, D.H.-T. *see* Einstein, R., 111, 199

Chang, G.-J. *see* Lin, C.-H., 113, 275

Chao, A.C., Zifferblatt, J.B., Wagner, J.A., Dong, Y.-J., Gruenert, D.C. and Gardner, P. Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169

Chapelain, B. *see* Galizzi, J.P., 113, 389

Chapman, R.W. *see* Bolser, D.C., 113, 1344

Charoonroje, P., Tokumitsu, Y. and Nomura, Y. Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569

Chaudieu, I. *see* Clarke, P.B.S., 111, 397

Chauveau, M. *see* Larrazet, F., 113, 649

Checler, F. *see* Barelli, H., 112, 127

Chemtob, S. *see* Li, D.Y., 112, 59

Chen, C.-C. *see* Young, M.-L., 113, 69

Chen, C.C. *see* Wang, J.P., 113, 945

Chen, H., Jassar, B.S., Kurenni, D.E. and Smith, P.A. Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55

Chen, J.R. *see* Wiley, J.S., 112, 946

Chen, S., Inoue, R., Inomata, H. and Ito, Y. Role of cyclic AMP-induced C1 conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137

Chern, J.-W. *see* Lee, Y.-M., 112, 1031

Chess-Williams, R. *see* Otter, D.J., 113, 576

Chi, L. *see* Friedrichs, G.S., 113, 209

Chiba, K., Moriyama, S., Ishigai, Y., Fukuzawa, A., Irie, K. and Shibano, T. Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837

Chiba, S. *see* Ren, L.-M., 111, 1321

Chieng, B. and Christie, M.J. Hyperpolarization by opioids acting on μ -receptors of a sub-population of rat periaqueductal gray neurones *in vitro*, 113, 121

Chieng, B. and Christie, M.J. Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Chiesi, M. *see* Yard, N.J., 113, 1001

Chin, J.P.F., Kaye, D.M., Hurlston, R.M., Angus, J.A., Jennings, G.L. and Dart, A.M. Effects of dietary marine oil supplementation on

reactivity of human buttock subcutaneous arteries and forearm veins *in vitro*, 112, 566

Chiou, L.C. and Chang, C.C. Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Chiyotani, A., Tamaoki, J., Takeuchi, S., Kondo, M., Isono, K. and Konno, K. Stimulation by menthol of C1 secretion via a Ca^{2+} -dependent mechanism in canine airway epithelium, 112, 571

Chizh, B.A., Cumberbatch, M.J. and Headley, P.M. A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843

Chong, N.W.S. and Sugden, D. Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Chopra, L.C., Twort, C.H.C. and Ward, J.P.T. Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237

Chou, C.C. *see* Alemayehu, A., 111, 205

Chou, S.-P. *see* Yang, C.M., 111, 21

Chrapusta, S.J. *see* Karoum, F., 113, 1391

Christie, M.J. *see* Chieng, B., 113, 121

Christie, M.J. *see* Chieng, B., 113, 303

Christodoulou, C. *see* Buchan, K.W., 112, 1251

Christofoli, F.L. and Wood, J.D. Electrophysiological subtypes of inhibitory P_1 purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703

Chudziak, F. *see* Schwanstecher, M., 113, 903

Chuliá, S., Ivorra, M.D., Lugnier, C., Vila, E., Noguera, M.A. and D'Ocon, P. Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377

Chung, K.F. *see* Nichol, G.M., 111, 918

Chung, S.C., McDonald, T.V. and Gardner, P. Inhibition by SK&F 96365 of Ca^{2+} current, IL-2 production and activation in T lymphocytes, 113, 861

Church, J., Fletcher, E.J., Baxter, K. and MacDonald, J.F. Blockade by ifenprodil of high voltage-activated Ca^{2+} channels in rat and mouse cultured hippocampal pyramidal neurones: comparison with N-methyl-D-aspartate receptor antagonist actions, 113, 499

Church, M.K. *see* Walsh, D.A., 113, 1407

Ciabattoni, G. *see* Currò, D., 113, 541

Cilia, M. *see* Calapai, G., 111, 1328

Cipolla-Neto, J. *see* Ferreira, Z.S., 112, 107

Claing, A., Bkaily, G., Berthiaume, N., Sirois, P., Rola-Pleszczynski, M. and D'Orléans-Juste, P. Role of R -calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202

Claing, A. *see* D'Orléans-Juste, P., 113, 1257

Clapp, L.H. *see* Evans, A.M., 111, 972

Clark, K.L. *see* Buchan, K.W., 112, 1251

Clarke, D.E. *see* Waikar, M.V., 111, 213

Clarke, P.B.S., Chaudieu, I., El-Bizri, H., Boksa, P., Quik, M., Esplin, B.A. and Čapek, R. The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397

Clarke, P.B.S., Reuben, M. and El-Bizri, H. Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695

Clarke, P.B.S. *see* El-Bizri, H., 111, 406

Clarke, P.B.S. *see* El-Bizri, H., 111, 414

Clarke, P.B.S. *see* El-Bizri, H., 113, 917

Clayton, R.A. *see* Nally, J.E., 113, 1328

Clozel, M. *see* Filep, J.G., 113, 845

Coates, W.J. *see* Turner, N.C., 111, 1198

Coatney, R.W. *see* Alemayehu, A., 111, 205

Coccini, T. *see* Tonini, M., 113, 1

Cocks, T.M. *see* Kilpatrick, E.V., 112, 557

Cocks, T.M. *see* Stork, A.P., 113, 1093

Cocks, T.M. *see* Stork, A.P., 113, 1099

Coderre, T.J. *see* Fundytus, M.E., 113, 1215

Coëffier, E., Joseph, D. and Vargaftig, B.B. Role of interleukin-5 in enhanced migration of eosinophils from airways of immunized guinea-pigs, 113, 749

Cohen, R.B. *see* Dickinson, K.E.J., 113, 179

Colado, M.I. and Green, A.R. A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chloroformiazole, dizocilpine and other protective compounds, 111, 131

Cole, P.J. *see* Kanthakumar, K., 112, 493

Colella, A. *see* Modesti, P.A., 112, 81

Coleman, H.A. *see* Tonta, M.A., 112, 525

Colledge, W.H. *see* Cuthbert, A.W., 112, 31

Colling, E. *see* Molderings, G.J., 111, 733

Collins, P. *see* Jiang, C., 111, 711

Conigrave, A.D. *see* Gibb, C.A., 111, 1135

Connolly, G.P. Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357

Connor, H.E. *see* Beattie, D.T., 112, 262

Conti, A. *see* Dionisotti, S., 112, 659

Cook, D.I. *see* Gibb, C.A., 111, 1135

Cools, A.R. *see* Prinssen, E.P.M., 112, 769

Coppes, R.P., Brouwer, F., Freie, I., Smit, J. and Zaagsma, J. Sustained prejunctional facilitation of noradrenergic neurotransmission by adrenaline as a co-transmitter in the portal vein of freely moving rats, 113, 342

Cordeiro, R.S.B. *see* Pires, A.L.A., 113, 994

Corder, R. *see* Murphy, L.J., 113, 137

Corder, R. *see* Wellings, R.P., 111, 515

Corman, B. *see* Thorin, E., 113, 363

Corradetti, R. *see* Passani, M.B., 112, 695

Correia-de-Sá, P. and Ribeiro, J.A. Potentiation by tonic A_{2A} -adenosine receptor activation of CGRP-facilitated [^3H]-ACh release from rat motor nerve endings, 111, 582

Cortijo, J., Sanz, C.M., Villagrasa, V., Morcillo, E.J. and Small, R.C. The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769

Costentin, J. *see* Dubuc, I., 112, 352

Costoli, A. *see* Modesti, P.A., 112, 81

Coussy, L. *see* Galizzi, J.P., 113, 389

Couture, R. *see* Picard, P., 112, 240

Couture, R. *see* Prat, A., 112, 250

Cox, B.F., Greenland, B.D., Perrone, M.H. and Merkel, L.A. Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A_2 - but not to an A_1 -agonist in the dog, 111, 1233

Cox, H.M. and Tough, I.R. Calcitonin gene-related peptide receptors in human gastrointestinal epithelia, 113, 1243

Crack, B.E., Beukers, M.W., McKechnie, K.C.W., IJzerman, A.P. and Leff, P. Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2X} -purinoceptor ligands, 113, 1432

Crespo, M.S. *see* Pellón, M.I., 112, 1101

Cristol, J.-P. *see* Wellings, R.P., 111, 515

Cross, A.J. *see* Baldwin, H.A., 112, 188

Cross, A.J. *see* Watson, W.P., 112, 1017

Croxtall, J. *see* Mitchell, J.A., 113, 1008

Crozier, B. *see* Su, J., 113, 395

Cruwys, S.C., Garrett, N.E., Perkins, M.N., Blake, D.R. and Kidd, B.L. The role of bradykinin B_1 receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940

Cuevas, J. and Adams, D.J. Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663

Culliford, S.J. *see* Ellory, J.C., 111, 903

Cumberbatch, M.J. *see* Chizh, B.A., 112, 843

Cundell, D.R. *see* Kanthakumar, K., 112, 493

Cunha, F.Q. *see* Braga da Motta, J.I., 112, 111

Cunningham, J. *see* Neal, M., 113, 1085

Cunningham, J.R. *see* Neal, M.J., 113, 789

Cunningham, S.M.C., Mihara, S. and Lees, G.M. Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Currò, D., Preziosi, P., Ragazzoni, E. and Ciabattoni, G. Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541

Curtis, A.R. *see* Alexander, S.P.H., 113, 1501

Cuthbert, A.W., Hickman, M.E., MacVinish, L.J., Evans, M.J., Colledge, W.H., Ratcliff, R., Seale, P.W. and Humphrey, P.P.A. Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31

D

da Silva, W.D. *see* Tambourgi, D.V., 112, 723
 Dale, M.M. *see* Bahl, A.K., 113, 809
 Dalla Rosa, C. *see* Gallico, L., 112, 795
 Dallaire, N. *see* Marleau, S., 112, 654
 Daly, S.A. and Waddington, J.L. The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839
 Daniel, E.E. *see* Barelli, H., 112, 127
 Daniel, E.E. *see* Low, A.M., 112, 604
 Darlington, C.L. *see* Gilchrist, D.P.D., 111, 358
 Dart, A.M. *see* Chin, J.P.F., 112, 566
 Dashwood, M.R., Allen, S.P., Luu, T.N. and Muddle, J.R. The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
 Davenport, A.P., Kuc, R.E., Fitzgerald, F., Maguire, J.J., Berryman, K. and Doherty, A.M. [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
 Davenport, A.P., Kuc, R.E., Hoskins, S.L., Karet, F.E. and Fitzgerald, F. [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
 Davenport, A.P. *see* Gardiner, S.M., 112, 477
 Davenport, A.P. *see* Maguire J.J., 113, 49
 Davies, D.S. *see* Shaunak, S., 113, 151
 Davies, S.N. *see* Shuker, M.A., 111, 951
 Davis, A.J. and Perkins, M.N. The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
 Davis, E.A. *see* James, G.M., 111, 370
 Davis, G. *see* Huang, C., 111, 111
 Dawid-Milner, M.S. *see* St Lambert, J.H., 113, 159
 Dawson, D.A., Graham, D.I., McCulloch, J. and Macrae, I.M. Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a N-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247
 de Brum Fernandes, A.J. *see* Sirois, M.G., 111, 1111
 de Fátima D. Furtado, M. *see* Tambourgi, D.V., 112, 723
 de Freitas, M.C.W. *see* Tambourgi, D.V., 112, 723
 de Lima, W.T. *see* Sirois, M.G., 111, 1111
 de Lorenzi, F.G., Bridal, T.R. and Spinelli, W. Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
 De Man, J.G., Boeckxstaens, G.E., Herman, A.G. and Pelckmans, P.A. Effect of potassium channel blockade and α₂-adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341
 De Meyer, G.R.Y., Bult, H., Üstünes, L., Kockx, M., Jordaeans, F.H., Zonnekeyn, L.L. and Herman, A.G. Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471
 de Montigny, C. *see* Monnet, F.P., 112, 709
 de Nucci, G. *see* Grassi-Kassis, D.M., 113, 675
 de Nucci, G. *see* Lopes-Martins, R.A.B., 113, 81
 de Wildt, D.J. *see* Kappelle, A.C., 111, 887
 Debotto, P. *see* Bova, S., 111, 1067
 Debonnel, G. *see* Monnet, F.P., 112, 709
 Deckert, V., Pruneau, D. and Elghozi, J.-L. Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939
 DeFeo, P.A. *see* Keith, R.A., 113, 379
 DeGennaro, F.C. *see* Bolser, D.C., 113, 1344
 Dejana, E. *see* Tranchina, S., 111, 575
 Del Maschio, A. *see* Tranchina, S., 111, 575
 del Pozo, B.F. *see* Pérez-Vizcaíno, F., 113, 1281
 Del Río, J. *see* Ramírez, M.J., 111, 419
 Del Río, J. *see* Ramírez, M.J., 113, 77
 Del Río, J. *see* Zazpe, A., 113, 968
 Del Rio-Lacheze, C. *see* Gué, M., 111, 930
 Delamere, F., Holland, E., Patel, S., Bennett, J., Pavord, I. and Knox, A. Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983
 Delaney, C.L. *see* Dickinson, K.E.J., 113, 179
 Delaunois, A., Gustin, P., Dassy-Doize, C. and Ansay, M. Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
 DelliPizzi, A., Hilchey, S.D. and Bell-Quiley, C.P. Natriuretic action of angiotensin(1-7), 111, 1
 Delpón, E. *see* Pérez, O., 111, 717
 Delumeau, J.C. *see* Hubert, J.P., 113, 261
 den Hertog, A. *see* Henning, R.H., 111, 459
 Denduchis, B. *see* Bacman, S., 111, 271
 Derrien, M., Durieux, C. and Roques, B.P. Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltrindole, 111, 956
 Desai, K.M., Warner, T.D., Bishop, A.E., Polak, J.M. and Vane, J.R. Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197
 Desai, K.M., Warner, T.D. and Vane, J.R. 5-HT₃ receptors do not mediate vagally-induced relaxation or contraction of the isolated stomach of the guinea-pig, 111, 346
 DeSiato, M.A. *see* Farmer, S.G., 112, 461
 Dassy-Doize, C. *see* Delaunois, A., 113, 973
 Dev, K.K. *see* Barnes, J.M., 113, 339
 Di Giannuario, A. *see* Pieretti, S., 113, 1416
 Di Rosa, M. *see* Brignola, G., 113, 1372
 Dickel, C. *see* Schwanstecher, C., 111, 302
 Dickel, C. *see* Schwanstecher, M., 113, 903
 Dickinson, K.E.J., Cohen, R.B., Skwisch, S., Delaney, C.L., Serafino, R.P., Poss, M.A., Gu, Z., Ryono, D.E., Moreland, S. and Powell, J.R. BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179
 Diéguez, G. *see* Fernández, N., 113, 550
 Dilly, K.W. *see* Donoso, P., 111, 455
 Dionisotti, S., Conti, A., Sandoli, D., Zocchi, C., Gatta, F. and Ongini, E. Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
 Dive, V. *see* Barelli, H., 112, 127
 Doble, A. *see* Hubert, J.P., 113, 261
 D'Ocon, P. *see* Chuliá, S., 113, 1377
 Doherty, A.M. *see* Davenport, A.P., 111, 4
 Dohi, Y., Kojima, M. and Sato, K. Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673
 Doi, K. *see* Kaneda, M., 111, 1355
 Dolan, J.S. *see* Turner, N.C., 111, 1198
 Dolly, J.O. *see* Hall, A., 113, 959
 Domenici, M.R. *see* Pieretti, S., 113, 1416
 Donat, F. *see* Roccon, A., 111, 145
 Donello, J.E. *see* Regan, J.W., 112, 377
 Dong, Y.-J. *see* Chao, A.C., 112, 169
 Donigi-Gale, D. *see* Herd, C.M., 111, 7
 Donigi-Gale, D. *see* Herd, C.M., 112, 292
 Donoso, M.V., Salas, C., Sepúlveda, G., Lewin, J., Fournier, A. and Huidobro-Toro, J.P. Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473
 Donoso, P., O'Neill, S.C., Dilly, K.W., Negretti, N. and Eisner, D.A. Comparison of the effects of caffeine and other methylxanthines on [Ca²⁺]_i in rat ventricular myocytes, 111, 455
 Doolan, C.M. and Keenan, A.K. Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509
 D'Orléans-Juste, P., Clain, A., Télémque, S., Maurice, M.-C., Yano, M. and Gratton, J.-P. Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257
 D'Orléans-Juste, P. *see* Clain, A., 112, 1202
 dos Santos, M.C. *see* Tambourgi, D.V., 112, 723
 Douglas, G.J. *see* Paul, W., 111, 159
 Drew, G.M. *see* Balmforth, A.J., 112, 277
 Drew, G.M. *see* Hilditch, A., 111, 137
 Drieman, J.C., van Kan, F.J.P.M., Thijssen, H.H.W., van Essen, H., Smits, J.F.M. and Struijker Boudier, H.A.J. Regional haemodynamic effects of dopamine and its prodrugs L-dopa and gludopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
 Drieu la Rochelle, C., Grosset, A. and O'Connor, S.E. Comparison of

the haemodynamic profiles of elgodipine and nicardipine in the anaesthetized dog, 111, 49

Driscoll, E.M. *see* Lucchesi, B.R., 113, 1333

Droogmans, G. *see* Nilius, B., 112, 1049

Drumare, M.F. *see* Blin, N., 112, 911

Du Plooy, W.J., Hay, L., Kahler, C.P., Schutte, P.J. and Brandt, H.D. The dose-related hyper-and-hypokalaemic effects of salbutamol and its arrhythmogenic potential, 111, 73

du Souich, P. *see* Chamelian, M., 113, 317

Dubuc, I., Costentin, J., Terranova, J.P., Barnouin, M.C., Soubrié, P., Le Fur, G., Rostène, W. and Kitabgi, P. The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352

Duggan, A.W. *see* Lang, C.W., 112, 951

Dumont, P. *see* Trifilieff, A., 111, 1228

Durcan, M.J., Morgan, P.F., Van Etten, M.L. and Linnoila, M. Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

Durieux, C. *see* Derrien, M., 111, 956

Dusting, G.J. *see* Loke, K.E., 112, 630

Dykes, C.W. *see* Buchan, K.W., 112, 1251

E

e Silva, P.M.R. *see* Pires, A.L.A., 113, 994

Ebihara, S. *see* Uneyama, H., 111, 245

Eckly, A.E. and Lugnier, C. Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445

Eckman, D.M., Weinert, J.S., Buxton, I.L.O. and Keef, K.D. Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

Edvinsson, L. *see* Cardell, L.O., 111, 445

Edvinsson, L. *see* Ekelund, U., 112, 887

Edvinsson, L. *see* Gardiner, S.M., 112, 477

Edwards, G., Niederste-Hollenberg, A., Schneider, J., Noack, Th. and Weston, A.H. Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538

Edwards, J.S. and Ritter, J.M. Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926

Edwards, M.D. *see* Lees, G., 113, 642

Edwards, R.J. *see* Shaunak, S., 113, 151

Edwardson, J.M. *see* Koenig, J.A., 111, 1023

Egan, R.W. *see* Bolser, D.C., 113, 1344

Egeren, A.D.-V. *see* Manicourt, D.H., 113, 1113

Eglen, R.M. *see* Watson, N., 112, 179

Eikenburg, D.C. *see* Lakhani, P.P., 111, 1296

Einstein, R., Abdul-Hussein, N., Wong, T.-W., Chang, D.H.-T., Matthews, R. and Richardson, D.P. Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199

Eisner, D.A. *see* Donoso, P., 111, 455

Ekelund, U., Adner, M., Edvinsson, L. and Mellander, S. Effects of selective ET_B-receptor stimulation on arterial, venous and capillary functions in cat skeletal muscle, 112, 887

El-Bizri, H. and Clarke, P.B.S. Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

El-Bizri, H. and Clarke, P.B.S. Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

El-Bizri, H. and Clarke, P.B.S. Regulation of nicotinic receptors in rat brain following quasi-irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917

El-Bizri, H. *see* Clarke, P.B.S., 111, 397

El-Bizri, H. *see* Clarke, P.B.S., 111, 695

Elghozi, J.-L. *see* Deckert, V., 112, 939

Elhawary, A.M. and Pang, C.C.Y. α_1 -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819

Ellenbroek, B.A. *see* Prinsen, E.P.M., 112, 769

Elliott, J. *see* Rupniak, N.M.J., 113, 1487

Ellis, E.S. *see* Wardle, K.A., 112, 789

Ellory, J.C., Culliford, S.J., Smith, P.A., Wolowyk, M.W. and Knaus, E.E. Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903

Emslander, H.P. *see* Hauck, R.W., 113, 1043

Endo, M. *see* Kasai, Y., 112, 1132

Enokibori, M., Okamura, T. and Toda, N. Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77

Entzeroth, M. *see* Schambye, H.T., 113, 331

Erdemli, G. and Krnjević, K. Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

Erdmann, E. *see* Böhm, M., 112, 137

Ernst, G.E. *see* Keith, R.A., 113, 379

Escriba, P.V. *see* Olmos, G., 111, 997

Esplin, B.A. *see* Clarke, P.B.S., 111, 397

Evans, A.M., Clapp, L.H. and Gurney, A.M. Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972

Evans, M.J. *see* Cuthbert, A.W., 112, 31

Evans, R.H. *see* Siarey, R.J., 111, 497

Evans, R.J. and Kennedy, C. Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853

Evenden, J.L. The effect of 5-HT_{1A} receptor agonists on locomotor activity in the guinea-pig, 112, 861

F

Faccioli, L.H. *see* Weg, V.B., 113, 35

Facklam, M. and Bowery, N.G. Erratum, 111, 648

Fairbairn, C.E. *see* Regan, J.W., 112, 377

Fairbairn, S.M. *see* Teixeira, M.M., 113, 1363

Fallgren, B. *see* Gardiner, S.M., 113, 129

Fan, P. Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741

Fan, P. Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745

Fardin, V. *see* Floch, A., 111, 759

Farmer, S.G. and DeSiato, M.A. Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461

Farthing, M.J.G. *see* Battistini, B., 112, 1244

Fauchère, J.-L. *see* Féltetou, M., 112, 683

Featherstone, R. *see* Walsh, D.A., 113, 1407

Feifel, R. *see* Waelbroeck, M., 112, 505

Feldman, J. *see* Monassier, L., 111, 1347

Feldtrauer, J.-J. *see* Waldmeier, P.C., 113, 1515

Féltetou, M., Germain, M., Thurieau, C., Fauchère, J.-L. and Canet, E. Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683

Felici, L. *see* Lakhani, P.P., 111, 1296

Feniuk, W. *see* McKeen, E.S., 113, 628

Feolde, E. *see* Vigne, P., 112, 775

Ferdinand, P. *see* Szilvassy, Z., 112, 999

Ferlito, M. *see* Squadrito, F., 113, 1153

Fernandes, M.H. *see* Soares-da-Silva, P., 112, 611

Fernández, N., Monge, L., García-Villalón, A.L., García, J.L., Gómez, B. and Diéguez, G. Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550

Ferrando, R. *see* Alegret, M., 112, 551

Ferreira, A.T. *see* Silva, E.G., 113, 1022

Ferreira, S.H. *see* Braga da Motta, J.I., 112, 111

Ferreira, T. *see* Baldwin, H.A., 112, 188

Ferreira, Z.S., Cipolla-Neto, J. and Markus, R.P. Presence of P₂-purinoceptors in the rat pineal gland, 112, 107

Ferry, C.B. *see* Kelly, S.S., 111, 747

Fickling, S.A. *see* MacAllister, R.J., 112, 43

Field, J.L., Butt, S.K., Morton, I.K.M. and Hall, J.M. Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607

Figueira, M.A. *see* Hernandez, J., 111, 1316

Filep, J.G., Clozel, M., Fournier, A. and Földes-Filep, É. Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

Filep, J.G., Fournier, A. and Földes-Filep, É. Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A

receptor, platelet-activating factor and thromboxane A₂, 112, 963
 Fiorelli, G. *see* Cavari, S., 113, 254
 Fischer, B. *see* Bo, X., 112, 1151
 Fitzgerald, F. *see* Davenport, A.P., 111, 4
 Fitzgerald, F. *see* Davenport, A.P., 113, 1303
 Fletcher, E.J. *see* Church, J., 113, 499
 Fletcher, S.R. *see* Rupniak, N.M.J., 113, 1487
 Floch, A., Fardin, V. and Cavero, I. Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759
 Flor, B. *see* Martínez, M.C., 113, 419
 Flower, R.J. *see* Perretti, M., 112, 801
 Földes-Filep, É. *see* Filep, J.G., 112, 963
 Földes-Filep, É. *see* Filep, J.G., 113, 845
 Fontela, T. *see* Hermida, O.G., 111, 861
 Forbes, I. *see* Kennett, G.A., 111, 797
 Forbes, I.J. *see* MacDonald, A., 112, 576
 Ford, A.P.D.W. *see* Waikar, M.V., 111, 213
 Foreman, J.C. *see* Austin, C.E., 111, 969
 Foreman, J.C. *see* Bahl, A.K., 113, 809
 Foreman, J.C. *see* Friis, U.G., 111, 1179
 Forsyth, M.J. *see* Matthews, E.K., 113, 87
 Fournier, A. *see* Battistini, B., 111, 1009
 Fournier, A. *see* Battistini, B., 112, 1244
 Fournier, A. *see* Donoso, M.V., 111, 473
 Fournier, A. *see* Filep, J.G., 112, 963, 113, 845
 Fox, A.J. *see* Ward, J.K., 111, 1095
 Fox-Threlkeld, J.E.T. *see* Barelli, H., 112, 127
 Franceschetti, G.P. *see* Tonini, M., 113, 1
 Franks, N.P. *see* Hall, A.C., 112, 906
 Frediani-Neto, E. *see* Silva, E.G., 113, 1022
 Freedman, S.B. *see* Rupniak, N.M.J., 113, 1487
 Freedman, S.B. *see* Seabrook, G.R., 111, 391
 Freedman, S.B. *see* Seabrook, G.R., 111, 1061
 Freedman, S.B. *see* Seabrook, G.R., 112, 728
 Freie, I. *see* Coppes, R.P., 113, 342
 Frelin, C. *see* Vigne, P., 112, 775
 Friedrichs, G.S., Chi, L., Green, A.L. and Lucchesi, B.R. Anti-fibrillatory effects of clofilium in the rabbit isolated heart, 113, 209
 Friend, S.J. *see* Kenny, B.A., 111, 1003
 Friis, U.G., Johansen, T., Hayes, N.A. and Foreman, J.C. IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179
 Frosini, M. *see* Palmi, M., 112, 449
 Frossard, N. *see* Larrazet, F., 113, 649
 Fryer, A.D., Yarkony, K.A. and Jacoby, D.B. The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in para-influenza virus-infected guinea-pigs, 112, 588
 Fu, W.-M. and Huang, F.-L. Potentiation by endogenously released ATP of spontaneous transmitter secretion at developing neuromuscular synapses in *Xenopus* cell cultures, 111, 880
 Fukuda, T. *see* Yasumatsu, H., 111, 1170
 Fukui, K. *see* Kaneda, M., 111, 1355
 Fukuroda, T., Ozaki, S., Ihara, M., Ishikawa, K., Yano, M. and Nishikibe M. Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336
 Fukushima, H. *see* Kashiwabara, T., 111, 278
 Fukuzawa, A. *see* Chiba, K., 112, 837
 Fulton, D., McGiff, J.C. and Quilley, J. Role of K⁺ channels in the vasodilator response to bradykinin in the rat heart, 113, 954
 Funayama, K. *see* Furukawa, K.-I., 113, 233
 Fundytus, M.E. and Coderre, T.J. Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
 Fung, H.-L. *see* Tabrizi-Fard, M.A., 111, 394
 Furió, V. *see* Pellón, M.I., 112, 1101
 Furman, B.L. *see* Song, W., 113, 1083
 Furness, J.B. *see* Johnson, P.J., 111, 1240
 Furness, J.B. *see* Yuan, S.Y., 112, 1095
 Furukawa, K.-I., Funayama, K., Ohkura, M., Oshima, Y., Tu, A.T. and Ohizumi, Y. Ca²⁺ release induced by myotoxin *a*, a radio-labellable probe having novel Ca²⁺ release properties in sarcoplasmic reticulum, 113, 233
 Furukawa, T. *see* Katsuragi, T., 113, 795

Fyhrquist, F. *see* Mervaala, E.M.A., 111, 1189

G

Gadre, A. *see* Kennett, G.A., 111, 797
 Gaffen, Z. *see* Allawi, H.S., 113, 282
 Gaginella, T.S. *see* Capasso, F., 113, 1127
 Gaginella, T.S. *see* Izzo, A.A., 113, 228
 Gale, J.D., Grossman, C.J., Whitehead, J.W.F., Oxford, A.W., Bunce, K.T. and Humphrey, P.P.A. GR113808: a novel, selective antagonist with high affinity at the 5-HT₄ receptor, 111, 332
 Gajlione, A. *see* Varney, M.A., 112, 390
 Galizzi, J.P., Bodinier, M.C., Chapelain, B., Ly, S.M., Coussy, L., Giraud, S., Neliat, G. and Jean, T. Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389
 Gallacher, D. *see* MacDonald, A., 112, 576
 Gallico, L., Borghi, A., Dalla Rosa, C., Ceserani, R. and Tognella, S. Mogusine: a novel peripheral non-narcotic antitussive drug, 112, 795
 Galvan, M. *see* Leishman, D.J., 111, 318
 Galzin, A.M. *see* Briand, V., 112, 1195
 García, J.L. *see* Fernández, N., 113, 550
 García-Sevilla, J.A. *see* García-Vallejo, P., 111, 1140
 García-Sevilla, J.A. *see* Olmos, G., 111, 997
 García-Vallejo, P., Barturen, F. and García-Sevilla, J.A. Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
 García-Villalón, A.L. *see* Fernández, N., 113, 550
 Gardiner, S.M., Kemp, P.A., Fallgren, B. and Bennett, T. Effects of chronic infusions of α -trinisol on regional and cardiac haemodynamics in conscious rats, 113, 129
 Gardiner, S.M., Kemp, P.A., March, J.E., Bennett, T., Davenport, A.P. and Evinsson, L. Effects of an ET₁-receptor antagonist, FR139317, on regional haemodynamic responses to endothelin-1 and [Ala^{11,15}]Ac-endothelin-1 (6-21) in conscious rats, 112, 477
 Gardiner, S.M., Kemp, P.A., March, J.E. and Bennett, T. Effects of bosentan (Ro 47-0203), an ET_A, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
 Gardiner, S.M., Rakhit, T., Kemp, P.A., March, J.E. and Bennett, T. Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589
 Gardiner, S.M. *see* Kemp, P.A., 111, 325
 Gardiner, S.M. *see* Waller, J., 112, 1057
 Gardner, C.J. *see* Hilditch, A., 111, 137
 Gardner, P. *see* Chao, A.C., 112, 169
 Gardner, P. *see* Chung, S.C., 113, 861
 Garland, C.J. *see* Parsons, S.J.W., 113, 1275
 Garland, C.J. *see* Seager, J.M., 111, 525
 Garland, C.J. *see* Waldron, G.J., 112, 831
 Garrett, N.E. *see* Cruwys, S.C., 113, 940
 Gaspar, V. *see* Low, A.M., 113, 1263
 Gaster, L.M. *see* Wardle, K.A., 112, 789
 Gatta, F. *see* Dionisotti, S., 112, 659
 Gean, P.-W. *see* Huang, C.-C., 113, 1029
 Gebicke-Haerter, P.J. *see* Langosch, J.M., 113, 29
 Gebicke-Haerter, P.J. *see* Nörenberg, W., 111, 942
 Geppetti, P. *see* Ricciardolo, F.L.M., 113, 508
 Geppetti, P. *see* Ricciardolo, F.L.M., 113, 1147
 Germain, M. *see* Félix, M., 112, 683
 Ghaleh, B. *see* Béa, M.-L., 111, 981
 Ghelardini, C. *see* Malmberg-Aiello, P., 111, 1269
 Ghiglione, M. *see* Hermida, O.G., 111, 861
 Giachetti, A. *see* Astolfi, M., 111, 570
 Giachetti, A. *see* Maggi, C.A., 112, 150
 Gibb, C.A., Singh, S., Cook, D.I., Poronnik, P. and Conigrave, A.D. A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST₈₈₅, 111, 1135
 Gibson, A., McFadzean, I., Tucker, J.F. and Wayman, C. Variable potency of norepinephrine-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
 Gibson, W.J., Roques, T.W. and Young, J.M. Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-

hydroxymethyl-propan-1,3-diol HCl, 111, 1262
 Giembycz, M.A. *see* Souness, J.E., 111, 1081
 Gies, J.-P. *see* Trifilieff, A., 111, 1228
 Gil, D.W. *see* Regan, J.W., 112, 377
 Gilchrist, D.P.D., Darlington, C.L. and Smith, P.F. A dose-response analysis of the beneficial effects of the ACTH-(4-9) analogue, Org 2766, on behavioural recovery following unilateral labyrinthectomy in guinea-pig, 111, 358
 Gillespie, J.S. *see* Liu, X., 111, 1287
 Gilroy, D.W. *see* Tomlinson, A., 113, 693
 Giotti, A. *see* Malmberg-Aiello, P., 111, 1269
 Girard, D. *see* Berry, C.N., 113, 1209
 Giraud, S. *see* Galizzi, J.P., 113, 389
 Gispen, W.H. *see* Kappelle, A.C., 111, 887
 Giudicelli, J.-F. *see* Béa, M.-L., 111, 981
 Giuliani, S. *see* Ahluwalia, A., 111, 1017
 Giuliani, S. *see* Maggi, C.A., 111, 687
 Giuliani, S. *see* Maggi, C.A., 112, 150
 Gleizes-Escalà, C. *see* Gué, M., 111, 930
 Glen, A. *see* Kennett, G.A., 111, 797
 Glowinski, J. *see* Hubert, J.P., 113, 261
 Gniadecka, M. *see* Gniadecki, R., 113, 439
 Gniadecki, R., Gniadecka, M. and Serup, J. Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D₃, 113, 439
 Goldie, R.G., Grayson, P.S., Knott, P.G., Self, G.J. and Henry, P.J. Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749
 Goldie, R.G. *see* Henry, P.J., 112, 1188
 Goldie, R.G. *see* Spina, D., 111, 29
 Goldie, R.G. *see* Spina, D., 112, 895
 Goldie, R.G. *see* Spina, D., 112, 901
 Golomb, E. *see* Abassi, Z.A., 113, 204
 Gómez, B. *see* Fernández, N., 113, 550
 Gooderham, N.J. *see* Shaunak, S., 113, 151
 Goodstadt, L.J. *see* Kozlowski, R.Z., 111, 250
 Göthert, M. *see* Molderings, G.J., 111, 733
 Goto, K. *see* Yasuda, N., 113, 21
 Gottschlich, R. *see* Barber, A., 111, 843
 Gottschlich, R. *see* Barber, A., 113, 1317
 Graf, P.D. *see* Ricciardolo, F.L.M., 113, 508
 Graham, D.I. *see* Dawson, D.A., 113, 247
 Grantham, C.J., Main, M.J. and Cannell, M.B. Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells, 111, 483
 Grassi-Kassisse, D.M., Antunes, E., Withrington, P.G. and de Nucci, G. Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
 Gratton, J.-P. *see* D'Orléans-Juste, P., 113, 1257
 Gray, D.W. *see* Challiss, R.A.J., 112, 231
 Grayson, P.S. *see* Goldie, R.G., 112, 749
 Greber, S., Schwarzer, C. and Sperk, G. Neuropeptide Y inhibits potassium-stimulated glutamate release through Y₂ receptors in rat hippocampal slices *in vitro*, 113, 737
 Green, A.L. *see* Friedrichs, G.S., 113, 209
 Green, A.R. *see* Baldwin, H.A., 112, 188
 Green, A.R. *see* Colado, M.I., 111, 131
 Green, A.R. *see* Watson, W.P., 112, 1017
 Greengrass, P.M. *see* Kenny, B.A., 111, 1003
 Greenland, B.D. *see* Cox, B.F., 111, 1233
 Greger, R. *see* Pavenstädt, H., 113, 7
 Grégoire, G. *see* Pacaud, P., 113, 457
 Greiner, H.E. *see* Barber, A., 111, 843
 Greiner, H.E. *see* Barber, A., 113, 1317
 Grewal, S. *see* Kennett, G.A., 111, 797
 Griesbacher, T., Sutliff, R.L. and Lembeck, F. Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004
 Griesbacher, T. Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547
 Griesbacher, T. *see* Legat, F.J., 112, 453
 Griffin, G. *see* Pertwee, R.G., 111, 968
 Griffith, T.M. *see* Hutcheson, I.R., 113, 190
 Griffith, T.M. *see* Randall, M.D., 112, 49
 Grimsditch, D. *see* Turner, N.C., 111, 1198
 Groß, G. *see* Michel, M.C., 111, 533, 539
 Gronier, B. *see* Monnet, F.P., 112, 709
 Groschner, K. *see* Kühberger, E., 113, 1289
 Grosset, A. *see* Drieu la Rochelle, C., 111, 49
 Grossman, C.J. *see* Gale, J.D., 111, 332
 Grous, M. and Barnette, M. Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259
 Grubb, B.D. *see* Lang, C.W., 112, 951
 Gruenert, D.C. *see* Chao, A.C., 112, 169
 Grundemar, L. *see* Kahl, U., 111, 1129
 Gryglewski, R.J. *see* Korbut, R., 112, 289
 Gu, Z. *see* Dickinson, K.E.J., 113, 179
 Gué, M., Gleizes-Escalà, C., Del Rio-Lacheze, C., Junien J.-L. and Buéno, L. Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930
 Guh, J.-H. *see* Ko, F.-N., 112, 1174
 Guild, S.B. *see* McMullan, B.W., 113, 171
 Guitart, X. *see* Ballarín, M., 113, 425
 Guo, J.-Z. *see* Hosoki, R., 113, 1141
 Gurney, A.M. *see* Evans, A.M., 111, 972
 Gustafsson, L.E. *see* Iversen, H.H., 113, 1088
 Gustafsson, L.E. *see* Persson, M.G., 111, 825
 Gustin, P. *see* Delaunois, A., 113, 973
 Gylfe, E. *see* Sjödin, L., 112, 1037

H

Haazen, L. *see* Manicourt, D.H., 113, 1113
 Håkanson, R. *see* Wang, Z.-Y., 111, 179
 Håkanson, R. *see* Wang, Z.-Y., 111, 1342
 Hales, C.N. *see* Lee, K., 111, 632
 Hall, A., Stow, J., Sorensen, R., Dolly, J.O. and Owen, D. Blockade by dendrotoxin homologues of voltage-dependent K⁺ currents in cultured sensory neurones from neonatal rats, 113, 959
 Hall, A.C., Lieb, W.R. and Franks, N.P. Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906
 Hall, I.P. *see* McCreathe, G., 113, 823
 Hall, J.M., Mitchell, D. and Morton, I.K.M. Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985
 Hall, J.M. and Brain, S.D. Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
 Hall, J.M. *see* Field, J.L., 113, 607
 Hama, H. *see* Yasuda, N., 113, 21
 Hamaguchi, A. *see* Kim, S., 113, 662
 Hamburger, M. *see* Hay, A.J.B., 112, 9
 Hanft, G. *see* Michel, M.C., 111, 533
 Hanft, G. *see* Michel, M.C., 111, 539
 Hänninen, V. *see* Santti, E., 113, 1231
 Harada, M. *see* Arimura, A., 111, 339
 Haralabopoulos, G. *see* Pipili-Synetos, E., 111, 894
 Harden, T.K. *see* Boyer, J.L., 113, 614
 Harris, J.G. *see* Perretti, M., 112, 801
 Harting, J. *see* Barber, A., 111, 843
 Harting, J. *see* Barber, A., 113, 1317
 Hashimoto, M. *see* Shinozuka, K., 113, 1203
 Hassérian, H., Vaca, L. and Kunze, D.L. Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118
 Hata, F. *see* Suthamnatpong, N., 112, 676
 Hatano, A., Takahashi, H., Tamaki, M., Komeyama, T., Koizumi, T. and Takeda, M. Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
 Hattori, K. *see* Shinozuka, K., 113, 1203
 Hauck, R.W., Schulz, Ch., Emslander, H.P. and Böhm, M. Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043
 Hay, A.J.B., Hamburger, M., Hostettmann, K. and Hoult, J.R.S. Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylene-butyrolactone functions, 112, 9

Hay, L. *see* Du Plooy, W.J., 111, 73
 Hayaishi, O. *see* Minami, T., 112, 735
 Hayes, N.A. *see* Friis, U.G., 111, 1179
 Haynes, J.M. *see* James, G.M., 111, 370
 Headley, P.M. *see* Chizh, B.A., 112, 843
 Hecker, M., Pörsti, I., Bara, A.T. and Busse, R. Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
 Hecker, M. *see* Bauersachs, J., 113, 1548
 Hecker, M. *see* Pörsti, I., 111, 652
 Heeps, G. *see* MacDonald, A., 112, 576
 Heijnen, C.J. *see* Benschop, R.J., 113, 1311
 Hellewell, P.G. *see* Teixeira, M.M., 111, 811
 Hellewell, P.G. *see* Teixeira, M.M., 111, 1205
 Hellewell, P.G. *see* Teixeira, M.M., 112, 332
 Hellewell, P.G. *see* Teixeira, M.M., 113, 1363
 Henley, J.M. *see* Barnes, J.M., 113, 339
 Henning, R.H., Nelemans, S.A., van den Akker, J. and den Hertog, A. Induction of Na^+/K^+ -ATPase activity by long-term stimulation of nicotinic acetylcholine receptors in C2C12 myotubes, 111, 459
 Henry, P.J. and Goldie, R.G. ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
 Henry, P.J. Inhibitory effects of nordihydroguaiaretic acid on ET_A -receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
 Henry, P.J. *see* Goldie, R.G., 112, 749
 Herbert, M.K. and Holzer, P. Interleukin-18 enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
 Herd, C.M., Donigi-Gale, D., Shoupe, T.S., Burroughs, D.A., Yeadon, M. and Page, C.P. Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292
 Herd, C.M., Donigi-Gale, D., Shoupe, T.S., Kilfeather, S.A., Okiji, S.A. and Page, C.P. Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7
 Herity, N.A., Allen, J.D., Silke, B. and Adgey, A.A.J. Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
 Herman, A.G. *see* De Man, J.G., 112, 341
 Herman, A.G. *see* De Meyer, G.R.Y., 112, 471
 Hermida, O.G., Fontela, T., Ghiglione, M. and Utenthal, L.O. Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
 Hernández, F., Alexander, S.P.H. and Kendall, D.A. Natriuretic peptide-induced cyclic GMP accumulation in adult guinea-pig cerebellar slices, 113, 216
 Hernandez, J., Pinto, F., Figueira, M.A. and Ribeiro, J.A. Evidence for a cooperation between adenosine A_2 receptors and β_1 -adrenoceptors on cardiac automaticity in the isolated right ventricle of the rat, 111, 1316
 Hey, J.A. *see* Bolser, D.C., 113, 1344
 Hickman, M.E. *see* Cuthbert, A.W., 112, 31
 Hilchey, S.D. *see* DelliPizzi, A., 111, 1
 Hilditch, A., Hunt, A.A.E., Gardner, C.J., Twissell, D.J., Polley, J., Travers, A., Drew, G.M., Middlemiss, D., Ross, B.C. and Robertson, M.J. Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137
 Hill, A. *see* Parsons, S.J.W., 113, 1275
 Hill, R.G. *see* Longmore, J., 112, 176
 Hill, R.G. *see* Rupniak, N.M.J., 113, 1487
 Hill, R.G. *see* Tan, K.K.C., 111, 703
 Hill, S.J. *see* Alexander, S.P.H., 111, 185
 Hill, S.J. *see* Alexander, S.P.H., 113, 1501
 Hill, S.J. *see* Iredale, P.A., 111, 1252
 Hill, S.J. *see* McCreathe, G., 113, 823
 Hill, S.J. *see* Marsh, K.A., 112, 934
 Hill, S.J. *see* Peakman, M.-C., 111, 191, 113, 1554
 Hillier, K. *see* Tam, F.S.-F., 113, 143
 Hirano, K. *see* Ushio-Fukai, M., 113, 1173
 Hirasawa, N. *see* Watanabe, M., 111, 123
 Hirasawa, Y. *see* Kita, Y., 113, 5
 Hirasawa, Y. *see* Kita, Y., 113, 385
 Hirasawa, Y. *see* Kita, Y., 113, 1137
 Hirayama, Y. *see* Ramnarine, S.I., 113, 1183
 Hirokawa, K., O'Shaughnessy, K., Moore, K., Ramrakha, P. and Wilkins, M.R. Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396
 Hirokawa, K., O'Shaughnessy, K.M., Ramrakha, P. and Wilkins, M.R. Inhibition of nitric oxide synthesis in vascular smooth muscle by retinoids, 113, 1448
 Hirst, B.H. *see* Thwaites, D.T., 113, 1050
 Hirst, R.A. *see* Atcheson, R., 111, 787
 Hisayama, T. *see* Moritoki, H., 111, 655
 Hitri, A. *see* Karoum, F., 113, 1391
 Hittinger, L. *see* Su, J., 113, 395
 Hjorth, S.A. *see* Schambye, H.T., 113, 331
 Ho, J.K.S. *see* Qian, Y.-m., 113, 369
 Hodgson, W.C. *see* James, G.M., 111, 370
 Hodson, H.F. *see* Salas, E., 112, 1071
 Hogg, R.C., Wang, Q. and Large, W.A. Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977
 Hogg, R.C., Wang, Q. and Large, W.A. Effects of C1 channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333
 Hoggie, M. *see* Richard, V., 113, 869
 Holland, E. *see* Delamere, F., 111, 983
 Hollenberg, M.D. *see* Laniyonu, A., 113, 403
 Holzer, P. and Jocić, M. Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
 Holzer, P. *see* Herbert, M.K., 111, 681
 Holzer, P. *see* Maggi, C.A., 112, 161
 Holzer, P. *see* Pethö, G., 113, 1036
 Höнак, D. *see* Löscher, W., 112, 97
 Höнак, D. *see* Löscher, W., 113, 1349
 Hope, P.J. *see* Lang, C.W., 112, 951
 Horie, S., Yano, S. and Watanabe, K. Differential effects of Na^+/K^+ -ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87
 Horn, J.P. *see* Shen, W.-X., 113, 898
 Hoskins, S.L. *see* Davenport, A.P., 113, 1303
 Hosokawa, M. *see* Suthamnatpong, N., 112, 676
 Hosoki, R., Yanagisawa, M., Guo, J.-Z., Yoshioka, K., Machara, T. and Otsuka, M. Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141
 Hosoki, R. *see* Suzuki, H., 113, 310
 Hosseinzadeh, H. and Stone, T.W. The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316
 Hostettmann, K. *see* Hay, A.J.B., 112, 9
 Hou, Y.-S. *see* Ko, F.-N., 112, 1174
 Hoult, J.R.S. *see* Hay, A.J.B., 112, 9
 Hourani, S.M.O. *see* Bailey, S.J., 112, 219
 Hourani, S.M.O. *see* Johnson, C.R., 113, 1191
 Hourani, S.M.O. *see* Peachey, J.A., 113, 1249
 Hoyle, C.H.V. *see* Ziganshin, A.U., 111, 923
 Hoyle, C.H.V. *see* Ziganshin, A.U., 113, 669
 Hsia, H.-C. *see* Yang, C.M., 111, 21
 Hsia, H.-C. *see* Yang, C.M., 112, 781
 Hsieh, J.-T. *see* Yang, C.M., 111, 21
 Hsieh, J.-T. *see* Yang, C.M., 111, 777
 Hsieh, J.-T. *see* Yang, C.M., 112, 781
 Hsu, K.S. *see* Kang, J.J., 112, 1216
 Hsu, M.F. *see* Wang, J.P., 113, 945
 Huang, C., Davis, G. and Johns, E.J. Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111
 Huang, C.-C. and Gean, P.-W. Paired-pulse depression of the *N*-methyl-D-aspartate receptor-mediated synaptic potentials in the amygdala, 113, 1029
 Huang, F.-L. *see* Fu, W.-M., 111, 880
 Hubert, J.P., Delumeau, J.C., Glowinski, J., Prémont, J. and Doble, A. Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
 Hughes, S.R. and Brain, S.D. Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from

capsaicin-sensitive nerves in rabbit skin, 111, 425
 Huidobro-Toro, J.P. *see* Donoso, M.V., 111, 473
 Hulin, P. *see* Belachgar, F., 112, 465
 Humphrey, P.P.A. *see* Cuthbert, A.W., 112, 31
 Humphrey, P.P.A. *see* Gale, J.D., 111, 332
 Humphrey, P.P.A. *see* McKeen, E.S., 113, 628
 Humphrey, P.P.A. *see* Matthews, E.K., 113, 87
 Humphrey, P.P.A. *see* Trezise, D.J., 112, 282
 Humphrey, P.P.A. *see* Trezise, D.J., 113, 463
 Humphries, R.G., Tomlinson, W., Ingall, A.H., Cage, P.A. and Leff, P. FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2T}-purinoceptors, 113, 1057
 Hunt, A.A.E. *see* Hilditch, A., 111, 137
 Hunt, J.M., Redman, R.S. and Silinsky, E.M. Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753
 Hunter, J.C., Atwal, P., Woodruff, G.N. and Singh, L. Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002
 Hurlston, R.M. *see* Chin, J.P.F., 112, 566
 Hussain, A.S., Poklewsko-Koziell, M., Brien, J.F., Marks, G.S. and Nakatsu, K. Nitric oxide formation from glyceryl trinitrate by rabbit aortic strip: detection by rabbit taenia coli concurrent with vasorelaxation, 113, 1508
 Hussain, S.N.A. *see* Ward, M.E., 112, 65
 Hutcheson, I.R., Smith, J.A. and Griffith, T.M. Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolaemic rabbits, 113, 190
 Hutson, P.H. *see* Tricklebank, M.D., 113, 729
 Huupponen, R. *see* Santti, E., 113, 1231
 Huwiler, A. and Pfeilschifter, J. Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455
 Hyodo, M. *see* Minami, T., 112, 735

I

Iacono, G. and Vassalle, M. Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289
 Iben, L.G., Mahle, C.D. and Yocca, F.D. Differential sensitivity of ³H-agonist binding pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400
 Ichikawa, S. *see* Kitazawa, T., 111, 288
 Ichinose, M. *see* Yamauchi, H., 112, 958
 Igawa, Y. *see* Ishizuka, O., 113, 111
 Ihara, M. *see* Fukuroda, T., 113, 336
 Iino, M. *see* Kasai, Y., 112, 1132
 Ilzerman, A.P. *see* Crack, B.E., 113, 1432
 Ikeda, M. *see* Nakazawa, K., 113, 296
 Illes, P. *see* Langosch, J.M., 113, 29
 Illes, P. *see* Nörenberg, W., 111, 942
 Im, W.B., Pregenzer, J.F. and Thomsen, D.R. Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
 Imagawa, J.-i. *see* Kamei, K., 113, 889
 Imagawa, M. *see* Moritoki, H., 111, 655
 Imai, M. *see* Muto, S., 111, 173
 Imaizumi, Y. *see* Muraki, K., 111, 1154
 Imaizumi, Y. *see* Yokota, Y., 113, 324
 Ingall, A.H. *see* Humphries, R.G., 113, 1057
 Inomata, H. *see* Chen, S., 112, 1137
 Inoue, H. *see* Yamauchi, H., 112, 958
 Inoue, K. *see* Koizumi, S., 112, 992
 Inoue, K. *see* Nakazawa, K., 113, 296
 Inoue, R. *see* Chen, S., 112, 1137
 Ioculano, M. *see* Squadrato, F., 113, 912
 Ioculano, M. *see* Squadrato, F., 113, 1153
 Iravani, M.M. and Zar, M.A. Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
 Iravani, M.M. and Zar, M.A. Presence of neuropeptide Y in the rat seminal vesicle and its effects on noradrenaline- and nerve-induced contractions, 113, 877
 Iredale, P.A., Alexander, S.P.H. and Hill, S.J. Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium

mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252
 Irie, K. *see* Chiba, K., 112, 837
 Ishida, M. *see* Saitoh, T., 113, 165
 Ishigai, Y. *see* Chiba, K., 112, 837
 Ishii, A. *see* Kitazawa, T., 111, 288
 Ishikawa, J. *see* Yamauchi, H., 112, 958
 Ishikawa, K. *see* Fukuroda, T., 113, 336
 Ishikawa, K. *see* Sato, Y., 112, 625
 Ishimoto, H. *see* Nakahata, N., 112, 299
 Ishizuka, O., Igawa, Y., Lecci, A., Maggi, C.A., Mattiasson, A. and Andersson, K.-E. Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
 Isono, K. *see* Chiyotani, A., 112, 571
 Ito, S. *see* Itoh, T., 111, 165
 Ito, S. *see* Minami, T., 112, 735
 Ito, S. *see* Nakazato, Y., 111, 935
 Ito, S. *see* Ohta, T., 112, 972
 Ito, Y. *see* Chen, S., 112, 1137
 Itoh, T., Ito, S., Shafiq, J. and Suzuki, H. Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
 Itoh, T., Suzuki, A. and Watanabe, Y. Effect of a peptide inhibitor of protein kinase C on G-protein-mediated increase in myofilament Ca²⁺-sensitivity in rabbit arterial skinned muscle, 111, 311
 Iversen, H.H., Gustafsson, L.E., Leone, A.M. and Wiklund, N.P. Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088
 Ivorra, M.D. *see* Chuliá, S., 113, 1377
 Iwao, H. *see* Kim, S., 113, 662
 Izawa, T. *see* Kashiwabara, T., 111, 278
 Izzo, A.A., Gaginella, T.S., Mascolo, N. and Capasso, F. Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228
 Izzo, A.A. *see* Capasso, F., 113, 1127

J

Jacobson, K.A. *see* Bo, X., 112, 1151
 Jacobson, K.A. *see* Boyer, J.L., 113, 614
 Jacoby, D.B. *see* Fryer, A.D., 112, 588
 Jakab, I. *see* Szilvassy, Z., 112, 999
 Jakschik, J. *see* Molderings, G.J., 111, 733
 Jakubsk, J. and Tuček, S. Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529
 James, G.M., Hodgson, W.C., Davis, E.A. and Haynes, J.M. Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370
 Jamieson, G.P. *see* Wiley, J.S., 112, 946
 Jane, D.E., St. J. Jones, P.L., Pook, P.C.-K., Tse, H.-W. and Watkins, J.C. Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809
 Jansen, F.P., Wu, T.S., Voss, H.-P., Steinbusch, H.W.M., Vollinga, R.C., Rademaker, B., Bast, A. and Timmerman, H. Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenopropit, to rat brain, 113, 355
 Jassar, B.S. *see* Chen, H., 113, 55
 Javan, C.M. *see* Shaunik, S., 113, 151
 Jean, T. *see* Galizzi, J.P., 113, 389
 Jennings, G.L. *see* Chin, J.P.F., 112, 566
 Jhanwar-Uniyal, M. *see* Santti, E., 113, 1231
 Jiang, C. and Collins, P. Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N^G-monomethyl-L-arginine but not glibenclamide, 111, 711
 Jino, H., Usui, H., Temma, S., Shirahase, H. and Kurashiki, K. An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
 Jobling, P. *see* Shen, W.-X., 113, 898
 Jocić, M. *see* Holzer, P., 112, 1181
 Jocić, M. *see* Pethö, G., 113, 1036
 Johansen, T. *see* Friis, U.G., 111, 1179
 John, G.W. *see* Le Grand, B., 112, 195
 Johns, E. *see* Wu, X.C., 112, 360

Johns, E. *see* Wu, X.C., 112, 369
 Johns, E.J. *see* Chamienia, A.L., 112, 214
 Johns, E.J. *see* Huang, C., 111, 111
 Johns, E.J. *see* Wong, P.S.K., 112, 881
 Johnson, C.R. and Hourani, S.M.O. Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191
 Johnson, I.R. *see* Rouse, W., 113, 1064
 Johnson, M. *see* Kanthakumar, K., 112, 493
 Johnson, P.J., Bornstein, J.C., Furness, J.B., Woppard, D.J. and Orrman-Rossiter, S.L. Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240
 Johnson, R.J. *see* Sirois, M.G., 111, 1111
 Johnstone, A.P. *see* Bogle, R.G., 111, 1257
 Jones, H.B. *see* McAuliffe, S.J.G., 112, 272
 Jones, R.L. *see* Qian, Y.-m., 113, 369
 Jones, R.L. *see* Wise, H., 113, 581
 Jones, R.M. *see* Lees, G., 113, 642
 Jordae, F.H. *see* De Meyer, G.R.Y., 112, 471
 Jordan, D. *see* Shepheard, S.L., 111, 616
 Jordan, R. *see* Raeburn, D., 113, 1423
 Joseph, D. *see* Coëffier, E., 113, 749
 Joulin, Y. *see* Bouaziz, H., 113, 717
 Jun, T. and Wennmalm, Å. NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
 Junien, J.-L. *see* Gué, M., 111, 930
 Juorio, A.V. *see* Berry, M.D., 113, 1159
 Juorio, A.V. *see* Zhu, M.-Y., 112, 23

K

Kaakkola, S. *see* Törnwall, M., 112, 13
 Kaeffer, N. *see* Richard, V., 113, 869
 Kahl, U., Langel, Ü., Bartfai, T. and Grundemar, L. Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
 Kahler, C.P. *see* Du Plooy, W.J., 111, 73
 Kähönen, M., Mäkinen, H., Arvola, P. and Pörsti, I. Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925
 Källström, B.-L., Sjöberg, J. and Waldeck, B. The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687
 Kamei, K., Yoshida, S., Imagawa, J.-i., Nabata, H. and Kuriyama, H. Regional and species differences in glyburide-sensitive K^+ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889
 Kanaide, H. *see* Aoki, H., 111, 489, 989
 Kanaide, H. *see* Ushio-Fukai, M., 113, 1173
 Kaneda, M., Fukui, K. and Doi, K. Activation of chloride current by P_2 -purinoreceptors in rat ventricular myocytes, 111, 1355
 Kang, J.-J., Hsu, K.S. and Lin-Shiau, S.Y. Effects of bipyridylum compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle, 112, 1216
 Kanthakumar, K., Cundell, D.R., Johnson, M., Wills, P.J., Taylor, G.W., Cole, P.J. and Wilson, R. Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493
 Kappelle, A.C., Biessels, G., Bravenboer, B., van Buren, T., Traber, J., de Wildt, D.J. and Gispens, W.H. Beneficial effect of the Ca^{2+} antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887
 Karet, F.E. *see* Davenport, A.P., 113, 1303
 Karlsson, J.-A. *see* Raeburn, D., 113, 1423
 Karmazyn, M. *see* Khandoudi, N., 112, 1007
 Karoum, F., Chrapusta, S.J., Brinjak, R., Hitri, A. and Wyatt, R.J. Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
 Karppanen, H. *see* Mervaala, E.M.A., 111, 1189
 Karppanen, H. *see* Mervaala, E.M.A., 112, 640
 Kasai, Y., Iino, M., Tsutsumi, O., Taketani, Y. and Endo, M. Effects of cyclopiazonic acid on rhythmic contractions in uterine smooth muscle bundles of the rat, 112, 1132
 Kasakov, L., Belai, A., Vlaskovska, M. and Burnstock, G. Noradrenergic-nitroergic interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403
 Kasamatsu, K. *see* Kawabata, A., 111, 503
 Kase, H. *see* Kurokawa, M., 113, 43
 Kashiwabara, T., Ogawa, N., Izawa, T. and Fukushima, H. Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278
 Kask, A. *see* Törnwall, M., 112, 13
 Kasuya, Y. *see* Yasuda, N., 113, 21
 Kathmann, M. *see* Schlicker, E., 112, 1043
 Kathmann, M. *see* Schlicker, E., 113, 657
 Katsuragi, T., Ogawa, S. and Furukawa, T. Contribution of intra- and extracellular Ca^{2+} to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
 Kaumann, A.J. *see* Taylor, E.M., 111, 264
 Kawabata, A., Kasamatsu, K., Umeda, N. and Takagi, H. The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503
 Kawabata, A., Manabe, S., Manabe, Y. and Takagi, H. Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
 Kawabata, A., Manabe, S. and Takagi, H. Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817
 Kawai, Y. and Ohhashi, T. Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635
 Kaye, D.M. *see* Chin, J.P.F., 112, 566
 Kedzie, K.M. *see* Regan, J.W., 112, 377
 Keef, K.D. *see* Eckman, D.M., 111, 1053
 Keenan, A.K. *see* Doolan, C.M., 111, 509
 Kehl, S.J. Block by capsaicin of voltage-gated K^+ currents in melanotrophs of the rat pituitary, 112, 616
 Keiser, H.R. *see* Abassi, Z.A., 113, 204
 Keith, R.A., Mangano, T.J., DeFeo, P.A., Ernst, G.E. and Warawa, E.J. Differential inhibition of neuronal calcium entry and [3 H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
 Kellock, N.A. *see* Plevin, R., 112, 311
 Kelly, S.S. and Ferry, C.B. The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747
 Kemp, J.A. *see* Brown, N.A., 113, 600
 Kemp, J.A. *see* Cartmell, J., 111, 364
 Kemp, J.A. *see* Seabrook, G.R., 111, 391
 Kemp, J.A. *see* Seabrook, G.R., 111, 1061
 Kemp, P.A., Gardiner, S.M., March, J.E., Bennett, T. and Rubin, P.C. Effects of N^G -nitro-L-arginine methyl ester on regional haemodynamic responses to $MgSO_4$ in conscious rats, 111, 325
 Kemp, P.A. *see* Gardiner, S.M., 111, 589
 Kemp, P.A. *see* Gardiner, S.M., 112, 477
 Kemp, P.A. *see* Gardiner, S.M., 112, 823
 Kemp, P.A. *see* Gardiner, S.M., 113, 129
 Kendall, D.A. *see* Alexander, S.P.H., 111, 185
 Kendall, D.A. *see* Alexander, S.P.H., 113, 1501
 Kendall, D.A. *see* Cartmell, J., 111, 364
 Kendall, D.A. *see* Hernández, F., 113, 216
 Kendall, D.A. *see* Wright, I.K., 112, 1146
 Kengatharan, M. *see* Mackay, D., 111, 227
 Kennedy, C. *see* Evans, R.J., 113, 853
 Kennedy, C. *see* McLaren, G.J., 111, 913
 Kennedy, I. *see* Trezise, D.J., 112, 282
 Kennedy, I. *see* Trezise, D.J., 113, 463
 Kennett, G.A., Wood, M.D., Glen, A., Grewal, S., Forbes, I., Gadre, A. and Blackburn, T.P. *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797
 Kennett, G.A. *see* Wardle, K.A., 112, 789
 Kenny, B.A., Naylor, A.M., Greengrass, P.M., Russell, M.J., Friend, S.J., Read, A.M. and Wyllie, M.G. Pharmacological properties of the cloned $\alpha_{1A/1B}$ -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral cortex and vas deferens, 111, 1003
 Khandoudi, N., Moffat, M.P. and Karmazyn, M. Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison

with phorbol ester, 112, 1007
 Khawaja, A.M. *see* Paul, W., 111, 159
 Kidd, B.L. *see* Cruwys, S.C., 113, 940
 Kiff, R.J., Moss, D.W. and Moncada, S. Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 496
 Kigoshi, S. *see* Kohno, Y., 112, 1167
 Kilfeather, S.A. *see* Herd, C.M., 111, 7
 Kilpatrick, E.V. and Cocks, T.M. Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557
 Kim, S., Ohta, K., Harnaguchi, A., Omura, T., Tominaga, K., Yukimura, T., Miura, K., Tanaka, M. and Iwao, H. AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662
 Kipnis, T.L. *see* Tambourgi, D.V., 112, 723
 Kirk, I.P. *see* Kurokawa, M., 113, 43
 Kirkpatrick, K.A. *see* Kurokawa, M., 113, 43
 Kirstetter, P. *see* Moreau, R., 112, 649
 Kita, Y., Hirasawa, Y., Yoshida, K. and Maeda, K. Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385
 Kita, Y., Ozaki, R., Sakai, S., Sugimoto, T., Hirasawa, Y., Ohtsuka, M., Senoh, H., Yoshida, K. and Maeda, K. Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137
 Kita, Y., Sugimoto, T., Hirasawa, Y., Yoshida, K. and Maeda, K. Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5
 Kitabgi, P. *see* Dubuc, I., 112, 352
 Kitazawa, T., Ichikawa, S., Yokoyama, T., Ishii, A. and Shuto, K. Stimulating action of KW-5139 (Leu¹³-motilin) on gastrointestinal motility in the rabbit, 111, 288
 Kitchen, I. *see* Peachey, J.A., 113, 1249
 Klaus, W. *see* Rump, A.F.E., 112, 757
 Kleene, S.J. Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469
 Knapp, J. *see* Michel, M.C., 113, 240
 Knaus, E.E. *see* Ellory, J.C., 111, 903
 Knock, G.A. *see* Walsh, D.A., 112, 435
 Knott, P.G. *see* Goldie, R.G., 112, 749
 Knowles, M. *see* Seabrook, G.R., 112, 728
 Knowles, M.R. *see* Seabrook, G.R., 111, 1061
 Knox, A. *see* Delamere, F., 111, 983
 Ko, F.-N., Guh, J.-H., Yu, S.-M., Hou, Y.-S., Wu, Y.-C. and Teng, C.-M. (-)-Discretamine, a selective α_{1D}-adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
 Ko, F.-N. *see* Lin, C.-H., 113, 275
 Ko, F.-N. *see* Lin, C.-H., 113, 1464
 Kobayashi, S. *see* Aoki, H., 111, 489
 Kobayashi, S. *see* Aoki, H., 111, 989
 Kockx, M. *see* De Meyer, G.R.Y., 112, 471
 Koenig, J.A. and Edwardson, J.M. Routes of delivery of muscarinic acetylcholine receptors to the plasma membrane in NG108-15 cells, 111, 1023
 Kohno, Y., Saito, H., Takita, M., Kigoshi, S. and Muramatsu, I. Heterogeneity of α₁-adrenoceptor subtypes involved in adrenergic contractions of dog blood vessels, 112, 1167
 Koizumi, S., Watano, T., Nakazawa, K. and Inoue, K. Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
 Koizumi, S. *see* Nakazawa, K., 113, 296
 Koizumi, T. *see* Hatano, A., 113, 723
 Kojda, G., Beck, J.K., Meyer, W. and Noack, E. Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533
 Kojima, M. *see* Dohi, Y., 111, 673
 Kokubun, S. *see* Suzuki, H., 112, 117
 Koltai, M. *see* Szilvassy, Z., 112, 999
 Komeichi, H. *see* Moreau, R., 112, 649
 Komeyama, T. *see* Hatano, A., 113, 723
 Kondo, M. *see* Chiyotani, A., 112, 571
 Kondoh, S. *see* Tanabe, J.-i., 113, 1480
 Kondoh, W. *see* Moritoki, H., 111, 655
 Konno, K. *see* Chiyotani, A., 112, 571
 Kooyman, A.R., van Hooft, J.A., Vanderheijden, P.M.L. and Vijverberg, H.P.M. Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541
 Korbut, R., Warner, T.D., Gryglewski, R.J. and Vane, J.R. The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289
 Kotsonis, P. and Majewski, H. Facilitation by procaterol, a β-adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
 Kötting, A. *see* Michel, M.C., 112, 499
 Koulu, M. *see* Santti, E., 113, 1231
 Kozlowski, R.Z., Goodstadt, L.J., Twist, V.W. and Powell, T. Modulation of cardiac L-type Ca²⁺ channels by GTPγS in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
 Kreutner, W. *see* Bolser, D.C., 113, 1344
 Krier, J. *see* Venkova, K., 112, 1237
 Krnjević, K. *see* Erdemli, G., 113, 411
 Kruck, Z.L. *see* Wieczorek, W.J., 111, 829
 Kuc, R.E. *see* Davenport, A.P., 111, 4
 Kuc, R.E. *see* Davenport, A.P., 113, 1303
 Kuc, R.E. *see* Maguire, J.J., 113, 49
 Kühberger, E., Groschner, K., Kukovetz, W.R. and Brunner, F. The role of myoendothelial cell contact in non-nitric oxide-, non-prostanoid-mediated endothelium-dependent relaxation of porcine coronary artery, 113, 1289
 Kuijpers, G.A.J., Vergara, L.A., Calvo, S. and Yadid, G. Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
 Kukovetz, W.R. *see* Kühberger, E., 113, 1289
 Kunz, H.M.D. and Pfeilschifter, J. Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1
 Kunze, D.L. *see* Hasséssian, H., 112, 1118
 Kuo, J.S. *see* Liao, J.-F., 111, 439
 Kurahashi, K. *see* Jino, H., 111, 435
 Kurenný, D.E. *see* Chen, H., 113, 55
 Kurihara, T. *see* Suzuki, H., 113, 310
 Kuriyama, H. *see* Kamei, K., 113, 889
 Kuroda, A. *see* Saito, K., 111, 465
 Kurokawa, M., Kirk, I.P., Kirkpatrick, K.A., Kase, H. and Richardson, P.J. Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
 Kurosawa, A. *see* Arimura, A., 111, 339
 Kurz, A.K. *see* Bültmann, R., 111, 151
 Kus, M. *see* Williams, F.M., 111, 1123
 Kwan, C.-Y. *see* Leung, Y.-M., 113, 767
 Kwan, C.Y. *see* Low, A.M., 112, 604
 Kwon, O.-J. *see* Mitchell, J.A., 113, 1008

L

Laakso, J. *see* Mervaala, E.M.A., 111, 1189
 Laakso, J. *see* Mervaala, E.M.A., 112, 640
 Lach, E. *see* Trifilieff, A., 111, 1228
 Lacroix, J.S., Ulman, L.G. and Potter, E.K. Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y₂ prejunctional receptor, 113, 479
 Ladinsky, H. *see* Sagrada, A., 113, 711
 Lagente, V. *see* Pires, A.L.A., 113, 994
 Laguna, J.C. *see* Alegret, M., 112, 551
 Lakhiani, P.P., Amenta, F., Napoleone, P., Felici, L. and Eikenburg, D.C. Pharmacological characterization and anatomical localization of prejunctional β-adrenoceptors in the rat kidney, 111, 1296
 Lamb, J. *see* Turner, N.C., 111, 1047
 Lamb, J. *see* Turner, N.C., 111, 1198
 Lambert, D.G. *see* Atcheson, R., 111, 787
 Lamberti, C. *see* Malmberg-Aiello, P., 111, 1269
 Lambrecht, G. *see* McLaren, G.J., 111, 913
 Lambrecht, G. *see* Waelbroeck, M., 112, 505
 Lambrecht, G. *see* Windscheif, U., 113, 1015
 Lambrecht, G. *see* Ziganshin, A.U., 111, 923
 Lang, C.W., Hope, P.J., Grubb, B.D. and Duggan, A.W. Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951
 Lang, R.J. *see* Zhang, Y., 113, 431
 Langel, Ü. *see* Kahl, U., 111, 1129

Langosch, J.M., Gebicke-Haerter, P.J., Nörenberg, W. and Illes, P. Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29

Langosch, J.M. *see* Nörenberg, W., 111, 942

Laniyonu, A., Saifeddine, M., Ahmad, S. and Hollenberg, M.D. Regulation of vascular and gastric smooth muscle contractility by perva-nadate, 113, 403

Laplace, M. *see* Su, J., 113, 395

Large, W.A. *see* Hogg, R.C., 111, 1333

Large, W.A. *see* Hogg, R.C., 112, 977

Larrazet, F., Chauveau, M., Weber, S., Lockhart, A. and Frossard, N. Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649

Lasheras, B. *see* Ramírez, M.J., 111, 419

Lasheras, B. *see* Ramírez, M.J., 113, 77

László, F., Whittle, B.J.R. and Moncada, S. Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131

Laszlo, F., Whittle, B.J.R. and Moncada, S. Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309

Lawrence, L. *see* Paul, W., 111, 159

Le Fur, G. *see* Dubuc, I., 112, 352

Le Grand, B., Marty, A., Vié, B., Patoiseau, J.-F. and John, G.W. Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractures in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195

Lebrec, D. *see* Moreau, R., 112, 649

Lecci, A. *see* Ahluwalia, A., 111, 1017

Lecci, A. *see* Ishizuka, O., 113, 111

Lecoffre, C. *see* Berry, C.N., 113, 1209

Lécrivain, A. *see* Chamelian, M., 113, 317

Lee, F.-Y. *see* Liao, J.-F., 111, 439

Lee, K., Ozanne, S.E., Hales, C.N. and Ashford, M.L.J. Mg^{2+} -dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632

Lee, S.-S. *see* Wu, M.-H., 113, 221

Lee, S.S. *see* Pak, J.-M., 112, 1209

Lee, W.S., Moussaoui, S.M. and Moskowitz, M.A. Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

Lee, Y.-M., Chern, J.-W. and Yen, M.-H. Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031

Lees, G., Munday, I.T., Edwards, M.D. and Jones, R.M. *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Lees, G.M. *see* Cunningham, S.M.C., 113, 883

Leeson, P.D. *see* Tricklebank, M.D., 113, 729

Lefebvre, R.A. *see* Barbier, A.J.M., 111, 1280

Leff, P. *see* Crack, B.E., 113, 1432

Leff, P. *see* Humphries, R.G., 113, 1057

Legat, F.J., Griesbacher, T. and Lembeck, F. Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453

Leishman, D.J., Boeijinga, P.H. and Galvan, M. Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorsolateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318

Lembeck, F. *see* Griesbacher, T., 112, 1004

Lembeck, F. *see* Legat, F.J., 112, 453

Leone, A.M. *see* Iversen, H.H., 113, 1088

Leung, Y.-M., Kwan, C.-Y. and Loh, T.-T. Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767

Leurs, R., Smit, M.J., Menge, W.M.B.P. and Timmerman, H. Pharmacological characterization of the human histamine H₂ receptor stably expressed in Chinese hamster ovary cells, 112, 847

Levin, R.M. *see* Tammela, T.L.J., 113, 195

Lewin, J. *see* Donoso, M.V., 111, 473

Lewis, S.A. *see* Raeburn, D., 113, 1423

Li, C.G. *see* Rand, M.J., 111, 1089

Li, D.Y., Varma, D.R. and Chemtob, S. Ontogenetic increase in PGE₂ and PGF_{2 α} receptor density in brain microvessels of pigs, 112, 59

Li, P., Penner, S.B. and Smyth, D.D. Attenuated renal response to moxonidine and rilmenidine in one kidney-one clip hypertensive rats, 112, 200

Liao, J.-F., Yu, P.-C., Lin, H.-C., Lee, F.-Y., Kuo, J.S. and Yang, M.C.-M. Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439

Lieb, W.R. *see* Hall, A.C., 112, 906

Likungu, J. *see* Molderings, G.J., 111, 733

Lin, C.-H., Chang, G.-J., Su, M.-J., Wu, Y.-C., Teng, C.-M. and Ko, F.-N. Pharmacological characteristics of liriiodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275

Lin, C.-H., Yang, C.-M., Ko, F.-N., Wu, Y.-C. and Teng, C.-M. Antimuscarinic action of liriiodenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle, 113, 1464

Lin, H.-C. *see* Liao, J.-F., 111, 439

Lin, Y.-Q. *see* Nicolarakis, P.J., 111, 521

Lin-Shiau, S.Y. *see* Kang, J.J., 112, 1216

Linnoila, M. *see* Durcan, M.J., 112, 855

Little, H.J. *see* Rabbani, M., 111, 89

Little, H.J. *see* Watson, W.P., 112, 1017

Liu, X., Gillespie, J.S. and Martin, W. Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of S-nitrosothiols, 111, 1287

Lluch, S. *see* Martínez, M.C., 113, 419

Lochot, S. *see* Berry, C.N., 113, 1209

Lock, K.R. *see* Alemayehu, A., 111, 205

Lockhart, A. *see* Larrazet, F., 113, 649

Loh, T.-T. *see* Leung, Y.-M., 113, 767

Loirand, G. *see* Pacaud, P., 113, 457

Loizzo, A. *see* Pieretti, S., 113, 1416

Loke, J.C.P. *see* Low, A.M., 112, 604

Loke, K.E., Sobey, C.G., Dusting, G.J. and Woodman, O.L. Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630

Long, S.K. *see* Siarey, R.J., 111, 497

Longhurst, P.A. *see* Tammela, T.L.J., 113, 195

Longmore, J., Razzaque, Z., Shaw, D. and Hill, R.G. Differences in the effects of NK₁-receptor antagonists, (±)-CP 96,345 and CP 99,994, on agonist-induced responses in guinea-pig trachea, 112, 176

Longmore, J. *see* Tan, K.K.C., 111, 703

Lonovics, J. *see* Szilvassy, Z., 112, 999

Lopes-Martins, R.A.B., Antunes, E., Oliva, M.L.V., Sampaio, C.A.M., Burton, J. and de Nucci, G. Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81

Lopez-Belmonte, J. and Whittle, B.J.R. The involvement of endothelial dysfunction, nitric oxide and prostanoids in the rat gastric microcirculatory responses to endothelin-1, 112, 267

Löscher, W., Wlaź, P., Rundfeldt, C., Baran, H. and Hönack, D. Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

Löscher, W. and Hönack, D. Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

Losinski, A. *see* Alexander, S.P.H., 111, 185

Low, A.M., Bowdish, D.M., Prashad, T.R. and Gaspar, V. Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263

Low, A.M., Loke, J.C.P., Kwan, C.Y. and Daniel, E.E. Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604

Lucarini, J.M. *see* Pruneau, D., 111, 1029

Lucchelli, A., Santagostino-Barbone, M.G., Barbieri, A. and Tonini, M. A pharmacological analysis of receptors mediating the excitatory response to 5-hydroxytryptamine in the guinea-pig isolated trachea, 112, 763

Lucchesi, B.R. *see* Friedrichs, G.S., 113, 209

Lucchesi, B.R., Rote, W.E., Driscoll, E.M. and Mu, D.-X. Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Luciani, S. *see* Bova, S., 111, 1067
 Lues, I. *see* Barber, A., 111, 843
 Lugnier, C. *see* Chuliá, S., 113, 1377
 Lugnier, C. *see* Eckly, A.E., 113, 445
 Lung, M.A. Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411
 Luo, S.-F. *see* Yang, C.M., 111, 21
 Luo, S.-F. *see* Yang, C.M., 112, 781
 Luscombe, G.P. *see* Scott, G., 111, 97
 Lustig, L. *see* Bacman, S., 111, 271
 Luu, T.N. *see* Dashwood, M.R., 112, 386
 Ly, S.M. *see* Galizzi, J.P., 113, 389
 Lydford, S.J. and McKechnie, K. Characterization of the prostaglandin E₂ sensitive (EP)-receptor in the rat isolated trachea, 112, 133

M

McAllister, G. *see* Seabrook, G.R., 111, 391
 McAllister, G. *see* Seabrook, G.R., 111, 1061
 McAllister, G. *see* Seabrook, G.R., 112, 728
 MacAllister, R.J., Fickling, S.A., Whitley, G.St.J. and Vallance, P. Metabolism of methylarginines by human vasculature; implications for the regulation of nitric oxide synthesis, 112, 43
 McAuliffe, S.J.G., Moors, J.A. and Jones, H.B. Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272
 McBean, G.J. Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, o-aminoadipate, 113, 536
 McCall, R. *see* Nally, J.E., 111, 1163
 McColl, S.R. *see* McDonald, P.P., 111, 852
 McCreath, G., Hall, I.P. and Hill, S.J. Agonist-induced desensitization of histamine H₁ receptor-mediated inositol phospholipid hydrolysis in human umbilical vein endothelial cells, 113, 823
 McCulloch, J. *see* Dawson, D.A., 113, 247
 MacDermot, J. *see* Shaunak, S., 113, 151
 MacDonald, A., Forbes, I.J., Gallacher, D., Heeps, G. and McLaughlin, D.P. Adrenocceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576
 MacDonald, J.F. *see* Church, J., 113, 499
 McDonald, P.P., McColl, S.R., Braquet, P. and Borgeat, P. Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852
 McDonald, R.L., Vaughan, P.F.T. and Peers, C. Muscarinic (M₁) receptor-mediated inhibition of K⁺-evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L- and N-type Ca²⁺ channels, 113, 621
 McDonald, T.V. *see* Chung, S.C., 113, 861
 McFadzean, I. *see* Gibson, A., 113, 1494
 McFerran, B.W. and Guild, S.B. Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
 McGarry, S.J. and Williams, A.J. Activation of the sheep cardiac sarcoplasmic reticulum Ca²⁺-release channel by analogues of sulmazole, 111, 1212
 McGiff, J.C. *see* Fulton, D., 113, 954
 McGrath, J.C. *see* Nally, J.E., 111, 1163
 McGrath, J.C. *see* Nally, J.E., 113, 1328
 McGrath, J.C. *see* Smith, G.C.S., 111, 1245
 Machado, A. *see* Santiago, M., 111, 837
 Mackay, D. A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219
 Mackay, D. and Kengatharan, M. pK_i values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227
 McKechnie, K. *see* Lydford, S.J., 112, 133
 McKechnie, K.C.W. *see* Crack, B.E., 113, 1432
 McKeen, E.S., Feniuk W. and Humphrey, P.P.A. Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628
 MacLagan, J. *see* Wessler, I., 113, 1221
 McLaren, G.J., Lambrecht, G., Mutschler, E., Bäumert, H.G., Sneddon, P. and Kennedy, C. Investigation of the actions of PPADS, a novel P_{2X}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913
 McLaughlin, D.P. *see* MacDonald, A., 112, 576
 Maclouf, J. *see* Zuany-Amorim, C., 112, 1230
 McMullen, D. *see* Moreland, S., 112, 704
 McMurdo, L., Thiemermann, C. and Vane, J.R. The effects of the endothelin ET_A receptor antagonist, FR 139317, on infarct size in a rabbit model of acute myocardial ischaemia and reperfusion, 112, 75
 McPherson, G.A. and Piekarska, A.E. Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
 Macrae, I.M. *see* Dawson, D.A., 113, 247
 MacVinish, L.J. *see* Cuthbert, A.W., 112, 31
 Maeda, K. *see* Kita, Y., 113, 5
 Maeda, K. *see* Kita, Y., 113, 385
 Maeda, K. *see* Kita, Y., 113, 1137
 Maehara, T. *see* Hosoki, R., 113, 1141
 Maestrone, E., Magnelli, V., Nobile, M. and Usai, C. Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
 Maeyama, K. *see* Yamauchi, H., 112, 958
 Maggi, C.A., Giuliani, S. and Santicioli, P. Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
 Maggi, C.A., Patacchini, R., Bartho, L., Holzer, P. and Santicioli, P. Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161
 Maggi, C.A., Patacchini, R., Meini, S., Quartara, L., Sisto, A., Potier, E., Giuliani, S. and Giachetti, A. Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
 Maggi, C.A. *see* Ahluwalia, A., 111, 1017
 Maggi, C.A. *see* Astolfi, M., 111, 570
 Maggi, C.A. *see* Belvisi, M.G., 111, 103
 Maggi, C.A. *see* Ishizuka, O., 113, 111
 Maggi, C.A. *see* Meini, S., 111, 739
 Maggi, C.A. *see* Meini, S., 112, 1123
 Maggi, C.A. *see* Santicioli, P., 113, 588
 Maggi, C.A. *see* Zagorodnyuk, V., 112, 1077
 Magnelli, V. *see* Maestrone, E., 111, 283
 Maguire, J.J., Kuc, R.E., O'Reilly, G. and Davenport, A.P. Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49
 Maguire, J.J. *see* Davenport, A.P., 111, 4
 Mahle, C.D. *see* Iben, L.G., 113, 1400
 Maillard, M. *see* Bo, X., 112, 1151
 Mailman, D. Differential effects of luminal L-arginine and N^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
 Main, M.J. *see* Grantham, C.J., 111, 483
 Majewski, H. *see* Kotsonis, P., 113, 781
 Makki, T., Talom, R.T., Niederhoffer, N., Amin, F., Tankosic, P., Mertes, P.-M. and Atkinson, J. Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555
 Mäkinen, H. *see* Kähönen, M., 112, 925
 Malcangio, M. and Bowery, N.G. Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
 Malcangio, M. and Bowery, N.G. Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_A receptor system, 113, 1561
 Mallet, A.I. *see* Murphy, L.J., 113, 137
 Malmberg-Aiello, P., Lamberti, C., Ghelardini, C., Giotti, A. and Bartolini, A. Role of histamine in rodent antinociception, 111, 1269
 Malva, J.O., Carvalho, A.P. and Carvalho, C.M. Modulation of dopamine and noradrenaline release and of intracellular Ca²⁺ concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
 Manabe, S. *see* Kawabata, A., 112, 547
 Manabe, S. *see* Kawabata, A., 112, 817
 Manabe, Y. *see* Kawabata, A., 112, 547
 Manahan-Vaughan, D., Anwyl, R. and Rowan, M.J. Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083
 Mangano, T.J. *see* Keith, R.A., 113, 379

Manicourt, D.H., Egeren, A.D.-V., Haazen, L. and Nagant de Duxchaisnes, C. Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Mann, G.E. *see* Baydoun, A.R., 112, 487

Männistö, P.T. *see* Törnwall, M., 112, 13

Manzini, S. *see* Astolfi, M., 111, 570

Maragoudakis, M.E. *see* Pipili-Synetos, E., 111, 894

March, J.E. *see* Gardiner, S.M., 111, 589

March, J.E. *see* Gardiner, S.M., 112, 477

March, J.E. *see* Gardiner, S.M., 112, 823

March, J.E. *see* Kemp, P.A., 111, 325

Marchionni, D. *see* Roccon, A., 111, 145

Marks, G.S. *see* Hussain, A.S., 113, 1508

Markus, R.P. *see* Ferreira, Z.S., 112, 107

Marleau, S., Dallaire, N., Poubelle, P.E. and Borgeat, P. Metabolic disposition of leukotriene B₄ (LTB₄) and oxidation-resistant analogues of LTB₄ in conscious rabbits, 112, 654

Marsal, J. *see* Cantí, C., 112, 19

Marsh, K.A. and Hill, S.J. Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934

Marthan, R. *see* Savineau, J.-P., 111, 978

Martí, E. *see* Cantí, C., 112, 19

Martin, D.K., Nakaya, Y., Wyse, K.R. and Campbell, T.J. Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873

Martín, M.L. *see* Morán, A., 113, 1358

Martin, S.W. and Broadley, K.J. Effects of chronic intravenous infusions of doxepamine and isoprenaline to rats on D₁-, B₁- and B₂-receptor-mediated responses, 112, 595

Martin, W. *see* Liu, X., 111, 1287

Martínez, M.C., Vila, J.M., Aldasoro, M., Medina, P., Flor, B. and Lluch, S. Relaxation of human isolated mesenteric arteries by vasopressin and desmopressin, 113, 419

Martins, M.A. *see* Pires, A.L.A., 113, 994

Marty, A. *see* Le Grand, B., 112, 195

Maruyama, M. *see* Saitoh, T., 113, 165

Marwood, R. *see* Rupniak, N.M.J., 113, 1487

Masaki, T. *see* Yasuda, N., 113, 21

Mascolo, N. *see* Capasso, F., 113, 1127

Mascolo, N. *see* Izzo, A.A., 113, 228

Mason, R. *see* Scott, G., 111, 97

Masumura, S. *see* Shinozuka, K., 113, 1203

Mathie, A. *see* Wooltorton, J.R.A., 111, 377

Mathie, R.T. *see* Browse, D.J., 113, 987

Matsuda, T. *see* Yokota, Y., 113, 324

Matsuda, Y. *see* Rutherford, R.A.D., 113, 931

Matsuki, N. *see* Torii, Y., 111, 431

Matthews, E.K., Seaton, E.D., Forsyth, M.J. and Humphrey, P.P.A. Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Matthews, M.L. *see* Sillence, M.N., 111, 866

Matthews, R. *see* Einstein, R., 111, 199

Mattiasson, A. *see* Ishizuka, O., 113, 111

Mauler, F. *see* Barber, A., 111, 843

Mauler, F. *see* Barber, A., 113, 1317

Maurice, M.-C. *see* D'Orléans-Juste, P., 113, 1257

Maynard, K.I. and Burnstock, G. Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123

Mazzaglia, G. *see* Calapai, G., 111, 1328

Medina, P. *see* Martínez, M.C., 113, 419

Meini, S., Patacchini, R. and Maggi, C.A. Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Meini, S. and Maggi, C.A. Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123

Meini, S. *see* Astolfi, M., 111, 570

Meini, S. *see* Maggi, C.A., 112, 150

Meisner, M. *see* Tschaikowsky, K., 113, 664

Mellander, S. *see* Ekelund, U., 112, 887

Menge, W.M.B.P. *see* Leurs, R., 112, 847

Mercuri, N.B., Bonci, A., Calabresi, P., Stratta, F., Stefani, A. and Bernardi, G. Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831

Merkel, L.A. *see* Cox, B.F., 111, 1233

Merlos, M. *see* Alegret, M., 112, 551

Mertès, P.-M. *see* Makki, T., 111, 555

Mervaala, E.M.A., Laakso, J., Vapaatalo, H. and Karppanen, H. Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

Mervaala, E.M.A., Paakkari, I., Laakso, J., Nevala, R., Teräväinen, T.M.-L., Fyhrquist, F., Vapaatalo, H. and Karppanen, H. Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Messori, E. *see* Tonini, M., 113, 1

Mettauer, B. *see* Monassier, L., 111, 1347

Meyer, W. *see* Kojda, G., 112, 533

Michael, J. *see* Wu, X.C., 112, 360

Michael, J. *see* Wu, X.C., 112, 369

Michel, M.C., Hanft, G. and Groß, G. Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539

Michel, M.C., Hanft, G. and Groß, G. Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533

Michel, M.C., Knapp, J. and Ratjen, H. Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240

Michel, M.C. and Kötting, A. Rapid desensitization of adrenaline- and neuropeptide Y-stimulated Ca^{2+} mobilization in HEL-cells, 112, 499

Mickel, S.J. *see* Waldmeier, P.C., 113, 1515

Middlemiss, D. *see* Balmforth, A.J., 112, 277

Middlemiss, D. *see* Hilditch, A., 111, 137

Mihara, S. *see* Cunningham, S.M.C., 113, 883

Milne, A. *see* Venkova, K., 112, 1237

Minami, T., Nishihara, I., Uda, R., Ito, S., Hyodo, M. and Hayaishi, O. Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735

Minc-Golomb, D., Tsarfaty, I. and Schwartz, J.P. Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720

Minck, K.-O. *see* Barber, A., 113, 1317

Minshall, E. *see* Spina, D., 112, 901

Mirrlees, D.J. *see* Taylor, P.D., 111, 42

Misra, A. *see* Watson, W.P., 112, 1017

Mistry, R. *see* Challiss, R.A.J., 112, 231

Mitchell, D. *see* Hall, J.M., 112, 985

Mitchell, J.A., Belvisi, M.G., Akarasereenont, P., Robbins, R.A., Kwon, O.-J., Croxtall, J., Barnes, P.J. and Vane, J.R. Induction of cyclooxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008

Mitchell, J.A. *see* Akarasereenont, P., 113, 1522

Mitchell, J.A. *see* Tomlinson, A., 113, 693

Miura, K. *see* Kim, S., 113, 662

Miura, M. *see* Yamauchi, H., 112, 958

Miyake, K., Takagi, N. and Takeo, S. Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226

Miyamoto, M.D. *see* Provan, S.D., 111, 1103

Mizuno, K. *see* Nakahata, N., 112, 299

Mizusawa, H. *see* Omote, M., 111, 233

Modesti, P.A., Cecioni, I., Colella, A., Costoli, A., Paniccia, R. and Serneri, G.G.N. Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

Moffat, M.P. *see* Khandoudi, N., 112, 1007

Molderings, G.J., Colling, E., Likungu, J., Jakschik, J. and Göthert, M. Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

Momose, K. *see* Shimizu, S.-i., 113, 564

Monassier, L., Tibiriça, E., Roegel, J.-C., Mettauer, B., Feldman, J. and Bousquet, P. Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347

Moncada, S. *see* Kiff, R.J., 113, 496

Moncada, S. *see* Laszlo, F., 111, 1309

Moncada, S. *see* László, F., 113, 1131
 Moncada, S. *see* Salas, E., 112, 1071
 Monge, L. *see* Fernández, N., 113, 550
 Monnet, F.P., Debonnel, G., Bergeron, R., Gronier, B. and de Montigny, C. The effects of sigma ligands and of neuropeptide Y on N-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709
 Moore, A.R. *see* Tomlinson, A., 113, 693
 Moore, K. *see* Hirokawa, K., 112, 396
 Moore, P.K. *see* Allawi, H.S., 113, 282
 Moors, J.A. *see* McAuliffe, S.J.G., 112, 272
 Morán, A., Velasco, C., Salvador, T., Martín, M.L. and San Román, L. Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358
 Morcillo, E.J. *see* Cortijo, J., 111, 769
 Moreau, R., Komeichi, H., Kirstetter, P., Yang, S., Aupetit-Faisant, B., Cailmail, S. and Lebrec, D. Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649
 Moreland, S., McMullen, D., Abbo-Offei, B. and Seymour, A. Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature, 112, 704
 Moreland, S. *see* Dickinson, K.E.J., 113, 179
 Morgan, N.G. *see* Chan, S.L.F., 112, 1065
 Morgan, P.F. *see* Durcan, M.J., 112, 855
 Morice, A.H. *see* Thompson, J.S., 113, 1121
 Morimoto, Y. *see* Yasumatsu, H., 111, 1170
 Moritoki, H., Hisayama, T., Takeuchi, S., Kondoh, W. and Imagawa, M. Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
 Moriyama, S. *see* Chiba, K., 112, 837
 Moro, M.A. *see* Salas, E., 112, 1071
 Morris, J.L. Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105
 Morrison, K.J. *see* Boulanger, C.M., 112, 519
 Morton, I.K.M. *see* Field, J.L., 113, 607
 Morton, I.K.M. *see* Hall, J.M., 112, 985
 Moshiri, A. *see* Schwanstecher, M., 113, 903
 Moskowitz, M.A. *see* Lee, W.S., 112, 920
 Moss, D.W. *see* Kiff, R.J., 113, 496
 Moussaoui, S.M. *see* Lee, W.S., 112, 920
 Mu, D.-X. *see* Lucchessi, B.R., 113, 1333
 Muddle, J.R. *see* Dashwood, M.R., 112, 386
 Mue, S. *see* Tanabe, J.-i., 113, 1480
 Mue, S. *see* Watanabe, M., 111, 123
 Mukai, J., Asai, T., Naka, M. and Tanaka T. Separation and characterization of a novel isoenzyme of cyclic nucleotide phospho-diesterase from rat cerebrum, 111, 389
 Müller-Ehmsen, J. *see* Böhm, M., 112, 137
 Mulvany, M.J. *see* Buus, N.H., 112, 579
 Munakata, M. *see* Nishikawa, M., 112, 1160
 Munday, I.T. *see* Lees, G., 113, 642
 Muraki, K., Bolton, T.B., Imaizumi, Y. and Watanabe, M. Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154
 Muramatsu, I. *see* Kohno, Y., 112, 1167
 Murphy, L.J., Corder, R., Mallet, A.I. and Turner, A.J. Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
 Murphy, O.E. *see* Baxter, G.S., 112, 323
 Murphy, T.V. *see* Seager, J.M., 111, 525
 Murray, K.J. *see* Turner, N.C., 111, 1047
 Murray, K.J. *see* Turner, N.C., 111, 1198
 Murray, R.D. *see* Barber, A., 111, 843
 Murray, R.D. *see* Barber, A., 113, 1317
 Mutafova-Yambolieva, V.N. and Radomirov, R.G. Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109
 Muto, S., Imai, M. and Asano, Y. Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na⁺ and K⁺ transport properties in the rabbit cortical collecting duct, 111, 173
 Mutschler, E. *see* McLaren, G.J., 111, 913
 Mutschler, E. *see* Waelbroeck, M., 112, 505
 Mutschler, E. *see* Windscheif, U., 113, 1015
 Mutschler, E. *see* Ziganshin, A.U., 111, 923
 Myers, J. *see* Seabrook, G.R., 111, 1061
 Myers, J. *see* Seabrook, G.R., 112, 728
 Mynett, K.J. *see* Bailey, C.J., 111, 793
 Mynett, K.J. *see* Bailey, C.J., 112, 671

N

Nabata, H. *see* Kamei, K., 113, 889
 Nadel, J.A. *see* Ricciardolo, F.L.M., 113, 508
 Nadel, J.A. *see* Ricciardolo, F.L.M., 113, 1147
 Naderali, E.K. and Poyser, N.L. The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103
 Nagant de Duxchaisnes, C. *see* Manicourt, D.H., 113, 1113
 Nagao, T. *see* Okuyama, R., 113, 451
 Nagy, I. *see* Szilvassy, Z., 112, 999
 Nahmias, C. *see* Blin, N., 112, 911
 Nahorski, S.R. *see* Challiss, R.A.J., 112, 231
 Naka, M. *see* Mukai, J., 111, 389
 Nakahata, N., Ishimoto, H., Mizuno, K., Ohizumi, Y. and Nakanishi, H. Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299
 Nakajima, N. *see* Yamauchi, H., 112, 958
 Nakane, T. *see* Ren, L.-M., 111, 1321
 Nakanishi, H. *see* Nakahata, N., 112, 299
 Nakao, T. *see* Yasumatsu, H., 111, 1170
 Nakatsu, K. *see* Hussain, A.S., 113, 1508
 Nakaya, Y. *see* Martin, D.K., 111, 873
 Nakazato, Y., Tani, Y., Teraoka, H., Sugawara, T., Asano, T., Ohta, T. and Ito, S. Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
 Nakazato, Y. *see* Ohta, T., 112, 972
 Nakazawa, K., Inoue, K., Koizumi, S., Ikeda, M. and Inoue, K. Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
 Nakazawa, K. *see* Koizumi, S., 112, 992
 Nally, J.E., Clayton, R.A., Thomson, N.C. and McGrath, J.C. The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328
 Nally, J.E., McCall, R., Young, L.C., Wakelam, M.J.O., Thomson, N.C. and McGrath, J.C. Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163
 Napoleone, P. *see* Lakhani, P.P., 111, 1296
 Naylor, A.M. *see* Kenny, B.A., 111, 1003
 Neal, M. and Cunningham, J. Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085
 Neal, M.J., Paterson, S.J. and Cunningham, J.R. Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789
 Negretti, N. *see* Donoso, P., 111, 455
 Nelemans, S.A. *see* Henning, R.H., 111, 459
 Neliat, G. *see* Galizzi, J.P., 113, 389
 Nevala, R. *see* Mervaala, E.M.A., 111, 1189
 Newman, M.J. *see* Wong, P.S.K., 112, 881
 Nichol, G.M., Parsons, G.H. and Chung, K.F. Effect of sodium metabolite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918
 Nicolarakis, P.J., Lin, Y.-Q. and Bennett, M.R. Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521
 Niederhoffer, N. *see* Makki, T., 111, 555
 Niederste-Hollenberg, A. *see* Edwards, G., 113, 1538
 Nijkamp, F.P. *see* Benschop, R.J., 113, 1311
 Nilius, B., Sehrer, J. and Droogmans, G. Permeation properties and

modulation of volume-activated Cl^- currents in human endothelial cells, 112, 1049

Nisato, D. *see* Roccon, A., 111, 145

Nishihara, I. *see* Minami, T., 112, 735

Nishikawa, M., Munakata, M. and Akaike, N. Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160

Nishikibe M. *see* Fukuroda, T., 113, 336

Nishimura, J. *see* Aoki, H., 111, 489

Nishimura, J. *see* Aoki, H., 111, 989

Noack, E. *see* Kojda, G., 112, 533

Noack, Th. *see* Edwards, G., 113, 1538

Nobile, M. *see* Maestrone, E., 111, 283

Noguera, M.A. *see* Chuliá, S., 113, 1377

Nomoto, M. *see* Shimizu, S.-i., 113, 564

Nomura, Y. *see* Charoonroje, P., 113, 569

Nörenberg, W., Langosch, J.M., Gebicke-Haerter, P.J. and Illes, P. Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942

Nörenberg, W. *see* Langosch, J.M., 113, 29

Norman, K.E., Argenbright, L.W., Williams, T.J. and Rossi, A.G. Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117

Norman, K.E. *see* Teixeira, M.M., 113, 1363

Nourshargh, S. *see* Sanz, M.J., 113, 513

Nuti, S. *see* Solito, E., 112, 347

O

Obrenovitch, T.P., Urenjak, J. and Zilkha, E. Intracerebral microdialysis combined with recording of extracellular field potential: a novel method for investigation of depolarizing drugs *in vivo*, 113, 1295

O'Connor, S.E. *see* Drieu la Rochelle, C., 111, 49

O'Donnell, L.J.D. *see* Battistini, B., 112, 1244

Ogawa, N. *see* Kashiwabara, T., 111, 278

Ogawa, S. *see* Katsuragi, T., 113, 795

Ohhashi, T. *see* Kawai, Y., 112, 635

Ohizumi, Y. *see* Furukawa, K.-I., 113, 233

Ohizumi, Y. *see* Nakahata, N., 112, 299

Ohkura, M. *see* Furukawa, K.-I., 113, 233

Ohta, K. *see* Kim, S., 113, 662

Ohta, T., Ito, S. and Nakazato, Y. All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972

Ohta, T. *see* Nakazato, Y., 111, 935

Ohtsuka, M. *see* Kita, Y., 113, 1137

Ohuchi, K. *see* Tanabe, J.-i., 113, 1480

Ohuchi, K. *see* Watanabe, M., 111, 123

Okamura, T. *see* Enokibori, M., 111, 77

Okiji, S.A. *see* Herd, C.M., 111, 7

Oku, Y. *see* Saito, K., 111, 465

Okuyama, R., Adachi-Akahane, S. and Nagao, T. Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca^{2+} current in guinea-pig heart, 113, 451

Olianas, M.C. *see* Onali, P., 113, 775

Oliva, M.L.V. *see* Lopes-Martins, R.A.B., 113, 81

Olmos, G., Alemany, R., Escriba, P.V. and García-Sevilla, J.A. The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997

Omote, M. and Mizusawa, H. Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233

Omura, T. *see* Kim, S., 113, 662

Onali, P., Aasen, A.J. and Olianas, M.C. Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775

O'Neill, S.C. *see* Donoso, P., 111, 455

Ong, R. *see* Yang, C.M., 111, 21

Ong, R. *see* Yang, C.M., 111, 777

Ong, R. *see* Yang, C.M., 112, 781

Ongini, E. *see* Dionisotti, S., 112, 659

Oon, B.B. *see* Taylor, P.D., 111, 35

Opgenorth, T.J. *see* Pollock, D.M., 111, 729

O'Reilly, G. *see* Maguire, J.J., 113, 49

O'Reilly, S. *see* Bolser, D.C., 113, 1344

Oriowo, M.A. Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699

Orrman-Rossiter, S.L. *see* Johnson, P.J., 111, 1240

O'Shaughnessy, K. *see* Hirokawa, K., 112, 396

O'Shaughnessy, K.M. *see* Hirokawa, K., 113, 1448

Oshima, Y. *see* Furukawa, K.-I., 113, 233

Osuka, M. *see* Hosoki, R., 113, 1141

Osuka, M. *see* Suzuki, H., 113, 310

Otter, D.J. and Chess-Williams, R. The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576

Ouedraogo, S., Stoclet, J.-C. and Bucher, B. Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625

Ouedraogo, S., Tschöpl, M., Stoclet, J.-C. and Bucher, B. Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867

Owen, D. *see* Hall, A., 113, 959

Oxford, A.W. *see* Gale, J.D., 111, 332

Ozaki, R. *see* Kita, Y., 113, 1137

Ozaki, S. *see* Fukuroda, T., 113, 336

Ozanne, S.E. *see* Lee, K., 111, 632

P

Paakkari, I. *see* Mervaala, E.M.A., 111, 1189

Pacaud, P., Grégoire, G. and Loirand, G. Release of Ca^{2+} from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca^{2+} entry, 113, 457

Page, C.P. *see* Herd, C.M., 111, 7

Page, C.P. *see* Herd, C.M., 112, 292

Page, C.P. *see* Paul, W., 111, 159

Page, C.P. *see* Spina, D., 112, 901

Page, T. *see* Bailey, C.J., 112, 671

Paiava, A.C.M. *see* Silva, E.G., 113, 1022

Paiava, T.B. *see* Silva, E.G., 113, 1022

Pak, J.-M., Adeagbo, A.S.O., Triggle, C.R., Shaffer, E.A. and Lee, S.S. Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209

Palmi, M., Frosini, M., Becherucci, C., Sgaragli, G.P. and Parente, L. Increase of extracellular brain calcium involved in interleukin-18-induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449

Pang, C.C.Y. *see* Elhawary, A.M., 111, 819

Paniccia, R. *see* Modesti, P.A., 112, 81

Pannangpetch, P. *see* Woodman, O.L., 112, 443

Panten, U. *see* Schwanstecher, C., 111, 302

Panten, U. *see* Schwanstecher, M., 113, 903

Papp, J.G. *see* Vegh, A., 113, 1081

Papp, J.G. *see* Vegh, A., 113, 1167

Parente, L. *see* Palmi, M., 112, 449

Parente, L. *see* Solito, E., 112, 347

Parkington, H.C. *see* Tonta, M.A., 112, 525

Parratt, J. *see* Vegh, A., 113, 1167

Parratt, J.R. *see* Song, W., 113, 1083

Parratt, J.R. *see* Vegh, A., 113, 1081

Parsons, G.H. *see* Nichol, G.M., 111, 918

Parsons, S.J.W., Hill, A., Waldron, G.J., Plane, F. and Garland, C.J. The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275

Pasquale, C. *see* Pires, A.L.A., 113, 994

Passani, M.B., Pugliese, A.M., Azzurrini, M. and Corradetti, R. Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695

Patacchini, R. *see* Belvisi, M.G., 111, 103

Patacchini, R. *see* Maggi, C.A., 112, 150

Patacchini, R. *see* Maggi, C.A., 112, 161

Patacchini, R. *see* Meini, S., 111, 739

Patel, M., Sheehan, M.J. and Strong, P. Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741

Patel, S. *see* Delamere, F., 111, 983

Patel, S. *see* Rupniak, N.M.J., 113, 1487

Patel, S. *see* Seabrook, G.R., 111, 391

Patel, S. *see* Seabrook, G.R., 111, 1061

Patel, S. *see* Seabrook, G.R., 112, 728

Paterson, I.A. *see* Berry, M.D., 113, 1159

Paterson, I.A. *see* Zhu, M.-Y., 112, 23

Paterson, S.J. *see* Neal, M.J., 113, 789

Patoiseau, J.-F. *see* Le Grand, B., 112, 195
 Paul, W., Douglas, G.J., Lawrence, L., Khawaja, A.M., Perez, A.C., Schachter, M. and Page, C.P. Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159
 Pavenstt, H., Ruh, J., Greger, R. and Schollmeyer, P. Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7
 Pavord, I. *see* Delamere, F., 111, 983
 Payvandi, N. *see* Shaunak, S., 113, 151
 Peachey, J.A., Hourani, S.M.O. and Kitchen, I. The binding of 1,3-[³H]-dipropyl-8-cyclopentylxanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249
 Peakman, M.-C. and Hill, S.J. Adenosine A_{2B}-receptor-mediated cyclic AMP accumulation in primary rat astrocytes, 111, 191
 Peakman, M.-C. and Hill, S.J. Endogenous expression of histamine H₁ receptors functionally coupled to phosphoinositide hydrolysis in C6 glioma cells: regulation by cyclic AMP, 113, 1554
 Pearson, J.D. *see* Baydoun, A.R., 112, 487
 Peers, C. *see* McDonald, R.L., 113, 621
 Peglion, J.L. *see* Thollon, C., 112, 37
 Pelekmans, P.A. *see* De Man, J.G., 112, 341
 Pelln, M.I., Steil, A.A., Furi, V. and Crespo, M.S. Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101
 Penner, S.B. *see* Li, P., 112, 200
 Penner, S.B. and Smyth, D.D. Sodium excretion following central administration of an I₁ imidazoline receptor agonist, moxonidine, 112, 1089
 Peoples, R.W. and Weight, F.F. Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555
 Pepperl, D.J. *see* Regan, J.W., 112, 377
 Perez, A.C. *see* Paul, W., 111, 159
 Prez, O., Valenzuela, C., Delpn, E. and Tamargo, J. Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717
 Prez-Vizcao, F., del Pozo, B.F., Zaragoz, F. and Tamargo, J. Voltage- and time-dependent inhibitory effects on rat aortic and porcine coronary artery contraction induced by propafenone and quinidine, 113, 1281
 Peristeris, P. *see* Pipili-Synetos, E., 111, 894
 Perkins, M.N. *see* Cruwys, S.C., 113, 940
 Perkins, M.N. *see* Davis, A.J., 113, 63
 Perretti, M., Harris, J.G. and Flower, R.J. A role for endogenous histamine in interleukin-8-induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801
 Perrone, M.H. *see* Cox, B.F., 111, 1233
 Persson, M.G., Agvald, P. and Gustafsson, L.E. Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825
 Pertwee, R.G., Stevenson, L.A. and Griffin, G. Erratum, 111, 968
 Pesonen, U. *see* Santti, E., 113, 1231
 Pessina, A.C. *see* Bova, S., 111, 1067
 Pestana, M. and Soares-da-Silva, P. Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269
 Pestana, M. and Soares-da-Silva, P. The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417
 Peth, G., Joci, M. and Holzer, P. Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
 Pettipher, E.R., Salter, E.D. and Showell, H.J. Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
 Pfeilschifter, J. *see* Huwiler, A., 113, 1455
 Pfeilschifter, J. *see* Kunz, H.M.D., 112, 1
 Picard, P., Regoli, D. and Couture, R. Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240
 Picard, P. *see* Prat, A., 112, 250
 Piekarska, A.E. *see* McPherson, G.A., 112, 1223
 Pierce, K.L. *see* Regan, J.W., 112, 377
 Pieretti, S., Di Giannuario, A., Domenici, M.R., Sagratella, S., Capasso, A., Sorrentino, L. and Loizzo, A. Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents, 113, 1416
 Pinnock, R.D. *see* Boden, P.R., 112, 666
 Pinto, F. *see* Hernandez, J., 111, 1316
 Pinto-do-, P.C. *see* Soares-da-Silva, P., 112, 611
 Pipili-Synetos, E., Sakkoula, E., Haralabopoulos, G., Andriopoulou, P., Peristeris, P. and Maragoudakis, M.E. Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
 Pires, A.L.A., e Silva, P.M.R., Pasquale, C., Castro-Faria-Neto, H.C., Bozza, P.T., Cordeiro, R.S.B., Rae, G.A., Braquet, P., Lagente, V. and Martins, M.A. Long-lasting inhibitory activity of the hexapeptide BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994
 Pitcher, A. *see* Allawi, H.S., 113, 282
 Plane, F. *see* Parsons, S.J.W., 113, 1275
 Planelles, G. *see* Belachgar, F., 112, 465
 Plante, G.E. *see* Sirois, M.G., 111, 1111
 Plevin, R., Kellock, N.A., Wakelam, M.J.O. and Wadsworth, R. Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311
 Plumpton, C. *see* Tan, K.K.C., 111, 703
 Poklewsko-Koziell, M. *see* Hussain, A.S., 113, 1508
 Polak, J.M. *see* Desai, K.M., 113, 1197
 Polak, J.M. *see* Rutherford, R.A.D., 113, 931
 Polak, J.M. *see* Walsh, D.A., 112, 435
 Polak, J.M. *see* Walsh, D.A., 113, 1407
 Polley, J. *see* Hilditch, A., 111, 137
 Pollock, D.M. and Opgenorth, T.J. ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729
 Pook, P.C.-K., *see* Jane, D.E., 112, 809
 Poronnik, P. *see* Gibb, C.A., 111, 1135
 Prsti, I., Bara, A.T., Busse, R. and Hecker, M. Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
 Prsti, I. *see* Hecker, M., 111, 238
 Prsti, I. *see* Khnen, M., 112, 925
 Prsti, I. *see* Vuorinen, P., 112, 71
 Poss, M.A. *see* Dickinson, K.E.J., 113, 179
 Poston, L. *see* Taylor, P.D., 111, 35
 Poston, L. *see* Taylor, P.D., 111, 42
 Poston, L. *see* Taylor, P.D., 113, 801
 Potier, E. *see* Maggi, C.A., 112, 150
 Potter, E.K. *see* Lacroix, J.S., 113, 479
 Poubelle, P.E. *see* Marleau, S., 112, 654
 Powell, J.R. *see* Dickinson, K.E.J., 113, 179
 Powell, T. *see* Kozlowski, R.Z., 111, 250
 Poyer, N.L. *see* Naderali, E.K., 113, 103
 Prashad, T.R. *see* Low, A.M., 113, 1263
 Prat, A., Picard, P. and Couture, R. Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250
 Pregenzer, J.F. *see* Im, W.B., 112, 1025
 Prmont, J. *see* Hubert, J.P., 113, 261
 Pretolani, M. *see* Zuany-Amorim, C., 112, 1230
 Preziosi, P. *see* Curr, D., 113, 541
 Prinssen, E.P.M., Ellenbroek, B.A. and Cools, A.R. Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769
 Pritchard, M.C. *see* Boden, P.R., 112, 666
 Prost, J.-F. *see* Thollon, C., 112, 37
 Provan, S.D. and Miyamoto, M.D. Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103
 Pruneau, D., Luccarini, J.M., Robert, C. and Blichard, P. Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029
 Pruneau, D. *see* Deckert, V., 112, 939
 Pugliese, A.M. *see* Passani, M.B., 112, 695
 Pun, K.-T. *see* Balmforth, A.J., 112, 277
 Purkiss, J.R., West, D., Wilkes, L.C., Scott, C., Yarrow, P., Wilkinson, G.F. and Boarder, M.R. Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoreceptor agonists, 111, 1041

Purkiss, J.R., Wilkinson, G.F. and Boarder, M.R. Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y} -purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
 Puttick, R.M. *see* Takahashi, H., 111, 1147
 Pyne, N.J. *see* Burns, F., 113, 3

Q

Qian, Y.-m., Jones, R.L., Chan, K.-m., Stock, A.I. and Ho, J.K.S. Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369
 Quartara, L. *see* Maggi, C.A., 112, 150
 Quik, M. *see* Clarke, P.B.S., 111, 397
 Quilley, J. *see* Fulton, D., 113, 954

R

Rabbani, M., Wright, J., Butterworth, A.R., Zhou, Q. and Little, H.J. Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89
 Racké, K. *see* Wessler, I., 113, 1221
 Rademaker, B. *see* Jansen, F.P., 113, 355
 Radomirov, R.G. *see* Mutafova-Yambolieva, V.N., 112, 1109
 Radomski, M.W. *see* Salas, E., 112, 1071
 Rae, G.A. *see* Pires, A.L.A., 113, 994
 Raeburn, D., Underwood, S.L., Lewis, S.A., Woodman, V.R., Battram, C.H., Tomkinson, A., Sharma, S., Jordan, R., Souness, J.E., Webber, S.E. and Karlsson, J.-A. Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423
 Raeburn, D. *see* Souness, J.E., 111, 1081
 Rafeiner, K. *see* Waelbroeck, M., 112, 505
 Ragazzoni, E. *see* Currò, D., 113, 541
 Rakhit, T. *see* Gardiner, S.M., 111, 589
 Ralevic, V. *see* Windscheif, U., 113, 1015
 Ramage, A.G. *see* Shepheard, S.L., 111, 616
 Ramírez, M.J., Cenarruzabeitia, E., Del Río, J. and Lasheras, B. Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
 Ramírez, M.J., Del Río, J., Cenarruzabeitia, E. and Lasheras, B. On the nature of the 5-HT receptor subtype inhibition acetylcholine release in the guinea-pig ileum, 113, 77
 Ramnarine, S.I., Hirayama, Y., Barnes, P.J. and Rogers, D.F. 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183
 Ramrakha, P. *see* Hirokawa, K., 112, 396
 Ramrakha, P. *see* Hirokawa, K., 113, 1448
 Rand, M.J. and Li, C.G. Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus, 111, 1089
 Randall, M.D., Ujiiie, H. and Griffith, T.M. Modulation of vasodilatation to levocromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
 Ratcliff, R. *see* Cuthbert, A.W., 112, 31
 Ratjen, H. *see* Michel, M.C., 113, 240
 Raung, S.L. *see* Wang, J.P., 113, 945
 Razzaque, Z. *see* Longmore, J., 112, 176
 Read, A.M. *see* Kenny, B.A., 111, 1003
 Redman, R.S. *see* Hunt, J.M., 111, 753
 Redman, R.S. and Silinsky, E.M. Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375
 Regan, J.W., Bailey, T.J., Donello, J.E., Pierce, K.L., Pepperl, D.J., Zhang, D., Kedzie, K.M., Fairbairn, C.E., Bogardus, A.M., Woodward, D.F. and Gil, D.W. Molecular cloning expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377
 Regoli, D. *see* Picard, P., 112, 240
 Reichart, B. *see* Böhm, M., 112, 137
 Reichenspurner, H. *see* Böhm, M., 112, 137
 Reidemeister, S. *see* Schlicker, E., 112, 1043
 Reidemeister, S. *see* Schlicker, E., 113, 657
 Reinheimer, T. *see* Wessler, I., 113, 1221
 Ren, L.-M., Nakane, T. and Chiba, S. Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by

electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
 Renaud, N. *see* Su, J., 113, 395
 Reuben, M. *see* Clarke, P.B.S., 111, 695
 Reynia, S. *see* Teixeira, M.M., 111, 811
 Ribeiro, J.A. *see* Correia-de-Sá, P., 111, 582
 Ribeiro, J.A. *see* Hernandez, J., 111, 1316
 Ricciardolo, F.L.M., Nadel, J.A., Graf, P.D., Bertrand, C., Yoshihara, S. and Geppetti, P. Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
 Ricciardolo, F.L.M., Nadel, J.A., Yoishihara, S. and Geppetti, P. Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
 Richard, V., Kaeffer, N., Hogie, M., Tron, C., Blanc, T. and Thuillez, C. Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869
 Richards, N.T. *see* Wu, X.C., 112, 360
 Richards, N.T. *see* Wu, X.C., 112, 369
 Richardson, D.P. *see* Einstein, R., 111, 199
 Richardson, P.J. *see* Kurokawa, M., 113, 43
 Riezebos, J., Watts, I.S. and Vallance, P.J.T. Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
 Ritter, J.M. *see* Edwards, J.S., 113, 926
 Riva, L. *see* Briand, V., 112, 1195
 Rizzi, C.A. *see* Tonini, M., 113, 1
 Robbins, R.A. *see* Mitchell, J.A., 113, 1008
 Robert, C. *see* Pruneau, D., 111, 1029
 Robertson, M.J. *see* Hilditch, A., 111, 137
 Robichaud, A. *see* Chamelian, M., 113, 317
 Robinson, M. *see* Teixeira, M.M., 111, 811
 Roccon, A., Marchionni, D., Donat, F., Segondy, D., Cazaubon, C. and Nisato, D. A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145
 Rodrigues de Miranda, J.F. *see* Waelbroeck, M., 112, 505
 Roegel, J.-C. *see* Monassier, L., 111, 1347
 Rogers, D.F. *see* Ramnarine, S.I., 113, 1183
 Rola-Pleszczynski, M. *see* Claeys, A., 112, 1202
 Roller, P.P. *see* Abassi, Z.A., 113, 204
 Roques, B.P. *see* Derrien, M., 111, 956
 Roques, T.W. *see* Gibson, W.J., 111, 1262
 Ross, B.C. *see* Hilditch, A., 111, 137
 Rossi, A.G. *see* Norman, K.E., 111, 117
 Rossi, A.G. *see* Teixeira, M.M., 111, 811
 Rossi, A.G. *see* Teixeira, M.M., 112, 332
 Rossi, A.G. *see* Teixeira, M.M., 113, 1363
 Rossi, G. *see* Bova, S., 111, 1067
 Rostène, W. *see* Dubuc, I., 112, 352
 Rote, W.E. *see* Lucchessi, B.R., 113, 1333
 Rothwell, L.A. *see* Bowery, B., 112, 873
 Rouru, J. *see* Santti, E., 113, 1231
 Rouse, W. and Johnson, I.R. Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
 Rowan, M.J. *see* Manahan-Vaughan, D., 112, 1083
 Rowbotham, D.J. *see* Atcheson, R., 111, 787
 Rowley, M. *see* Tricklebank, M.D., 113, 729
 Rubin, P.C. *see* Kemp, P.A., 111, 325
 Rubino, A. and Burnstock, G. Recovery after dietary vitamin E supplementation of impaired endothelial function in vitamin E-deficient rats, 112, 515
 Rügheimer, E. *see* Tschaikowsky, K., 113, 664
 Ruh, J. *see* Pavenstädt, H., 113, 7
 Rump, A.F.E., Acar, D. and Klaus, W. A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757
 Rump, L.C. and von Kügelgen, I. A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
 Rundfeldt, C. *see* Löscher, W., 112, 97
 Rupniak, N.M.J., Patel, S., Marwood, R., Webb, J., Traynor, J.R., Elliott, J., Freedman, S.B., Fletcher, S.R. and Hill, R.G. Anti-

nociceptive and toxic effects of (+)-epibatidine oxalate attributable to nicotinic agonist activity, 113, 1487

Russell, M.J. *see* Kenny, B.A., 111, 1003

Rutherford, R.A.D., Matsuda, Y., Wilkins, M.R., Polak, J.M. and Wharton, J. Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1, 113, 931

Ryon, D.E. *see* Dickinson, K.E.J., 113, 179

S

Safar, M. *see* Bouaziz, H., 113, 717

Sagrada, A., Schiavi, G.B., Cereda, E. and Ladinsky, H. Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

Sagratella, S. *see* Pieretti, S., 113, 1416

Saieddine, M. *see* Laniyonu, A., 113, 403

St. J. Jones, P.L. *see* Jane, D.E., 112, 809

St Lambert, J.H., Dawid-Milner, M.S., Silva-Carvalho, L. and Spyer, K.M. Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

Saito, H. *see* Kohno, Y., 112, 1167

Saito, H. *see* Torii, Y., 111, 431

Saito, K., Suetsugu, T., Oku, Y., Kuroda, A. and Tanaka, H. α_1 -Adrenoceptors in the conduction system of rat hearts, 111, 465

Saito, K. *see* Suzuki, H., 113, 310

Saitoh, T., Ishida, M., Maruyama, M. and Shinozaki, H. A novel antagonist, phenylbenzene ω -phosphono- α -amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord, 113, 165

Saitta, A. *see* Squadrito, F., 113, 912

Saitta, A. *see* Squadrito, F., 113, 1153

Sakai, S. *see* Kita, Y., 113, 1137

Sakkoula, E. *see* Pipili-Synetos, E., 111, 894

Salas, C. *see* Donoso, M.V., 111, 473

Salas, E., Moro, M.A., Askew, S., Hodson, H.F., Butler, A.R., Radomski, M.W. and Moncada, S. Comparative pharmacology of analogues of S-nitroso-N-acetyl-DL-penicillamine on human plate-lets, 112, 1071

Salmon, M. *see* Walsh, D.A., 113, 1407

Salter, E.D. *see* Pettipher, E.R., 113, 117

Salvador, T. *see* Morán, A., 113, 1358

Sampaio, C.A.M. *see* Lopes-Martins, R.A.B., 113, 81

San Román, L. *see* Morán, A., 113, 1358

Sandoli, D. *see* Dionisotti, S., 112, 659

Sanger, G.J. *see* Wardle, K.A., 112, 789

Santagostino-Barbone, M.G. *see* Lucchelli, A., 112, 763

Santiago, M., Machado, A. and Cano, J. Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837

Santicioli, P. and Maggi, C.A. Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588

Santicioli, P. *see* Ahluwalia, A., 111, 1017

Santicioli, P. *see* Maggi, C.A., 111, 687

Santicioli, P. *see* Maggi, C.A., 112, 161

Santti, E., Huupponen, R., Rouru, J., Hänninen, V., Pesonen, U., Jhanwar-Uniyal, M. and Koulu, M. Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

Sanz, C.M. *see* Cortijo, J., 111, 769

Sanz, M.J., Weg, V.B., Walsh, D.T., Williams, T.J. and Nourshargh, S. Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

Sato, K. *see* Dohi, Y., 111, 673

Sato, Y., Shibanoki, S., Sugahara, M. and Ishikawa, K. Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625

Savineau, J.-P. and Marthan, R. Diosmin-induced increase in sensitivity to Ca²⁺ of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Saxena, P.R. *see* Bax, W.A., 113, 1471

Saxena, P.R. *see* van Gelderen, E.M., 111, 961

Saywell, K. *see* Tricklebank, M.D., 113, 729

Scadding, G.K. *see* Austin, C.E., 111, 969

Scamps, F. and Vassort, G. Pharmacological profile of the ATP-mediated increase in L-type calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982

Scarpello, K.E. *see* Chan, S.L.F., 112, 1065

Scarr, E. *see* Berry, M.D., 113, 1159

Scatton, B. *see* Bonnafous, C., 113, 268

Schachter, M. *see* Paul, W., 111, 159

Schambye, H.T., Wijk, B.v., Hjorth, S.A., Wienen, W., Entzeroth, M., Bergsma, D.J. and Schwartz, T.W. Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Schiavi, G.B. *see* Sagrada, A., 113, 711

Schlemper, V. and Calixto, J.B. Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83

Schlicker, E., Kathmann, M., Reidemeister, S., Stark, H. and Schunack, W. Erratum, 113, 657

Schlicker, E., Kathmann, M., Reidemeister, S., Stark, H. and Schunack, W. Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043

Schneider, J. *see* Edwards, G., 113, 1538

Scholfield, C.N. *see* Yassin, M., 111, 1221

Schollmeyer, P. *see* Pavestadt, H., 113, 7

Schönhuber, F. *see* Tschaikowsky, K., 113, 664

Schrör, K. *see* Woditsch, I., 111, 1035

Schulz, Ch. *see* Hauck, R.W., 113, 1043

Schunack, W. *see* Schlicker, E., 112, 1043

Schunack, W. *see* Schlicker, E., 113, 657

Schutte, P.J. *see* Du Plooy, W.J., 111, 73

Schwanstecher, C., Dickel, C. and Panten, U. Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K⁺ channel in mouse B-cells, 111, 302

Schwanstecher, C. *see* Schwanstecher, M., 113, 903

Schwanstecher, M., Schwanstecher, C., Dickel, C., Chudziak, F., Moshiri, A. and Panten, U. Location of the sulphonylurea receptor at the cytoplasmic face of the B-cell membrane, 113, 903

Schwartz, J.P. *see* Minc-Golomb, D., 112, 720

Schwartz, T.W. *see* Schambye, H.T., 113, 331

Schwarzer, C. *see* Greber, S., 113, 737

Schwinger, R.H.G. *see* Böhm, M., 112, 137

Scott, C. *see* Purkiss, J.R., 111, 1041

Scott, G., Luscombe, G.P. and Mason, R. The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97

Scott, L.C. *see* Souness, J.E., 111, 1081

Scott, R.H. *see* Stapleton, S.R., 111, 57

Seabrook, G.R., Kemp, J.A., Freedman, S.B., Patel, S., Sinclair, H.A. and McAllister, G. Functional expression of human D₃ dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391

Seabrook, G.R., Knowles, M., Brown, N., Myers, J., Sinclair, H., Patel, S., Freedman, S.B. and McAllister, G. Pharmacology of high-threshold calcium currents in GH₄C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728

Seabrook, G.R., McAllister, G., Knowles, M.R., Myers, J., Sinclair, H., Patel, S., Freedman, S.B. and Kemp, J.A. Depression of high-threshold calcium currents by activation of human D₂ (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

Seabrook, G.R. *see* Bowery, B., 112, 873

Seabrook, G.R. *see* Brown, N.A., 113, 600

Seager, J.M., Murphy, T.V. and Garland, C.J. Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525

Seale, P.W. *see* Cuthbert, A.W., 112, 31

Seaton, E.D. *see* Matthews, E.K., 113, 87

Segondy, D. *see* Roccon, A., 111, 145

Sehrer, J. *see* Nilius, B., 112, 1049

Self, G.J. *see* Goldie, R.G., 112, 749

Sellers, A.J. and Ashford, M.L.J. Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619, 113, 659

Senoh, H. *see* Kita, Y., 113, 1137

Sepúlveda, G. *see* Donoso, M.V., 111, 473
 Serafino, R.P. *see* Dickinson, K.E.J., 113, 179
 Serner, G.G.N. *see* Modesti, P.A., 112, 81
 Serup, J. *see* Gniadecki, R., 113, 439
 Setoguchi, M. *see* Yasumatsu, H., 111, 1170
 Seyfried, C.A. *see* Barber, A., 111, 843
 Seyfried, C.A. *see* Barber, A., 113, 1317
 Seymour, A. *see* Moreland, S., 112, 704
 Sgaragli, G.P. *see* Palmi, M., 112, 449
 Shaffer, E.A. *see* Pak, J.-M., 112, 1209
 Shafiq, J. *see* Itoh, T., 111, 165
 Shankley, N.P. *see* Welsh, N.J., 112, 93
 Sharma, S. *see* Raeburn, D., 113, 1423
 Shaunak, S., Gooderham, N.J., Edwards, R.J., Payvandi, N., Javan, C.M., Baggett, N., MacDermot, J., Weber, J.N. and Davies, D.S. Infection by HIV-1 blocked by binding of dextrin 2-sulphate to the cell surface of activated human peripheral blood mononuclear cells and cultured T-cells, 113, 151
 Shaw, D. *see* Longmore, J., 112, 176
 Sheedy, W. *see* Thompson, J.S., 113, 1121
 Sheehan, M.J. *see* Anderson, R., 113, 1386
 Sheehan, M.J. *see* Patel, M., 113, 741
 Shen, W.-X., Jobling, P. and Horn, J.P. The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898
 Shepheard, S.L., Jordan, D. and Ramage, A.G. Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
 Shibano, T. *see* Chiba, K., 112, 837
 Shibanoiki, S. *see* Sato, Y., 112, 625
 Shimizu, S.-i., Nomoto, M., Yamamoto, T. and Momose, K. Reduction by N^G -nitro-L-arginine of H_2O_2 -induced endothelial cell injury, 113, 564
 Shinozaki, H. *see* Cartmell, J., 111, 364
 Shinozaki, H. *see* Saitoh, T., 113, 165
 Shinozuka, K., Hashimoto, M., Masumura, S., Bjur, R.A., Westfall, D.P. and Hattori, K. *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
 Shirahase, H. *see* Jino, H., 111, 435
 Shirato, K. *see* Yamauchi, H., 112, 958
 Shoaib, M., Benwell, M.E.M., Akbar, M.T., Stolerman, I.P. and Balfour, D.J.K. Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073
 Shock, A. *see* Teixeira, M.M., 111, 811
 Shoupe, T.S. *see* Herd, C.M., 111, 7
 Shoupe, T.S. *see* Herd, C.M., 112, 292
 Showell, H.J. *see* Pettipher, E.R., 113, 117
 Shuker, M.A., Bowser-Riley, F. and Davies, S.N. Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951
 Shuto, K. *see* Kitazawa, T., 111, 288
 Siarey, R.J., Long, S.K., Tulp, M.T.H. and Evans, R.H. The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497
 Silinsky, E.M. *see* Hunt, J.M., 111, 753
 Silinsky, E.M. *see* Redman, R.S., 113, 375
 Silke, B. *see* Herity, N.A., 112, 423
 Sillence, M.N. and Matthews, M.L. Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866
 Silva, E.G., Frediani-Neto, E., Ferreira, A.T., Paiva, A.C.M. and Paiva, T.B. Role of Ca^{2+} -dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022
 Silva-Carvalho, L. *see* St Lambert, J.H., 113, 159
 Simmons, N.L. *see* Thwaites, D.T., 113, 1050
 Simon, M. *see* Barber, A., 111, 843
 Simon, M. *see* Barber, A., 113, 1317
 Sinclair, H. *see* Seabrook, G.R., 111, 1061
 Sinclair, H. *see* Seabrook, G.R., 112, 728
 Sinclair, H.A. *see* Seabrook, G.R., 111, 391
 Singh, L. *see* Hunter, J.C., 112, 1002
 Singh, L. *see* Tricklebank, M.D., 113, 729
 Singh, S. *see* Gibb, C.A., 111, 1135
 Sirois, M.G., de Lima, W.T., de Brum Fernandes, A.J., Johnson, R.J., Plante, G.E. and Sirois, P. Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
 Sirois, P. *see* Claing, A., 112, 1202
 Sirois, P. *see* Sirois, M.G., 111, 1111
 Sisto, A. *see* Maggi, C.A., 112, 150
 Sjöberg, J. *see* Källström, B.-L., 113, 687
 Sjödin, L. and Gylfe, E. Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037
 Skwisch, S. *see* Dickinson, K.E.J., 113, 179
 Small, R.C. *see* Cortijo, J., 111, 769
 Smart, T.G. *see* Bowie, D., 111, 803
 Smit, J. *see* Coppes, R.P., 113, 342
 Smit, M.J. *see* Leurs, R., 112, 847
 Smith, G.C.S. and McGrath, J.C. Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
 Smith, J.A. *see* Hutcheson, I.R., 113, 190
 Smith, P.A., Williams, B.A. and Ashcroft, F.M. Block of ATP-sensitive K^+ channels in isolated mouse pancreatic β -cells by 2,3-butanedione monoxime, 112, 143
 Smith, P.A. *see* Chen, H., 113, 55
 Smith, P.A. *see* Ellory, J.C., 111, 903
 Smith, P.F. *see* Gilchrist, D.P.D., 111, 358
 Smits, J.F.M. *see* Drieman, J.C., 111, 1117
 Smyth, D.D. *see* Li, P., 112, 200
 Smyth, D.D. *see* Penner, S.B., 112, 1089
 Snarey, M. *see* Baldwin, H.A., 112, 188
 Sneddon, P. *see* McLaren, G.J., 111, 913
 Snook, M.B. *see* Wiley, J.S., 112, 946
 Soares-da-Silva, P., Fernandes, M.H. and Pinto-do-Ó, P.C. Cell inward transport of L-DOPA and 3-O-methyl-L-DOPA in rat renal tubules, 112, 611
 Soares-da-Silva, P. *see* Pestana, M., 112, 417
 Soares-da-Silva, P. *see* Pestana, M., 113, 1269
 Sobey, C.G. *see* Loke, K.E., 112, 630
 Soediono, P. and Burnstock, G. Contribution of ATP and nitric oxide to NANC inhibitory transmission in rat pyloric sphincter, 113, 681
 Solito, E., Nuti, S. and Parente, L. Dexamethasone-induced translocation of lipocortin (annexin) I to the cell membrane of U-937 cells, 112, 347
 Solsona, C. *see* Cantí, C., 112, 19
 Song, W., Furman, B.L. and Parratt, J.R. Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083
 Soo, S.-C. *see* Bogle, R.G., 111, 1257
 Sorenson, R. *see* Hall, A., 113, 959
 Sorrentino, L. *see* Pieretti, S., 113, 1416
 Soubrié, P. *see* Dubuc, I., 112, 352
 Souness, J.E., Villamil, M.E., Scott, L.C., Tomkinson, A., Giembycz, M.A. and Raeburn, D. Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
 Souness, J.E. *see* Raeburn, D., 113, 1423
 Southan, G.J. *see* Szabó, C., 112, 355
 Southan, G.J. *see* Szabó, C., 113, 757
 Späth, L. *see* von Kügelgen, I., 113, 815
 Sperk, G. *see* Greber, S., 113, 737
 Spina, D., Minshall, E., Goldie, R.G. and Page, C.P. The effect of allosteric antagonists in modulating muscarinic M_2 -receptor function in guinea-pig isolated trachea, 112, 901
 Spina, D. and Goldie, R.G. Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29
 Spina, D. and Goldie, R.G. Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895
 Spinelli, W. *see* de Lorenzi, F.G., 113, 527
 Spyer, K.M. *see* St Lambert, J.H., 113, 159
 Squadrito, F., Altavilla, D., Canale, P., Ioculano, M., Campo, G.M., Ammendolia, L., Ferlito, M., Zingarelli, B., Squadrito, G., Saitta, A. and Caputi, A.P. Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery

occlusion shock, 113, 1153

Squadrito, F., Altavilla, D., Canale, P., Ioculano, M., Campo, G.M., Ammendolia, L., Squadrito, G., Saitta, A., Calapai, G. and Caputi, A.P. Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912

Squadrito, F. *see* Calapai, G., 111, 1328

Squadrito, G. *see* Squadrito, F., 113, 912

Squadrito, G. *see* Squadrito, F., 113, 1153

Stapleton, S.R., Scott, R.H. and Bell, B.A. Effects of metabolic blockers on Ca^{2+} -dependent currents in cultured sensory neurones from neonatal rats, 111, 57

Stark, H. *see* Schlicker, E., 112, 1043

Stark, H. *see* Schlicker, E., 113, 657

Starke, K. *see* Bültmann, R., 111, 151

Starke, K. *see* Bültmann, R., 112, 690

Starke, K. *see* Bültmann, R., 113, 349

Starke, K. *see* von Kügelgen, I., 113, 815

Stefani, A. *see* Mercuri, N.B., 113, 831

Steil, A.A. *see* Pellón, M.I., 112, 1101

Steinbusch, H.W.M. *see* Jansen, F.P., 113, 355

Sterin-Borda, L. *see* Bacman, S., 111, 271

Stevens, P.A. *see* Burns, F., 113, 3

Stevenson, L.A. *see* Pertwee, R.G., 111, 968

Stewart, A.G. *see* Tomlinson, P.R., 111, 641

Stock, A.I. *see* Qian, Y.-m., 113, 369

Stoclet, J.-C. *see* Ouedraogo, S., 111, 625

Stoclet, J.-C. *see* Ouedraogo, S., 112, 867

Stolerman, I.P. *see* Shoaib, M., 111, 1073

Stone, T.W. *see* Hosseinzadeh, H., 112, 316

Stork, A.P. and Cocks, T.M. Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099

Stork, A.P. and Cocks, T.M. Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093

Stow, J. *see* Hall, A., 113, 959

Stratta, F. *see* Mercuri, N.B., 113, 831

Strichartz, G.R. *see* Wang, Z.-Y., 111, 179

Strichartz, G.R. *see* Wang, Z.-Y., 111, 1342

Strohmann, C. *see* Waelbroeck, M., 112, 505

Strong, P. *see* Anderson, R., 113, 1386

Strong, P. *see* Patel, M., 113, 741

Strosberg, A.D. *see* Blin, N., 112, 911

Struijker Boudier, H.A.J. *see* Drieman, J.C., 111, 1117

Su, J., Renaud, N., Carayon, A., Crozatier, B. and Hittinger, L. with the technical assistance of Laplace, M. Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395

Su, M.-J. *see* Lin, C.-H., 113, 275

Su, M.-J. *see* Wu, M.-H., 113, 221

Su, M.-J. *see* Young, M.-L., 113, 69

Suetsugu, T. *see* Saito, K., 111, 465

Sugahara, M. *see* Sato, Y., 112, 625

Sugawara, T. *see* Nakazato, Y., 111, 935

Sugden, D. *see* Chong, N.W.S., 111, 295

Sugidachi, A. *see* Watanabe, M., 111, 123

Sugimoto, T. *see* Kita, Y., 113, 5, 1137

Sumner, M.J. *see* Buchan, K.W., 112, 1251

Suthamnatpong, N., Hosokawa, M., Takeuchi, T., Hata, F. and Takewaki, T. Nitric oxide-mediated inhibitory response of rat proximal colon: independence from changes in membrane potential, 112, 676

Sutliff, R.L. *see* Griesbacher, T., 112, 1004

Suzuki, A. *see* Itoh, T., 111, 311

Suzuki, H., Yoshioka, K., Yanagisawa, M., Urayama, O., Kurihara, T., Hosoki, R., Saito, K. and Otsuka, M. Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310

Suzuki, H. and Kokubun, S. Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117

Suzuki, H. *see* Itoh, T., 111, 165

Suzuki, T. *see* Walsh, D.A., 112, 435

Szabó, C., Southan, G.J., Thiemermann, C. and Vane, J.R. The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Szabó, C., Southan, G.J., Wood, E., Thiemermann, C. and Vane, J.R. Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355

Szilvassy, J. *see* Szilvassy, Z., 112, 999

Szilvassy, Z., Ferdinand, P., Bor, P., Jakab, I., Szilvassy, J., Nagy, I., Lonovics, J. and Koltai, M. Loss of preconditioning in rabbits with vascular tolerance to nitroglycerin, 112, 999

Szwarc, R.S. *see* Baines, C.P., 113, 334

T

Tabrizchi, R. *see* Adeagbo, A.S.O., 111, 13

Tabrizi-Fard, M.A. and Fung, H.-L. Pharmacokinetics, plasma protein binding and urinary excretion of N^{ω} -nitro-L-arginine in rats, 111, 394

Tacke, R. *see* Waelbroeck, M., 112, 505

Takagi, H. *see* Kawabata, A., 111, 503

Takagi, H. *see* Kawabata, A., 112, 547

Takagi, H. *see* Kawabata, A., 112, 817

Takagi, N. *see* Miyake, K., 112, 226

Takahashi, H., Puttick, R.M. and Terrar, D.A. The effects of propofol and enflurane on single calcium channel currents of guinea-pig isolated ventricular myocytes, 111, 1147

Takahashi, H. *see* Hatano, A., 113, 723

Takahashi, M. *see* Ueyama, C., 112, 349

Takeda, M. *see* Hatano, A., 113, 723

Takehara, S. *see* Yasumatsu, H., 111, 1170

Takeo, S. *see* Miyake, K., 112, 226

Taketani, Y. *see* Kasai, Y., 112, 1132

Takeuchi, S. *see* Chiyotani, A., 112, 571

Takeuchi, S. *see* Moritoki, H., 111, 655

Takeuchi, T. *see* Suthamnatpong, N., 112, 676

Takewaki, T. *see* Suthamnatpong, N., 112, 676

Takita, M. *see* Kohno, Y., 112, 1167

Talom, R.T. *see* Makki, T., 111, 555

Tam, F.S.-F., Hillier, K. and Bunce, K.T. Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143

Tamaki, M. *see* Hatano, A., 113, 723

Tamaoki, J. *see* Chiyotani, A., 112, 571

Tamargo, J. *see* Pérez-Viccaño, F., 113, 1281

Tamargo, J. *see* Pérez-Viccaño, F., 113, 1281

Tambourgi, D.V., dos Santos, M.C., de Fátima D. Furtado, M., de Freitas, M.C.W., da Silva, W.D. and Kipnis, T.L. Pro-inflammatory activities in elapid snake venoms, 112, 723

Tammela, T.L.J., Briscoe, J.A.K., Levin, R.M. and Longhurst, P.A. Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195

Tan, K.K.C., Brown, M.J., Longmore, J., Plumpton, C. and Hill, R.G. Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703

Tanabe, J.-i., Watanabe, M., Kondoh, S., Mue, S. and Ohuchi, K. Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480

Tanabe, J.-i. *see* Watanabe, M., 111, 123

Tanaka, H. *see* Saito, K., 111, 465

Tanaka, M. *see* Kim, S., 113, 662

Tanaka, T. *see* Mukai, J., 111, 389

Tanda, K. *see* Williams, F.M., 111, 1123

Tani, Y. *see* Nakazato, Y., 111, 935

Tankosic, P. *see* Makki, T., 111, 555

Tare, M. *see* Tonta, M.A., 112, 525

Tastenoy, M. *see* Waelbroeck, M., 112, 505

Tatchum-Talom, R. *see* Atkinson, J., 111, 1184

Tate, J. *see* Abassi, Z.A., 113, 204

Tattersall, F.D. *see* Trickleton, M.D., 113, 729

Taylor, E.M. and Kaumann, A.J. Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline mesenteric circulation, 111, 264

Taylor, G.W. *see* Kanthakumar, K., 112, 493

Taylor, P.D., Oon, B.B., Thomas, C.R. and Poston, L. Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35

Taylor, P.D., Wickenden, A.D., Mirrlees, D.J. and Poston, L. Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42

Taylor, P.D. and Poston, L. The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801

Teixeira, M.M., Fairbairn, S.M., Norman, K.E., Williams, T.J., Rossi, A.G. and Hellewell, P.G. Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Teixeira, M.M., Reynia, S., Robinson, M., Shock, A., Williams, T.J., Williams, F.M., Rossi, A.G. and Hellewell, P.G. Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811

Teixeira, M.M., Rossi, A.G., Williams, T.J. and Hellewell, P.G. Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332

Teixeira, M.M. and Hellewell, P.G. Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205

Télémaque, S. *see* D'Orléans-Juste, P., 113, 1257

Temma, S. *see* Jino, H., 111, 435

Teng, C.-M. *see* Ko, F.-N., 112, 1174

Teng, C.-M. *see* Lin, C.-H., 113, 275

Teng, C.-M. *see* Lin, C.-H., 113, 1464

Teng, C.-M. *see* Yu, S.-M., 111, 906

Teraoka, H. *see* Nakazato, Y., 111, 935

Teräväinen, T.M.-L. *see* Mervaala, E.M.A., 111, 1189

Terranova, J.P. *see* Dubuc, I., 112, 352

Terrar, D.A. *see* Takahashi, H., 111, 1147

Terrón, J.A. *see* Vallalón, C.M., 113, 13

Tesfamariam, B. and Allen, G.T. β_1 - and β_2 -adrenoceptor antagonist activities of ICI-215001, a putative β_3 -adrenoceptor agonist, 112, 55

Thiemermann, C. *see* Akarasereenont, P., 113, 1522

Thiemermann, C. *see* McMurdo, L., 112, 75

Thiemermann, C. *see* Szabó, C., 112, 355

Thiemermann, C. *see* Szabó, C., 113, 757

Thiemermann, C. *see* Wellings, R.P., 111, 515

Thijssen, H.W. *see* Drieman, J.C., 111, 1117

Thollon, C., Cambarrat, C., Vian, J., Prost, J.-F., Peglion, J.L. and Vilaine, J.P. Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Thomas, C.R. *see* Taylor, P.D., 111, 35

Thompson, J.S., Sheedy, W. and Morice, A.H. Neutral endopeptidase (NEP) inhibition in rats with established pulmonary hypertension secondary to chronic hypoxia, 113, 1121

Thomsen, D.R. *see* Im, W.B., 112, 1025

Thomson, N.C. *see* Nally, J.E., 111, 1163

Thomson, N.C. *see* Nally, J.E., 113, 1328

Thorin, E., Capdeville-Atkinson, C., Cormann, B. and Atkinson, J. Sympathetic neurotransmission in the tail artery of aging rats, 113, 363

Thorin, E. and Atkinson, J. Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351

Thorn, L. *see* Tricklebank, M.D., 113, 729

Thuillez, C. *see* Richard, V., 113, 869

Thurieau, C. *see* Félicitou, M., 112, 683

Thwaites, D.T., Hirst, B.H. and Simmons, N.L. Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

Tibiriça, E. *see* Monassier, L., 111, 1347

Timmerman, H. *see* Jansen, F.P., 113, 355

Timmerman, H. *see* Leurs, R., 112, 847

Toda, N. *see* Enokibori, M., 111, 77

Tognella, S. *see* Gallico, L., 112, 795

Tokumitsu, Y. *see* Charoonroje, P., 113, 569

Tomaki, M. *see* Yamauchi, H., 112, 958

Tominaga, K. *see* Kim, S., 113, 662

Tomkinson, A. *see* Raeburn, D., 113, 1423

Tomkinson, A. *see* Souness, J.E., 111, 1081

Tomlinson, A., Appleton, I., Moore, A.R., Gilroy, D.W., Willis, D., Mitchell, J.A. and Willoughby, D.A. Cyclo-oxygenase and nitric oxide synthase isoforms in rat carrageenin-induced pleurisy, 113, 693

Tomlinson, P.R., Wilson, J.W. and Stewart, A.G. Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

Tomlinson, W. *see* Humphries, R.G., 113, 1057

Tonini, M., Messori, E., Franceschetti, G.P., Rizzi, C.A., Castoldi, A.F., Coccini, T. and Candura, S.M. Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1

Tonini, M. *see* Lucchelli, A., 112, 763

Tonta, M.A., Parkington, H.C., Tare, M. and Coleman, H.A. Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525

Torii, Y., Saito, H. and Matsuki, N. Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431

Törnwall, M., Kaakkola, S., Tuomainen, P., Kask, A. and Männistö, P.T. Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

Tough, I.R. *see* Cox, H.M., 113, 1243

Traber, J. *see* Kappelle, A.C., 111, 887

Tranchina, S., Bernasconi, S., Dejana, E. and Del Maschio, A. Inhibition of human monocyte adhesion to endothelial cells by the coumarin derivative, cloricromene, 111, 575

Travers, A. *see* Hilditch, A., 111, 137

Traynor, J.R. *see* Rupniak, N.M.J., 113, 1487

Treggiari, S. *see* Astolfi, M., 111, 570

Trendelenburg, M. *see* Bültmann, R., 113, 349

Trezise, D.J., Bell, N.J., Kennedy, I. and Humphrey, P.P.A. Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂ purinoceptor classification, 113, 463

Trezise, D.J., Kennedy, I. and Humphrey, P.P.A. The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α, β -methylene adenosine 5'-triphosphate, 112, 282

Tricklebank, M.D., Bristow, L.J., Hutson, P.H., Leeson, P.D., Rowley, M., Saywell, K., Singh, L., Tattersall, F.D., Thorn, L. and Williams, B.J. The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the N-methyl-D-aspartate (NMDA) receptor complex, 113, 729

Trifileff, A., Lach, E., Dumont, P. and Gies, J.-P. Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228

Triggle, C.R. *see* Adeagbo, A.S.O., 111, 13

Triggle, C.R. *see* Pak, J.-M., 112, 1209

Tron, C. *see* Richard, V., 113, 869

Tsarfaty, I. *see* Minc-Golomb, D., 112, 720

Tschaikowski, K., Meisner, M., Schönhuber, F. and Rügheimer, E. Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664

Tschöpl, M. *see* Ouedraogo, S., 112, 867

Tse, H.-W. *see* Jane, D.E., 112, 809

Tsutsumi, O. *see* Kasai, Y., 112, 1132

Tu, A.T. *see* Furukawa, K.-I., 113, 233

Tuček, S. *see* Jakubík, J., 113, 1529

Tucker, J.F. *see* Gibson, A., 113, 1494

Tulp, M.T.H. *see* Siarey, R.J., 111, 497

Tung, S.R. *see* Wang, Z.-Y., 111, 179

Tung, S.R. *see* Wang, Z.-Y., 111, 1342

Tuomainen, P. *see* Törnwall, M., 112, 13

Turner, A.J. *see* Murphy, L.J., 113, 137

Turner, N.C., Dolan, J.S., Grimsditch, D., Lamb, J., Worby, A., Murray, K.J., Coates, W.J. and Warrington, B.H. Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

Turner, N.C., Lamb, J., Worby, A. and Murray, K.J. Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Twissell, D.J. *see* Hilditch, A., 111, 137

Twist, V.W. *see* Kozlowski, R.Z., 111, 250

Twort, C.H.C. *see* Chopra, L.C., 113, 1237

U

Uda, R. *see* Minami, T., 112, 735
 Ujiie, H. *see* Randall, M.D., 112, 49
 Ulman, L.G. *see* Lacroix, J.S., 113, 479
 Umeda, N. *see* Kawabata, A., 111, 503
 Underwood, S.L. *see* Raeburn, D., 113, 1423
 Uneyama, C., Uneyama, H., Takahashi, M. and Akaike, N. Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca^{2+} oscillation, 112, 349
 Uneyama, C. *see* Uneyama, H., 111, 245
 Uneyama, H., Uneyama, C., Ebihara, S. and Akaike, N. Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245
 Uneyama, H. *see* Uneyama, C., 112, 349
 Urayama, O. *see* Suzuki, H., 113, 310
 Urenjak, J. *see* Obrenovitch, T.P., 113, 1295
 Usai, C. *see* Maestrone, E., 111, 283
 Ushio-Fukai, M., Hirano, K. and Kanaide, H. The effects of a novel vasodilator, LP-805, on cytosolic Ca^{2+} concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
 Üstünes, L. *see* De Meyer, G.R.Y., 112, 471
 Usui, H. *see* Jino, H., 111, 435
 Utenthal, L.O. *see* Hermida, O.G., 111, 861

V

Vaca, L. *see* Hasséssian, H., 112, 1118
 Valenzuela, C. *see* Pérez, O., 111, 717
 Vallance, P. *see* Bogle, R.G., 111, 1257
 Vallance, P. *see* MacAllister, R.J., 112, 43
 Vallance, P.J.T. *see* Riezebos, J., 111, 609
 van Buren, T. *see* Kappelle, A.C., 111, 887
 van den Akker, J. *see* Henning, R.H., 111, 459
 van Essen, H. *see* Drieman, J.C., 111, 1117
 Van Etten, M.L. *see* Durcan, M.J., 112, 855
 van Gelderen, E.M. and Saxena, P.R. Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961
 van Hooft, J.A. *see* Kooyman, A.R., 112, 541
 van Kan, F.J.P.M. *see* Drieman, J.C., 111, 1117
 van Tricht, C.L.J. *see* Bax, W.A., 113, 1471
 VanBavel, E. *see* Buus, N.H., 112, 579
 Vanderheijden, P.M.L. *see* Kooyman, A.R., 112, 541
 Vane, J.R. *see* Akarasereenont, P., 113, 1522
 Vane, J.R. *see* Battistini, B., 111, 1009
 Vane, J.R. *see* Battistini, B., 112, 1244
 Vane, J.R. *see* Desai, K.M., 111, 346
 Vane, J.R. *see* Desai, K.M., 113, 1197
 Vane, J.R. *see* Korbut, R., 112, 289
 Vane, J.R. *see* McMurdo, L., 112, 75
 Vane, J.R. *see* Mitchell, J.A., 113, 1008
 Vane, J.R. *see* Szabó, C., 112, 355
 Vane, J.R. *see* Szabó, C., 113, 7571
 Vane, J.R. *see* Warner, T.D., 112, 207
 Vane, J.R. *see* Wellings, R.P., 111, 515
 Vanhoutte, P.M. *see* Boulanger, C.M., 112, 519
 Vannucchi, S. *see* Cavari, S., 113, 254
 Vapaatalo, H. *see* Mervaala, E.M.A., 111, 1189
 Vapaatalo, H. *see* Mervaala, E.M.A., 112, 640
 Vapaatalo, H. *see* Vuorinen, P., 112, 71
 Vargaftig, B.B. *see* Braga da Motta, J.I., 112, 111
 Vargaftig, B.B. *see* Coëffier, E., 113, 749
 Vargaftig, B.B. *see* Zuany-Amorim, C., 112, 1230
 Varma, D.R. *see* Li, D.Y., 112, 59
 Varney, M.A., Galione, A. and Watson, S.P. Lithium-induced decrease in spontaneous Ca^{2+} oscillations in single GH₃ rat pituitary cells, 112, 390
 Vassalle, M. *see* Iacono, G., 113, 289
 Vassort, G. *see* Scamps, F., 113, 982
 Vaughan, P.F.T. *see* McDonald, R.L., 113, 621
 Vázquez, M. *see* Alegret, M., 112, 551
 Vegh, A., Papp, J.G. and Parratt, J. Attenuation of the antiarrhythmic

effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167
 Vegh, A., Papp, J.G. and Parratt, J.R. Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081
 Velasco, C. *see* Morán, A., 113, 1358
 Venkova, K., Milne, A. and Krier, J. Contractions mediated by α_1 -adrenoceptors and P₂-purinoceptors in a cat colon circular muscle, 112, 1237
 Vergara, L.A. *see* Kuijpers, G.A.J., 113, 471
 Vian, J. *see* Thollon, C., 112, 37
 Vié, B. *see* Le Grand, B., 112, 195
 Vigne, P., Feolde, E., Breitmayer, J.P. and Frelin, C. Characterization of the effects of 2-methylthio-ATP and 2-chloro-ATP on brain capillary endothelial cells: similarities to ADP and differences from ATP, 112, 775
 Vrijerberg, H.P.M. *see* Kooyman, A.R., 112, 541
 Vila, E. *see* Chuliá, S., 113, 1377
 Vila, J.M. *see* Martínez, M.C., 113, 419
 Vilaine, J.P. *see* Thollon, C., 112, 37
 Villagrassa, V. *see* Cortijo, J., 111, 769
 Villalón, C.M. and Terrón, J.A. The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13
 Villamil, M.E. *see* Souness, J.E., 111, 1081
 Vincent, J.P. *see* Barelli, H., 112, 127
 Vlaskovska, M. *see* Kasakov, L., 112, 403
 Vollinga, R.C. *see* Jansen, F.P., 113, 355
 von Kügelgen, I., Späth, L. and Starke, K. Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
 von Kügelgen, I. *see* Rump, L.C., 111, 65
 Voss, H.-P. *see* Jansen, F.P., 113, 355
 Vuorinen, P., Wu, X., Arvola, P., Vapaatalo, H. and Pörsti, I. Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71

W

Waddington, J.L. *see* Daly, S.A., 113, 839
 Wadsworth, R. *see* Plevin, R., 112, 311
 Waelbroeck, M., Camus, J., Tastenoy, M., Feifel, R., Mutschler, E., Tacke, R., Strohmann, C., Rafeiner, K., Rodrigues de Miranda, J.F. and Lambrecht, G. Binding and functional properties of hexacyclium and sila-hexacyclium derivatives to muscarinic receptor subtypes, 112, 505
 Wagner, J.A. *see* Chao, A.C., 112, 169
 Waikar, M.V., Ford, A.P.D.W. and Clarke, D.E. Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213
 Wakelam, M.J.O. *see* Nally, J.E., 111, 1163
 Wakelam, M.J.O. *see* Plevin, R., 112, 311
 Waldeck, B. *see* Killström, B.-L., 113, 687
 Waldmeier, P.C., Wicki, P., Feldtrauer, J.-J., Mickel, S.J., Bittiger, H. and Baumann, P.A. GABA and glutamate release affected by GABA_B receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
 Waldron, G.J. and Garland, C.J. Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831
 Waldron, G.J. *see* Parsons, S.J.W., 113, 1275
 Wallace, D.M. *see* Buchan, K.W., 112, 1251
 Wallace, P. *see* Allawi, H.S., 113, 282
 Waller, J., Gardiner, S.M. and Bennett, T. Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
 Walsh, D.A., Salmon, M., Featherstone, R., Wharton, J., Church, M.K. and Polak, J.M. Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea-pig lung, 113, 1407
 Walsh, D.A., Suzuki, T., Knock, G.A., Blake, D.R., Polak, J.M. and Wharton, J. AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435
 Walsh, D.T. *see* Sanz, M.J., 113, 513

Wang, J.P., Raung, S.L., Hsu, M.F. and Chen, C.C. Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945

Wang, Q. *see* Hogg, R.C., 111, 1333

Wang, Q. *see* Hogg, R.C., 112, 977

Wang, Z.-Y., Tung, S.R., Strichartz, G.R. and Håkanson, R. Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342

Wang, Z.-Y., Tung, S.R., Strichartz, G.R. and Håkanson, R. Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179

Warawa, E.J. *see* Keith, R.A., 113, 379

Warburton, P. *see* Balmforth, A.J., 112, 277

Ward, J.K., Fox, A.J., Barnes, P.J. and Belvisi, M.G. Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095

Ward, J.P.T. *see* Chopra, L.C., 113, 1237

Ward, M.E. and Hussain, S.N.A. Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65

Wardell, C.F. and Cunnane, T.C. Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975

Wardle, K.A., Ellis, E.S., Baxter, G.S., Kennett, G.A., Gaster, L.M. and Sanger, G.J. The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789

Warner, T.D., Alcock, G.H. and Vane, J.R. Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207

Warner, T.D. *see* Battistini, B., 111, 1009

Warner, T.D. *see* Battistini, B., 112, 1244

Warner, T.D. *see* Desai, K.M., 111, 346

Warner, T.D. *see* Desai, K.M., 113, 1197

Warner, T.D. *see* Korbut, R., 112, 289

Warner, T.D. *see* Wellings, R.P., 111, 515

Warrington, B.H. *see* Turner, N.C., 111, 1198

Wassenaar, C. *see* Bax, W.A., 113, 1471

Watanabe, K. *see* Horie, S., 112, 87

Watanabe, M., Arakida, Y., Tanabe, J.-i., Sugidachi, A., Hirasawa, N., Mue, S. and Ohuchi, K. Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123

Watanabe, M. *see* Muraki, K., 111, 1154

Watanabe, M. *see* Tanabe, J.-i., 113, 1480

Watanabe, M. *see* Yokota, Y., 113, 324

Watanabe, T. *see* Yamauchi, H., 112, 958

Watanabe, Y. *see* Itoh, T., 111, 311

Watano, T. *see* Koizumi, S., 112, 992

Watkins, J.C. *see* Jane, D.E., 112, 809

Watson, M.L. *see* Weg, V.B., 113, 35

Watson, N. and Eglen, R.M. Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179

Watson, S.P. *see* Varney, M.A., 112, 390

Watson, W.P., Misra, A., Cross, A.J., Green, A.R. and Little, H.J. The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017

Watts, I.S. *see* Buchan, K.W., 112, 1251

Watts, I.S. *see* Riezebos, J., 111, 609

Wayman, C. *see* Gibson, A., 113, 1494

Webb, J. *see* Rupniak, N.M.J., 113, 1487

Webber, S.E. *see* Raeburn, D., 113, 1423

Weber, J.N. *see* Shaunak, S., 113, 151

Weber, S. *see* Larrazet, F., 113, 649

Weg, V.B., Watson, M.L., Faccioli, L.H. and Williams, T.J. Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35

Weg, V.B. *see* Sanz, M.J., 113, 513

Weight, F.F. *see* Peoples, R.W., 113, 555

Weinert, J.S. *see* Eckman, D.M., 111, 1053

Weinhold, C. *see* Böhm, M., 112, 137

Wellings, R.P., Corder, R., Warner, T.D., Cristol, J.-P., Thiemermann, C. and Vane, J.R. Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515

Welsh, N.J., Shankley, N.P. and Black, J.W. Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93

Wennmalm, Å. *see* Jun, T., 113, 345

Wessler, I., Reinheimer, T., Brunn, G., Anderson, G.P., Maclagan, J. and Racké, K. β -Adrenoceptors mediate inhibition of [³H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221

West, D. *see* Purkiss, J.R., 111, 1041

Westfall, D.P. *see* Shinozuka, K., 113, 1203

Weston, A.H. *see* Edwards, G., 113, 1538

Wharton, J. *see* Rutherford, R.A.D., 113, 931

Wharton, J. *see* Walsh, D.A., 112, 435

Wharton, J. *see* Walsh, D.A., 113, 1407

White, D.G. *see* Buchan, K.W., 112, 1251

Whitehead, J.W.F. *see* Gale, J.D., 111, 332

Whitley, G.St.J. *see* Bogle, R.G., 111, 1257

Whitley, G.St.J. *see* MacAllister, R.J., 112, 43

Whittle, B.J.R. *see* Laszlo, F., 111, 1309

Whittle, B.J.R. *see* László, F., 113, 1131

Whittle, B.J.R. *see* Lopez-Belmonte, J., 112, 267

Wickenden, A.D. *see* Taylor, P.D., 111, 42

Wicki, P. *see* Waldmeier, P.C., 113, 1515

Wieczorek, W.J. and Kruk, Z.L. Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

Wienien, W. *see* Schambye, H.T., 113, 331

Wijk, B.v. *see* Schambye, H.T., 113, 331

Wiklund, N.P. *see* Iversen, H.H., 113, 1088

Wiley, J.S., Chen, J.R., Snook, M.B. and Jamieson, G.P. The P_{2X}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

Wilkes, L.C. *see* Purkiss, J.R., 111, 1041

Wilkins, M.R. *see* Hirokawa, K., 112, 396

Wilkins, M.R. *see* Hirokawa, K., 113, 1448

Wilkins, M.R. *see* Rutherford, R.A.D., 113, 931

Wilkinson, G.F. *see* Purkiss, J.R., 111, 723

Wilkinson, G.F. *see* Purkiss, J.R., 111, 1041

Williams, A.J. *see* McGarry, S.J., 111, 1212

Williams, B.A. *see* Smith, P.A., 112, 143

Williams, B.J. *see* Tricklebank, M.D., 113, 729

Williams, F.M., Kus, M., Tanda, K. and Williams, T.J. Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123

Williams, F.M. *see* Teixeira, M.M., 111, 811

Williams, J.L. *see* Baldwin, H.A., 112, 188

Williams, T.J. *see* Norman, K.E., 111, 117

Williams, T.J. *see* Sanz, M.J., 113, 513

Williams, T.J. *see* Teixeira, M.M., 111, 811

Williams, T.J. *see* Teixeira, M.M., 112, 332

Williams, T.J. *see* Teixeira, M.M., 113, 1363

Williams, T.J. *see* Weg, V.B., 113, 35

Williams, T.J. *see* Williams, F.M., 111, 1123

Willis, D. *see* Tomlinson, A., 113, 693

Willoughby, D.A. *see* Tomlinson, A., 113, 693

Wills, P.J. *see* Kanthakumar, K., 112, 493

Wilson, J.W. *see* Tomlinson, P.R., 111, 641

Wilson, R. *see* Kanthakumar, K., 112, 493

Windscheif, U., Ralevic, V., Bäumert, H.G., Mutschler, E., Lambrecht, G. and Burnstock, G. Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015

Wise, H. and Jones, R.L. Characterization of prostanoid receptors on rat neutrophils, 113, 581

Withington, P.G. *see* Grassi-Kassisse, D.M., 113, 675

Wlaź, P. *see* Löscher, W., 112, 97

Woditsch, I. and Schröer, K. Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035

Wolowyk, M.W. *see* Ellory, J.C., 111, 903

Wong, P.S.K., Barclay, P.L., Newman, M.J. and Johns, E.J. The influence of acetazolamide and amlodipine on the intracellular sodium

content of rat proximal tubular cells, 112, 881
 Wong, T.-W. *see* Einstein, R., 111, 199
 Wood, E. *see* Szabó, C., 112, 355
 Wood, J.D. *see* Christofi, F.L., 113, 703
 Wood, M.D. *see* Kennett, G.A., 111, 797
 Woodman, O.L. and Pannangpetch, P. Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443
 Woodman, O.L. *see* Loke, K.E., 112, 630
 Woodman, V.R. *see* Raeburn, D., 113, 1423
 Woodruff, G.N. *see* Boden, P., 112, 717
 Woodruff, G.N. *see* Boden, P.R., 112, 429
 Woodruff, G.N. *see* Boden, P.R., 112, 666
 Woodruff, G.N. *see* Hunter, J.C., 112, 1002
 Woodward, D.F. *see* Regan, J.W., 112, 377
 Woollard, D.J. *see* Johnson, P.J., 111, 1240
 Wooltorton, J.R.A. and Mathie, A. Erratum, 111, 377
 Worby, A. *see* Turner, N.C., 111, 1047
 Worby, A. *see* Turner, N.C., 111, 1198
 Wright, I.K., Amirthappan, S. and Kendall, D.A. Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
 Wright, J. *see* Rabbani, M., 111, 89
 Wu, M.-H., Su, M.-J., Lee, S.-S. and Young, M.-L. The electrophysiological effects of antiarrhythmic potential of a secoaporphine, *N*-allylsecoboldine, 113, 221
 Wu, M.-H. *see* Young, M.-L., 113, 69
 Wu, T.-S. *see* Yu, S.-M., 111, 906
 Wu, T.S. *see* Jansen, F.P., 113, 355
 Wu, X. *see* Vuorinen, P., 112, 71
 Wu, X.C., Johns, E., Michael, J. and Richards, N.T. Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360
 Wu, X.C., Richards, N.T., Michael, J. and Johns, E. Relative roles of nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid in the contractile responses of rat renal arcuate arteries, 112, 369
 Wu, Y.-C. *see* Ko, F.-N., 112, 1174
 Wu, Y.-C. *see* Lin, C.-H., 113, 275
 Wu, Y.-C. *see* Lin, C.-H., 113, 1464
 Wyatt, R.J. *see* Karoum, F., 113, 1391
 Wyllie, M.G. *see* Kenny, B.A., 111, 1003
 Wyse, K.R. *see* Martin, D.K., 111, 873

Y

Yadid, G. *see* Kuijpers, G.A.J., 113, 471
 Yamada, G. *see* Yasuda, N., 113, 21
 Yamamoto, T. *see* Shimizu, S.-i., 113, 564
 Yamamoto, Y. *see* Yasumatsu, H., 111, 1170
 Yamauchi, H., Miura, M., Ichinose, M., Ishikawa, J., Nakajima, N., Tomaki, M., Inoue, H., Maeyama, K., Watanabe, T. and Shirato, K. Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958
 Yanagisawa, M. *see* Hosoki, R., 113, 1141
 Yanagisawa, M. *see* Suzuki, H., 113, 310
 Yang, C.-M. *see* Lin, C.-H., 113, 1464
 Yang, C.M., Hsia, H.-C., Chou, S.-P., Ong, R., Hsieh, J.-T. and Luo, S.-F. Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
 Yang, C.M., Hsia, H.-C., Luo, S.-F., Hsieh, J.-T. and Ong, R. The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
 Yang, C.M., Yo, Y.-L., Hsieh, J.-T. and Ong, R. 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
 Yang, M.C.-M. *see* Liao, J.-F., 111, 439
 Yang, S. *see* Moreau, R., 112, 649
 Yano, M. *see* D'Orléans-Juste, P., 113, 1257
 Yano, M. *see* Fukuroda, T., 113, 336
 Yano, S. *see* Horie, S., 112, 87
 Yard, N.J., Chiesi, M. and Ball, H.A. Effect of cyclopiazonic acid, an inhibitor of sarcoplasmic reticulum Ca²⁺-ATPase, on the frequency-

dependence of the contraction-relaxation cycle of the guinea-pig isolated atrium, 113, 1001
 Yarkony, K.A. *see* Fryer, A.D., 112, 588
 Yarrow, P. *see* Purkiss, J.R., 111, 1041
 Yassin, M. and Scholfield, C.N. NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
 Yasuda, N., Kasuya, Y., Yamada, G., Hama, H., Masaki, T. and Goto, K. Loss of contractile activity of endothelin-1 induced by electrical field stimulation-generated free radicals, 113, 21
 Yasumatsu, H., Morimoto, Y., Yamamoto, Y., Takehara, S., Fukuda, T., Nakao, T. and Setoguchi, M. The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
 Yeadon, M. *see* Herd, C.M., 112, 292
 Yen, M.-H. *see* Lee, Y.-M., 112, 1031
 Yo, Y.-L. *see* Yang, C.M., 111, 777
 Yocca, F.D. *see* Iben, L.G., 113, 1400
 Yoishihara, S. *see* Ricciardolo, F.L.M., 113, 1147
 Yokota, Y., Imaizumi, Y., Asano, M., Matsuda, T. and Watanabe, M. Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
 Yokoyama, T. *see* Kitazawa, T., 111, 288
 Yoshida, K. *see* Kita, Y., 113, 5
 Yoshida, K. *see* Kita, Y., 113, 385
 Yoshida, K. *see* Kita, Y., 113, 1137
 Yoshida, S. *see* Kamei, K., 113, 889
 Yoshihara, S. *see* Ricciardolo, F.L.M., 113, 508
 Yoshioka, K. *see* Hosoki, R., 113, 1141
 Yoshioka, K. *see* Suzuki, H., 113, 310
 Young, J.M. *see* Arias-Montaño, J.A., 111, 598
 Young, J.M. *see* Gibson, W.J., 111, 1262
 Young, L.C. *see* Nally, J.E., 111, 1163
 Young, M.-L., Su, M.-J., Wu, M.-H. and Chen, C.-C. The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69
 Young, M.-L. *see* Wu, M.-H., 113, 221
 Yu, P.-C. *see* Liao, J.-F., 111, 439
 Yu, S.-M., Wu, T.-S. and Teng, C.-M. Pharmacological characterization of cinnamophilin, a novel dual inhibitor of thromboxane synthase and thromboxane A₂ receptor, 111, 906
 Yu, S.-M. *see* Ko, F.-N., 112, 1174
 Yuan, S.Y., Bornstein, J.C. and Furness, J.B. Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095
 Yukimura, T. *see* Kim, S., 113, 662

Z

Zaagsma, J. *see* Coppes, R.P., 113, 342
 Zagorodnyuk, V. and Maggi, C.A. Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
 Zar, M.A. *see* Iravani, M.M., 113, 95
 Zar, M.A. *see* Iravani, M.M., 113, 877
 Zaragozá, F. *see* Pérez-Vizcaíno, F., 113, 1281
 Zazpe, A., Artaiz, I. and Del Río, J. Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968
 Zhang, D. *see* Regan, J.W., 112, 377
 Zhang, Y. and Lang, R.J. Effects of intrinsic prostaglandins on the spontaneous contractile and electrical activity of the proximal renal pelvis of the guinea-pig, 113, 431
 Zhou, Q. *see* Rabbani, M., 111, 89
 Zhu, M.-Y., Juorio, A.V., Paterson, I.A. and Boulton, A.A. Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23
 Zhu, M.-Y. *see* Berry, M.D., 113, 1159
 Zifferblatt, J.B. *see* Chao, A.C., 112, 169
 Ziganshin, A.U., Hoyle, C.H.V., Lambrecht, G., Mutschler, E., Bäumert, H.G. and Burnstock, G. Selective antagonism by PPADS at P_{2X}-purinoceptors in rabbit isolated blood vessels, 111, 923
 Ziganshin, A.U., Hoyle, C.H.V., Ziganshina, L.E. and Burnstock, G. Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669

Ziganshina, L.E. *see* Ziganshin, A.U., 113, 669
Zilkha, E. *see* Obrenovitch, T.P., 113, 1295
Zingarelli, B. *see* Calapai, G., 111, 1328
Zingarelli, B. *see* Squadrito, F., 113, 1153
Zocchi, C. *see* Dionisotti, S., 112, 659

Zohn, I.E. *see* Boyer, J.L., 113, 614
Zonnekeyn, L.L. *see* De Meyer, G.R.Y., 112, 471
Zuany-Amorim, C., Vargaftig, B.B., Maclouf, J. and Pretolani, M.
Strain-dependency of leukotriene C₄ generation from isolated lungs
of immunized mice, 112, 1230

SUBJECT INDEX

A

A

A 68930, The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839

A10 cells, (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174

AA861, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123

Abstinence syndrome, Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215

Acetaldehyde, Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089

Acetylcholine, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

- Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
- Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103
- Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642
- Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895
- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
- The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275
- Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630
- Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Acetylcholine currents, Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935

Acetylcholine receptors, Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271

Acetylcholine receptors, muscarinic, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598

- Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93
- Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179
- Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160
- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55
- Routes of delivery of muscarinic acetylcholine receptors to the plasma membrane in NG108-15 cells, 111, 1023

Acetylcholine receptors, nicotinic, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

- Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695
- Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
- Induction of Na⁺/K⁺-ATPase activity by long-term stimulation of nicotinic acetylcholine receptors in C2C12 myotubes, 111, 459
- Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Potentiation by tonic A_{2A}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582
- Regulation of nicotinic receptors in rat brain following quasi-reversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917
- The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898

Acetylcholine release, β -Adrenoceptors mediate inhibition of [³H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221

- Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703
- Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789
- Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
- Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085
- On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77
- Potentiation by endogenously released ATP of spontaneous transmitter secretion at developing neuromuscular synapses in *Xenopus* cell cultures, 111, 880
- Potentiation by tonic A_{2A}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582
- Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753

Acetylcholinesterase, Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Acetyl salicylic acid, Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983

N'-Acetyl-5-hydroxytryptamine, Presence of P₂-purinoceptors in the rat pineal gland, 112, 107

Acid phosphatase, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N⁶-nitro-L-arginine methyl ester, 113, 1127

Acinar cells, Down-regulation of bombesin binding to guinea-pig

pancreatic acinar cells during homologous desensitization, 112, 1037

Acoustic rhinometry, Reduction by Hoe 140, the B₂ kinin receptor antagonist, of antigen-induced nasal blockage, 111, 969

ACPD, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

- Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364
- Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231

ACTH, Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171

ACTH-like peptides, A dose-response analysis of the beneficial effects of the ACTH-(4-9) analogue, Org 2766, on behavioural recovery following unilateral labyrinthectomy in guinea-pig, 111, 358

Actinomycin D, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

Actinonin, Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310

Action potentials, Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717

- Effects of intrinsic prostaglandins on the spontaneous contractile and electrical activity of the proximal renal pelvis of the guinea-pig, 113, 431
- The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747

Action potential duration, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527

- Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717

Adenine, Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca²⁺ oscillation, 112, 349

Adenine nucleotides, Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71

- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815

Adenosine, Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7

- Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031
- Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29
- The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316
- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123
- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N⁶-monomethyl-L-arginine but not glibenclamide, 111, 711
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Modulation of vasodilatation to levromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
- Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753
- Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137

Adenosine deaminase, Potentiation by tonic A_{2A}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582

Adenosine receptors, Adenosine A_{2B}-receptor-mediated cyclic AMP accumulation in primary rat astrocytes, 111, 191

- Characterization of the adenosine receptors mediating hypothermia in the conscious mouse, 113, 1386
- A comparison of A₂ adenosine receptor-induced cyclic AMP generation in cerebral cortex and relaxation of pre-contracted aorta, 111, 185
- Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂ but not to an A₁-agonist in the dog, 111, 1233

Adenosine A₁ receptors, A₁ adenosine receptor inhibition of cyclic AMP formation and radioligand binding in the guinea-pig cerebral cortex, 113, 1501

- Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159
- The binding of 1,3-[³H]-dipropyl-8-cyclopentylxanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249
- Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252
- Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703

Adenosine A₁ receptor agonists, Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031

Adenosine A₂ receptors, Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7

- Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- Evidence for a cooperation between adenosine A₂ receptors and B₁-adrenoceptors on cardiac automaticity in the isolated right ventricle of the rat, 111, 1316

Adenosine A_{2A} receptor antagonists, Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659

Adenosine A_{2A} receptors, Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43

- Potentiation by tonic A_{2A}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582

Adenosine A₃ receptors, Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741

Adenosine 5'-O-(3-thiophosphate) (ATP γ S), Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815

Adenosine 5'-o-2-iodiphosphate, Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117

Adenylyl cyclase, Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775

- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377
- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240

Adenylyl cyclase inhibition, Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614

Adhesion molecules, Inhibition of human monocyte adhesion to endothelial cells by the coumarin derivative, clorcromene, 111, 575

- Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117
- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Adipose tissue, Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866

- Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911
- ADP, FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2U} -purinoceptors, 113, 1057
- Adrenal chromaffin cells**, Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
- Adrenal cortex**, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179
- Adrenaline**, Sustained prejunctional facilitation of noradrenergic neurotransmission by adrenaline as a co-transmitter in the portal vein of freely moving rats, 113, 342
- Adrenergic motor transmission**, Presence of neuropeptide Y in the rat seminal vesicle and its effects on noradrenaline- and nerve-induced contractions, 113, 877
- Adrenergic responsiveness**, Beneficial effect of the Ca^{2+} antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887
- Adrenoceptors**, Adrenoceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576
- Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105
- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769
- Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154
- Adrenoceptor agonists**, Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159
- α -Adrenoceptors**, The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- The effects of perfusion rate and N^G -nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
- Heterogeneity of α_1 -adrenoceptor subtypes involved in adrenergic contractions of dog blood vessels, 112, 1167
- Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- The noradrenaline precursor L-*threo*-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- α -Adrenoceptor agonists**, Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649
- α_1 -Adrenoceptors**, α_1 -Adrenoceptors in the conduction system of rat hearts, 111, 465
- Contractions mediated by α_1 -adrenoceptors and P_2 -purinoceptors in a cat colon circular muscle, 112, 1237
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769
- Pharmacological properties of the cloned α_{1AD} -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral cortex and vas deferens, 111, 1003
- Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractions in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195
- Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439
- α_1 -Adrenoceptor activation**, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007
- α_1 -Adrenoceptor antagonists**, Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377
- Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractions in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195
- α_{1A} -Adrenoceptors**, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151
- Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539
- Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- α_{1B} -Adrenoceptors**, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151
- Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539
- Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- α_{1D} -Adrenoceptors**, (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
- α_1 -Adrenoceptor subtypes**, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151
- α_{1B} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819
- (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
- Heterogeneity of α_1 -adrenoceptor subtypes involved in adrenergic contractions of dog blood vessels, 112, 1167
- Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- Pharmacological properties of the cloned α_{1AD} -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral cortex and vas deferens, 111, 1003
- α_2 -Adrenoceptors**, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855
- Effect of potassium channel blockade and α_2 -adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341
- Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411
- Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485
- Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769
- Rapid desensitization of noradrenaline- and neuropeptide Y-stimulated Ca^{2+} mobilization in HEL-cells, 112, 499
- α_2 -Adrenoceptor antagonists**, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231
- β -Adrenoceptors**, β -Adrenoceptors mediate inhibition of [3 H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221
- The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311
- Effects of muscarinic M_2 and M_3 receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179
- Evidence for a cooperation between adenosine A_2 receptors and β_1 -adrenoceptors on cardiac automaticity in the isolated right ventricle of the rat, 111, 1316
- Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296
- Role of cyclic AMP-induced Cl⁻ conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137
- β -Adrenoceptors, atypical**, Adrenoceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576
- Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699
- β_1 - and β_2 -adrenoceptor antagonist activities of ICI-215001, a putative β_3 -adrenoceptor agonist, 112, 55
- Mediation of most atypical effects by species homologues of the β_3 -

adrenoceptor, 112, 911

β -Adrenoceptor agonists, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493

- Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687

β -Adrenoceptor antagonists, The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311

- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

β -Adrenoceptor ligands, Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866

β -Adrenoceptor subtypes, Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296

β_1 -Adrenoceptors, Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866

- Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D_1 -, β_1 - and β_2 -receptor-mediated responses, 112, 595
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769

β_2 -Adrenoceptors, Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866

- Effects of N^0 -nitro-L-arginine methyl ester on regional haemodynamic responses to $MgSO_4$ in conscious rats, 111, 325
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769

β_2 -Adrenoceptors, prejunctional, Sustained prejunctional facilitation of noradrenergic neurotransmission by adrenaline as a co-transmitter in the portal vein of freely moving rats, 113, 342

β_2 -Adrenoceptor agonists, Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199

- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

β_3 -Adrenoceptors, Adrenoceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576

- Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911
- Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

AF-DX116, Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Afferent nerve endings, Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681

Afferent nerve stimulation, Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181

Afferent neurones, Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547

- Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681

Affinity constants, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

- pK_1 values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

Afterload, Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334

α -Agatoxin IVA, Depression of high-threshold calcium currents by activation of human D_2 (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

Ageing, Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat, 111, 1184

- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363

Aggrecan, Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Airway hyperresponsiveness, Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292

- Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7

Airways, Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7
- Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649

Airways, guinea-pig, Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147

- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095

Alcohol, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Alcohols (aliphatic), Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089

Alcuronium, Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Aldose reductase inhibitors, The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801

- The effects of aldose reductase inhibition with ponarrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponarrestat, 111, 42

Allergen challenge, Reduction by Hoe 140, the B_2 kinin receptor antagonist, of antigen-induced nasal blockage, 111, 969

- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Allergic rhinitis, Reduction by Hoe 140, the B_2 kinin receptor antagonist, of antigen-induced nasal blockage, 111, 969

Allergy, Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205

- Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Role of interleukin-5 in enhanced migration of eosinophils from airways of immunized guinea-pigs, 113, 749

Allodynia, Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735

***N*-Allylsecoboldine**, The electrophysiological effects of antiarrhythmic potential of a secoaporphine, *N*-allylsecoboldine, 113, 221

Alzheimer's disease, Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Amacrine cells, Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789

Amastatin, Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570

6-Amidino-2-naphthol, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na⁺ and K⁺ transport properties in the rabbit cortical collecting duct, 111, 173

Amine uptake, Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579

Amino acids, Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188

α -Aminoadipate, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, α aminoadipate, 113, 536

Aminoglycosides, Decrease in calcium currents induced by

aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Aminopeptidases, Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570

3-Aminopropylphosphinic acid, Peripheral and central sites of action of GABA_A agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344

4-Aminopyridine, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103

Amitriptyline, Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199

AMPA, A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843

- Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

Amphetamines, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

- Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
- Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391

Amrinone, A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757

Amygdala, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

- Paired-pulse depression of the N-methyl-D-aspartate receptor-mediated synaptic potentials in the amygdala, 113, 1029

Amylase, Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037

Anaesthetics, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Analgesia, Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004

- Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents, 113, 1416
- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352
- The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503
- Role of histamine in rodent antinociception, 111, 1269

Anaphylaxis, Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

Anaphylaxis, passive cutaneous, Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205

- Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35

Angina, Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137

Angiogenesis, Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894

Angiotensin, Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137

- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

Angiotensin-converting enzymes, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

- Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837
- Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652

Angiotensin-converting enzyme inhibitors, Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717

- Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555
- Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Angiotensin peptides, Natriuretic action of angiotensin(1-7), 111, 1

- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652

Angiotensin receptors, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

Angiotensin receptor subtypes, Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Angiotensin receptor antagonism, Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652

Angiotensin receptor antagonists, Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Angiotensin receptor antagonists, non-peptide, Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Angiotensin AT₁ receptors, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662

- Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137
- Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Angiotensin AT₁, receptors, human, Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Angiotensin AT₁ receptor subtypes, Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195

- Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Angiotensin AT₁ receptor antagonists, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662

- Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137
- A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145

Angiotensin-II, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

- Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195
- Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277
- Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351
- Natriuretic action of angiotensin(1-7), 111, 1

Angiotensin II receptors, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

Anococcygeus muscle, Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089

- Noradrenergic-nitergic interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403
- Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494

Anoxia, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221

ANQ9040, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Antagonism, competitive, Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Antagonism, insurmountable, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

- Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Antagonism, surmountable, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

Antagonists, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

Antagonists, competitive, pK₁ values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

Anti-anginal drugs, Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137

Anti-arrhythmic drugs, Antifibrillatory effects of clofilium in the rabbit isolated heart, 113, 209

- Block of the delayed rectifier current (I_R) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
- Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717
- Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873
- The electrophysiological effects of antiarrhythmic potential of a secoaporphine, N-allylsecoboldine, 113, 221
- The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69
- Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673

Antibodies, monoclonal, Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703

Antibody microprobes, Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951

Anticonvulsant drugs, The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the N-methyl-D-aspartate (NMDA) receptor complex, 113, 729

- Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89

Antidepressant effects, Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltindole, 111, 956

Antidepressants, Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741

- The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97
- Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

Anti-fungal agents, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

Antigen, Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292

- Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958

Antigen-induced contractions, Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259

Anti-inflammatory, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

Antilaminin IgG, Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271

Antinociception, Antinociceptive and toxic effects of (+)-epibatidine oxalate attributable to nicotinic agonist activity, 113, 1487

- Central and peripheral actions of the novel κ-opioid receptor agonist, EMD 60400, 111, 837
- Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817
- Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002
- Modulation of morphine antinociception in the mouse by endogenous nitric oxide, 113, 1372
- The noradrenaline precursor L-*threo*-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α-adrenoceptors in the mouse, 111, 503
- A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 61753, 113, 1317
- Role of histamine in rodent antinociception, 111, 1269

Antiplatelet activity, Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385

Antiplatelet drugs, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

- Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Antipsychotics, The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839

Antispasmodic activity, Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

Antithrombotic agents, Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

Antitussive agents, Mogusine: a novel peripheral non-narcotic antitussive drug, 112, 795

- Peripheral and central sites of action of GABA_B agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344

Anxiety, *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797

Anxiolytic, Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

Aorta, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683

- Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
- (-)-Discretamine, a selective α_{1D}-adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
- The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Role of Ca²⁺-dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022
- Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604

Aorta, guinea-pig, A comparison of A₂ adenosine receptor-induced cyclic AMP generation in cerebral cortex and relaxation of pre-contracted aorta, 111, 185

Aorta, human, [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4

Aorta, rabbit, Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolaemic rabbits, 113, 190

Aorta, rat, Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370

– Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42

– *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203

– Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377

– Recovery after dietary vitamin E supplementation of impaired endothelial function in vitamin E-deficient rats, 112, 515

– Regulation of vascular and gastric smooth muscle contractility by pervaanadate, 113, 403

– Relaxation of rat thoracic aorta induced by the Ca^{2+} -ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655

– Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445

– Voltage- and time-dependent inhibitory effects on rat aortic and porcine coronary artery contraction induced by propafenone and quinidine, 113, 1281

Aortic distensibility, Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555

L-AP4, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

– Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364

Apamin, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607

– Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077

– Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958

– Nitric oxide-mediated inhibitory response of rat proximal colon: independence from changes in membrane potential, 112, 676

APNEA, Characterization of the adenosine receptors mediating hypothermia in the conscious mouse, 113, 1386

– Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741

Aprotinin, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81

Apyrase, Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Aqueous humour, Role of cyclic AMP-induced Cl⁻ conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137

Arachidonic acid, Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509

– Pharmacological characterization of the human histamine H₂ receptor stably expressed in Chinese hamster ovary cells, 112, 847

Arachidonic acid metabolites, Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548

Argatroban, Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

Arginine, Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159

D-Arginine, L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205

L-Arginine, Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817

– Differential effects of lumenal L-arginine and N^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304

– Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282

– Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547

– L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205

– Modulation of morphine antinociception in the mouse by endogenous nitric oxide, 113, 1372

– NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345

– Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035

– Reduction by N^G-nitro-L-arginine of H₂O₂-induced endothelial cell injury, 113, 564

– Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65

Arginine transport, Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487

Arginine vasopressin, Effect of dehydration and hyperosmolal hydration on lignocaine and metabolites disposition in conscious rabbits, 113, 317

Argininosuccinic acid, Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65

Aromatic L-amino acid decarboxylase, The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503

– Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23

Arrhythmias, Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031

– Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167

– Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083

– The dose-related hyper-and-hypokalaemic effects of salbutamol and its arrhythmogenic potential, 111, 73

– The electrophysiological effects of antiarrhythmic potential of a secoaporphine, N-allylsecoboldine, 113, 221

– Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081

– Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869

Arterial bed, rat mesenteric, Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoreceptors, 113, 1015

Arterial occlusion, Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912

Arteries, Effects of selective ET_B-receptor stimulation on arterial, venous and capillary functions in cat skeletal muscle, 112, 888

– Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609

– Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723

– Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525

Arteries, basilar, Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324

Arteries, carotid, Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029

– Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

Arteries, caudal, *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203

Arteries, cerebral, Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635

– Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939

Arteries, common carotid, Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699

Arteries, coronary, Enhancement of noradrenergic constriction of large

coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443

- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by *N*^G-monomethyl-L-arginine but not glibenclamide, 111, 711
- Loss of contractile activity of endothelin-1 induced by electrical field of stimulation-generated free radicals, 113, 21
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- The role of myoendothelial cell contact in non-nitric oxide-, non-prostanoid-mediated endothelium-dependent relaxation of porcine coronary artery, 113, 1289
- Voltage- and time-dependent inhibitory effects on rat aortic and porcine coronary artery contraction induced by propafenone and quinidine, 113, 1281

Arteries, coronary, bovine, Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659

Arteries, coronary, dog, Lack of importance of NO in β -adrenoceptor-mediated relaxation of large epicardial canine coronary arteries, 111, 981

Arteries, coronary, guinea-pig, Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

Arteries, coronary, human, Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471

- The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093

Arteries, coronary, porcine, Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557

- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652

Arteries, cutaneous, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105

Arteries, ear, rabbit, Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550

- Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233
- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123

Arteries, femoral, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683

- Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550

Arteries, femoral, rabbit, Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278

- The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173

Arteries, mesenteric, Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831

- Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579
- The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801
- Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71
- Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
- Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360
- Role of R-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439

Arteries, mesenteric, cat, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435

Arteries, mesenteric, human, Relaxation of human isolated mesenteric arteries by vasopressin and desmopressin, 113, 419

Arteries, mesenteric, rat, Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673

Arteries, mesenteric resistance, Endothelial function in the isolated perfused mesentery and aorta of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42

- Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35

Arteries, pulmonary, Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972

- Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β_1 - and β_2 -receptor-mediated responses, 112, 595
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311

Arteries, pulmonary, human, Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

- Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

Arteries, pulmonary, rabbit, Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336

Arteries, renal arcuate, Relative roles of nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid in the contractile responses of rat renal arcuate arteries, 112, 369

Arteries, renal, human, Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49

Arteries, saphenous, Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109

Arteries, splenic, Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321

Arteries, subcutaneous, Effects of dietary marine oil supplementation on reactivity of human buttock subcutaneous arteries and forearm veins *in vitro*, 112, 566

Arteries, tail, Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853

- Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439

Arteries, tail, rat, Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625

- Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867
- Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351
- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363

Arteries, temporal, Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77

Arteriovenous anastomoses, Inhibition of nitric oxide biosynthesis and

carotid arteriovenous anastomotic shunting in the pig, 111, 961

Arthritis, The role of bradykinin B₁ receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940

- Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_A receptor system, 113, 1561

Arthus reaction, Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117

- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Aspartate release, Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379

Aspirin, Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Asthma, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- The dose-related hyper-and-hypokalaemic effects of salbutamol and its arrhythmogenic potential, 111, 73
- The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

Astrocytes, Adenosine A_{2B}-receptor-mediated cyclic AMP accumulation in primary rat astrocytes, 111, 191

- The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997

Astrocytoma cells, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598

- Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299

Atenolol, Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925

- Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nifedipine, 113, 1064

Atherosclerosis, The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386

Atipamezole, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

ATP, Activation of chloride current by P₂-purinoceptors in rat ventricular myocytes, 111, 1355

- Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972
- Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853
- Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942
- Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29
- Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191
- Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569
- Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219

- Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493
- Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂ purinoceptor classification, 113, 463
- Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K⁺ channel in mouse β -cells, 111, 302
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2X}-purinoceptor ligands, 113, 1432
- Pharmacological profile of the ATP-mediated increase in L-type calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
- Potentiation by endogenously released ATP of spontaneous transmitter secretion at developing neuromuscular synapses in *Xenopus* cell cultures, 111, 880
- Release of Ca²⁺ from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca²⁺ entry, 113, 457
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455
- The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015

ATP, extracellular, Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296

- A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST_{GS}, 111, 1135

ATP derivatives, Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151

- Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191

ATP potentiation, Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473

ATP release, Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975

- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65

ATPase, Induction of Na⁺/K⁺-ATPase activity by long-term stimulation of nicotinic acetylcholine receptors in C2C12 myotubes, 111, 459

ATPase, calcium, Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655

ATPase, Na⁺,K⁺, Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87

ATP- γ -S, Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191

ATP-receptors, The P_{2X}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

ATP-sensitive potassium channels, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

- Antifibrillatory effects of clofibrate in the rabbit isolated heart, 113, 209
- Block of ATP-sensitive K⁺ channels in isolated mouse pancreatic β -cells by 2,3-butanedione monoxime, 112, 143
- Interaction of tolbutamide and cytosolic nucleotides in controlling the

ATP-sensitive K^+ channel in mouse β -cells, 111, 302

- Mg^{2+} -dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632

Atria, Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D_1 -, β_1 - and β_2 -receptor-mediated responses, 112, 595

Atria, guinea-pig, β_1 - and β_2 -adrenoceptor antagonist activities of ICI-215001, a putative β_3 -adrenoceptor agonist, 112, 55

Atria, rat, Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractions in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195

Atrial natriuretic peptide, Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146

- Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569
- Hydrolysis of iodine labelled urotilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204
- Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1, 113, 931
- The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328
- Neutral endopeptidase (NEP) inhibition in rats with established pulmonary hypertension secondary to chronic hypoxia, 113, 1121

Atrioventricular nodes, α_1 -Adrenoceptors in the conduction system of rat hearts, 111, 465

Atrium, isolated, Effect of cyclopiazonic acid, an inhibitor of sarcoplasmic reticulum Ca^{2+} -ATPase, on the frequency-dependence of the contraction-relaxation cycle of the guinea-pig isolated atrium, 113, 1001

Autonomic ganglia, Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55

Autoradiography, [^{125}I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303

- α_1 -Adrenoceptors in the conduction system of rat hearts, 111, 465
- AT_1 receptor characteristics of angiotensin analogue binding in human synovium, 112, 435
- Characterization of the binding of the first selective radiolabelled histamine H_3 -receptor antagonist, [^{125}I]-iodophenpropit, to rat brain, 113, 355
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1, 113, 931
- Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296
- Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749

Autoreceptors, Cyclothiazide unmasks AMPA-evoked stimulation of [3H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339

- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635

Azelastine, Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111

B

Baclofen, The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316

- The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497
- Peripheral and central sites of action of GABA_B agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344

BAPTA, Lithium-induced decrease in spontaneous Ca^{2+} oscillations in single GH₃ rat pituitary cells, 112, 390

- Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753

Barbiturates, Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89, 89

Barium influx, The P_{2X}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

Bay K 8644, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017

- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831

Behaviour, Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250

- Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

Benserazide, Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

Benzamides, Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143

Benzodiazepines, Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

- Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Benzodiazepine receptors, partial agonists, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170

Benzodiazepine receptors, peripheral, Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429

- Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Benzodiazepine sites, Benzodiazepine-induced intestinal motor disturbances in rats: mediation by w_2 (BZ₂) sites on capsaicin-sensitive afferent neurones, 113, 268

Benzoylbenzoic ATP, The P_{2X}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

Benzyltetrahydroisoquinolines, Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377

Bethanechol, Differential effects of Na^+, K^+ -ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87

Bile salts, Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209

BIM23027, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

BIM23056, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

BIMU 8, Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213

Binding studies, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

Binding/functional correlations, Binding and functional properties of hexocyclium and sila-hexocyclium derivatives to muscarinic receptor subtypes, 112, 505

Bipyridylum, Effects of bipyridylum compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle, 112, 1216

Bisoprolol, Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

Bladder, Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219

- Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195
- Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111

Bladder, urinary, Comparative studies on the affinities of ATP

derivatives for P_{2X} -purinoceptors in rat urinary bladder, 112, 1151

- Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669
- Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95

Bladder, urinary, human, Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1

Bladder, urinary, rat, Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Bladder afferents, Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017

Bleeding time, Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

Blood-brain barrier, Characterization of the effects of 2-methylthio-ATP and 2-chloro-ATP on brain capillary endothelial cells: similarities to ADP and differences from ATP, 112, 775

- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041

Blood flow, Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolaemic rabbits, 113, 190

- Differential effects of luminal L-arginine and N^G -nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961
- Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y₂ prejunctional receptor, 113, 479
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65

Blood flow, bronchial, Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918

Blood flow, carotid, The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13

- The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

Blood flow, cerebral, Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226

Blood flow, coronary, Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233

Blood flow, cutaneous, Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181

- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Blood flow, gastric, The involvement of endothelial dysfunction, nitric oxide and prostanooids in the rat gastric microcirculatory responses to endothelin-1, 112, 267

- Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036

Blood flow, intestinal, L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205

Blood flow, nerve, Beneficial effect of the Ca²⁺ antagonist, nifedipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887

- Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425

Blood pressure, Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

- AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662
- Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547
- Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137
- Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

- Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825
- Differential effects of luminal L-arginine and N^G -nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925
- Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature, 112, 704
- Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555
- Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358
- Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837
- NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
- A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567

Blood vessels, Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

- Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Heterogeneity of α_1 -adrenoceptor subtypes involved in adrenergic contractions of dog blood vessels, 112, 1167
- Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109

Blood vessels, isolated, Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 496

Blood vessels, pulmonary, Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H₃ receptors in the trachea, 111, 445

Blood vessels, rabbit, Selective antagonism by PPADS at P_{2X} -purinoceptors in rabbit isolated blood vessels, 111, 923

BMS-180560, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

BMS-182874, Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251

BMY-7378, Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400

BMY-14802, The effects of sigma ligands and of neuropeptide Y on N-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709

BN 50730, Long-lasting inhibitory activity of the tetrazepinic BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994

Bombesin, Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037

Bosentan, Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

- Effects of bosentan (Ro 47-0203), an ET_A, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869

BQ-123, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

- Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336
- BQ-788**, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336
- BQ-3020**, [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- 8Br-cyclic GMP**, Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- Bradycardic agents**, Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
- Bradykinin**, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683
- Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167
- Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228
- Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
- Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159
- Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029
- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
- Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
- Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983
- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
- The role of bradykinin B₁ receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940
- Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
- Role of K⁺ channels in the vasodilator response to bradykinin in the rat heart, 113, 954
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Des-Arg⁹-Bradykinin**, Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029
- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Bradykinin binding sites**, Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228
- Bradykinin receptors**, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- The role of bradykinin B₁ receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940
- Bradykinin receptor antagonists**, Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004
- Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
- Bradykinin receptor mechanisms**, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Bradykinin B₁ receptors**, Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029
- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389
- Bradykinin B₂ receptors**, Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167
- Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
- Bradykinin B₂ receptor antagonists**, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683
- Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- Reduction by Hoe 140, the B₂ kinin receptor antagonist, of antigen-

induced nasal blockage, 111, 969

- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Brain, A₁, adenosine receptor inhibition of cyclic AMP formation and radioligand binding in the guinea-pig cerebral cortex, 113, 1501

- Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400
- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347
- Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391

Brain, guinea-pig, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598

Brain, human, Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041

Brain, preoptic area, Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328

Brain, rat, The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997

- Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Separation and characterization of a novel isoenzyme of cyclic nucleotide phosphodiesterase from rat cerebrum, 111, 389

Brain ischaemia, Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a N-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247

- Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188

Brain slices, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

BRL 35135, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

BRL 37344, Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699

Bronchi, bovine, Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163

Bronchi, human, Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570

- The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687
- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

Bronchial hyperresponsiveness, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

Bronchioles, Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237

Bronchoconstriction, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Bronchodilation, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

- Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

Bronchopulmonary system, rat, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

Bronchospasm, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

Brown adipose tissue, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

Brush border membranes, Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509

BTS 54 505, The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97

α -Bungarotoxin, Regulation of nicotinic receptors in rat brain following quasi-reversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917

- The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898

Buspirone, Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

2,3-Butanedione monoxime, Block of ATP-sensitive K⁺ channels in isolated mouse pancreatic β -cells by 2,3-butanedione monoxime, 112, 143

γ -Butyrolactone, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

BW755C, Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360

C

C5a, Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117

C6 glioma cells, Endogenous expression of histamine H₁ receptors functionally coupled to phosphoinositide hydrolysis in C6 glioma cells: regulation by cyclic AMP, 113, 1554

Ca²⁺ ionophores, Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099

Ca²⁺-ATPase, Effect of cyclopiazonic acid, an inhibitor of sarcoplasmic reticulum Ca²⁺-ATPase, on the frequency-dependence of the contraction-relaxation cycle of the guinea-pig isolated atrium, 113, 1001

Caco-2 cells, Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

Cadaverine, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Cadmium, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

Caecum, Y₂-receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Caffeine, Activation of the sheep cardiac sarcoplasmic reticulum Ca²⁺-release channel by analogues of sulmazole, 111, 1212

- Ca²⁺ release induced by myotoxin α , a radio-labellable probe having novel Ca²⁺ release properties in sarcoplasmic reticulum, 113, 233
- Comparison of the effects of caffeine and other methylxanthines on

- [Ca²⁺]_i in rat ventricular myocytes, 111, 455
- The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103
- Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
- Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
- Calcitonin gene-related peptide (CGRP),** Calcitonin gene-related peptide receptors in human gastrointestinal epithelia, 113, 1243
- Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703
- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123
- The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262
- Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588
- Interleukin-18 enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
- Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425
- Potentiation by tonic A_{2a}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582
- Calcitonin gene-related peptide (CGRP) receptors,** Calcitonin gene-related peptide receptors in human gastrointestinal epithelia, 113, 1243
- Calcium,** Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
- Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037
- The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316
- The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
- Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539
- Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640
- Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449
- Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753
- Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169
- Stimulation by menthol of C1 secretion via a Ca²⁺-dependent mechanism in canine airway epithelium, 112, 571
- Studies on the mechanism of [³H]-noradrenaline release from SH-SY5Y cells: the role of CA²⁺ and cyclic AMP, 111, 787
- Calcium, cytoplasmic,** Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca²⁺ oscillation, 112, 349
- Calcium, cytosolic,** Dual effects of tetradrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767
- The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
- A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST₈₈₅, 111, 1135
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- Calcium, extracellular,** Contribution of intra- and extracellular Ca²⁺ to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
- Calcium, intracellular,** α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151
- All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972
- Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
- Characterization of prostanoid receptors on rat neutrophils, 113, 581
- Comparison of the effects of caffeine and other methylxanthines on [Ca²⁺]_i in rat ventricular myocytes, 111, 455
- Contribution of intra- and extracellular Ca²⁺ to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
- Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252
- Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57
- Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926
- Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
- Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
- Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Modulation of dopamine and noradrenaline release and of intracellular Ca²⁺ concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
- Release of Ca²⁺ from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca²⁺ entry, 113, 457
- Calcium antagonists,** Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579
- Calcium-activated chloride currents,** Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977
- Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57
- Calcium-activated potassium currents,** Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977
- Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca²⁺ oscillation, 112, 349
- Calcium/calmodulin-dependent protein kinase,** Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169
- Calcium channels,** Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375
- Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299
- Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells, 111, 483
- Inhibition by SK&F 96365 of Ca²⁺ current, IL-2 production and

- activation in T lymphocytes, 113, 861
- Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209
- Modulation of dopamine and noradrenaline release and of intracellular Ca^{2+} concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- Muscarinic (M_1) receptor-mediated inhibition of K^+ -evoked [3H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L - and N -type Ca^{2+} channels, 113, 621
- Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154
- Calcium channels, L -type**, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca^{2+} current in guinea-pig heart, 113, 451
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Modulation of cardiac L -type Ca^{2+} channels by GTP γ S in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
- Potentiation by endogenously released ATP of spontaneous transmitter secretion at developing neuromuscular synapses in *Xenopus* cell cultures, 111, 880
- Specific inhibition of Ca -activated K channels in red cells by selected dihydropyridine derivatives, 111, 903
- Calcium channels, N -type**, Fluspirilene block of N -type calcium current in NGF-differentiated PC12 cells, 111, 483
- Calcium channels, R -type**, Role of R -type calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Calcium channels, voltage-sensitive**, Differential inhibition of neuronal calcium entry and [3H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Calcium channel antagonists**, Beneficial effect of the Ca^{2+} antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887
- Comparison of the haemodynamic profiles of elgiodipine and nicardipine in the anaesthetized dog, 111, 49
- Contribution of intra- and extracellular Ca^{2+} to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
- The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017
- Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca^{2+} current in guinea-pig heart, 113, 451
- Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767
- Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111
- Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- Fluspirilene block of N -type calcium current in NGF-differentiated PC12 cells, 111, 483
- The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881
- Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377
- Pharmacological characteristics of liriodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275
- Calcium currents**, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375
- Depression of high-threshold calcium currents by activation of human D_2 (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061
- Effects of metabolic blockers on Ca^{2+} -dependent currents in cultured sensory neurones from neonatal rats, 111, 57
- The effects of propofol and enflurane on single calcium channel currents of guinea-pig isolated ventricular myocytes, 111, 1147
- The electrophysiological effects of antiarrhythmic potential of a secoaporphine, *N*-allylseboldine, 113, 221
- Functional expression of human D_3 dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391
- Pharmacological profile of the ATP-mediated increase in L -type calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Pharmacology of high-threshold calcium currents in GH_4C_1 pituitary cells and their regulation by activation of human D_2 and D_4 dopamine receptors, 112, 728
- Calcium dependence**, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- Calcium entry**, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151
- Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- Calcium influx**, Blockade of the inward rectifier potassium current by the Ca^{2+} -ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118
- Differential inhibition of neuronal calcium entry and [3H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103
- Calcium ions**, Dual effects of mastoparan on intracellular free Ca^{2+} concentrations in human astrocytoma cells, 112, 299
- Calcium mobilization**, Effects of a newly synthesized K^+ channel opener, Y-26763, on noradrenaline-induced Ca^{2+} mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- Rapid desensitization of adrenaline- and neuropeptide Y-stimulated Ca^{2+} mobilization in HEL-cells, 112, 499
- Calcium oscillations**, Lithium-induced decrease in spontaneous Ca^{2+} oscillations in single GH_3 rat pituitary cells, 112, 390
- Calcium release**, All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972
- Ca^{2+} release induced by myotoxin α , a radio-labellable probe having novel Ca^{2+} release properties in sarcoplasmic reticulum, 113, 233
- Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767
- Effects of bipyridylum compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle, 112, 1216
- Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- Release of Ca^{2+} from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca^{2+} entry, 113, 457
- Calcium-release channels, cardiac**, Activation of the sheep cardiac sarcoplasmic reticulum Ca^{2+} -release channel by analogues of sulmazole, 111, 1212
- Calcium sensitivity**, Diosmin-induced increase in sensitivity to Ca^{2+} of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978
- Importance of inositol (1,4,5)-trisphosphate, intracellular Ca^{2+} release and myofilament Ca^{2+} sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
- A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757
- Calcium transients**, Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
- Calcium uptake**, Effects of cyclopiazonic acid on rhythmic contractions in uterine smooth muscle bundles of the rat, 112, 1132
- Calmodulin**, Relaxation of rat thoracic aorta induced by the Ca^{2+} -ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Calmodulin antagonists**, Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469
- Calphostin C**, Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K^+ contracture in rat and guinea-pig aorta, 112, 604
- Cancer**, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113,

757

Capillaries, Effects of selective ET_B -receptor stimulation on arterial, 112, 888

Capillary pressure, Effects of selective ET_B -receptor stimulation on arterial, 112, 888

Capsaicin, Benzodiazepine-induced intestinal motor disturbances in rats: mediation by w_2 (BZ_2) sites on capsaicin-sensitive afferent neurones, 113, 268

- Block by capsaicin of voltage-gated K^+ currents in melanotrophs of the rat pituitary, 112, 616
- Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547
- Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017
- Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29
- Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703
- Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
- Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
- Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895

Captopril, Facilitation by procterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781

- Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310

Carbachol, All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972

- Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
- A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593
- Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351
- Variable potency of nitrergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494

Carbonic anhydrase inhibitors, The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881

5-Carboxamidotryptamine (5-CT), Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

Cardiac action potential, Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Cardiac asthma, Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649

Cardiac automaticity, Evidence for a cooperation between adenosine A_2 receptors and β -adrenoceptors on cardiac automaticity in the isolated right ventricle of the rat, 111, 1316

Cardiac conduction, The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69

Cardiac contractions, Comparison of the effects of caffeine and other methylxanthines on $[\text{Ca}^{2+}]_i$ in rat ventricular myocytes, 111, 455

- Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

Cardiac hypertrophy, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

- Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Cardiac inotropism, Effects of chronic infusions of α -trinisol on regional and cardiac haemodynamics in conscious rats, 113, 129

Cardiac muscles, Comparison of the effects of caffeine and other methylxanthines on $[\text{Ca}^{2+}]_i$ in rat ventricular myocytes, 111, 455

Cardiac pacing, Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081

Cardiac transplantation, Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137

Cardioprotection, A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757

Cardiovascular regulation, central, Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250

Cardiovascular system, Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

Carotid occlusion, Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443

Cartilage, Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Castor oil, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127

Catechol, Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154

Catecholamines, Antiarrhythmic effects of BN-063, a newly synthesized adenosine A_1 agonist, on myocardial ischaemia in rats, 112, 1031

- The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311
- Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154

Catecholamine overflow, Sustained prejunctional facilitation of noradrenergic neurotransmission by adrenaline as a co-transmitter in the portal vein of freely moving rats, 113, 342

Catecholamine secretion, Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935

Catechol O -methyl transferase inhibition, Comparison of two new inhibitors of catechol O -methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

Cationic channels, non-selective, Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942

Cationic polypeptides, Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29

Cations, divalent, Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P_2 purinoceptor classification, 113, 463

Caudate, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

L-CCG-I, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

- Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364

CD11/CD18, Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123

CD18, Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117

- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- cDNA**, Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377
- Celiprolol**, Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043
- Cell adhesion**, The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311
- Cell culture**, A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST₈₈, 111, 1135
- Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050
- Cell growth**, Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
- Cell proliferation**, Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
- Cell proliferation, smooth muscle**, Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- β -Cells**, Mg²⁺-dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632
- Cephalosporins**, Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050
- Cerebellar granule cells**, Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Cerebellar Purkinje neurones**, Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906
- Cerebellum**, Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Cerebellum, guinea-pig**, Natriuretic peptide-induced cyclic GMP accumulation in adult guinea-pig cerebellar slices, 113, 216
- Cerebral cortex**, Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226
- Cerebral cortex, guinea-pig**, A comparison of A₂ adenosine receptor-induced cyclic AMP generation in cerebral cortex and relaxation of pre-contracted aorta, 111, 185
- Cerebral cortex, rat**, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
- Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355
- Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231
- Cerebrospinal fluid**, Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449
- C-fibres**, Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310
- Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
- CGP 28014**, Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13
- (\pm)-CGP 12177, Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699
- CGS 15943**, Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- CGS 21680**, Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741
- Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
- CHAPS**, Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630
- Charybdotoxin**, Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233
- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Cheek pouch, hamster**, Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Chemotactic factor**, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Chewing**, The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839
- Chick embryos**, Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
- Chimeric peptides**, Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Chimeric receptors**, Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331
- Chlorethylclonidine**, Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- Chloride channels**, Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169
- Chloride channel blockers**, Effects of Cl channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333
- Chloride currents**, Activation of chloride current by P₂-purinoceptors in rat ventricular myocytes, 111, 1355
- Permeation properties and modulation of volume-activated Cl⁻ currents in human endothelial cells, 112, 1049
- Role of cyclic AMP-induced Cl⁻ conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137
- Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555
- Chloride currents, calcium-activated**, Effects of Cl channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333
- Chloride secretion**, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31
- Chloride transport**, IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179
- Chlorisondamine**, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414
- Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Regulation of nicotinic receptors in rat brain following quasi-irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917
- Chlormethiazole**, Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188
- A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131
- 2-Chloro-ATP**, Characterization of the effects of 2-methylthio-ATP and 2-chloro-ATP on brain capillary endothelial cells: similarities to ADP and differences from ATP, 112, 775

- The P_{2X} -purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946
- Chloroethylclonidine**, α_{1b} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819
- Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- CHO cells**, Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911
- Pharmacological characterization of the human histamine H_2 receptor stably expressed in Chinese hamster ovary cells, 112, 847
- Cholecystokinin**, Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429
- Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037
- Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930
- Cholecystokinin receptors**, Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltrindole, 111, 956
- Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429
- Evaluation of a series of novel CCK_B antagonists using a functional assay in the rat central nervous system, 112, 666
- Cholecystokinin receptor antagonists**, Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltrindole, 111, 956
- Cholecystokinin CCK_B receptor antagonists**, Evaluation of a series of novel CCK_B antagonists using a functional assay in the rat central nervous system, 112, 666
- [3 H]-**Choline release**, Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895
- Cholinergic neurones**, Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789
- Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085
- Cholinergic neurotransmission**, Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
- Cholinergic stimulation**, Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550
- Cholinergic synapses**, The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898
- Cholinesterase inhibitors**, Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695
- The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747
- Chondromalacia patellae**, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435
- Chorioallantoic membranes, chick**, Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
- Chromaffin cells**, Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- Cibacron blue**, Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117
- The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α , β -methylene adenosine 5'-triphosphate, 112, 282
- Pharmacological profile of the ATP-mediated increase in L-calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Cicaprost**, Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369
- Cilia**, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the cilium toxin, pyocyanin, 112, 493
- Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469
- Ciliotoxin**, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the cilium toxin, pyocyanin, 112, 493
- Cinanserin**, Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
- Cinnamophilin**, Pharmacological characterization of cinnamophilin, a novel dual inhibitor of thromboxane synthase and thromboxane A₂ receptor, 111, 906
- Cirazoline**, The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997
- The effects of perfusion rate and N^G-nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
- Circulation, bronchial**, Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649
- Circulation, coronary**, Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869
- Circulation, hepatic**, The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Circulation, splanchnic**, Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209
- Cirrhosis**, Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209
- Cisplatin**, Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431
- Citrulline metabolism**, Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487
- Citrulline transport**, Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487
- Clofilium**, Antifibrillatory effects of clofilium in the rabbit isolated heart, 113, 209
- Cloned receptors**, Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
- Pharmacological properties of the cloned $\alpha_{1A/D}$ -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral cortex and vas deferens, 111, 1003
- Clonidine**, Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
- Cloriciromene**, Inhibition of human monocyte adhesion to endothelial cells by the coumarin derivative, cloriciromene, 111, 575
- Clostridium histolyticum**, Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
- Clotrimazole**, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- Clozapine**, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873
- The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839
- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769
- CNQX**, Cyclothiazide unmasks AMPA-evoked stimulation of [3 H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- Cobra venom factor**, Pro-inflammatory activities in elapid snake venoms, 112, 723
- Cocaine**, Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
- Codeine**, Moguisteine: a novel peripheral non-narcotic antitussive drug, 112, 795
- Collagen**, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662
- Collagenase**, Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
- Collagenous protein biosynthesis**, Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894

Collecting duct cells, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na^+ and K^+ transport properties in the rabbit cortical collecting duct, 111, 173

Colon, Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077

Colon, cat, Contractions mediated by α_1 -adrenoceptors and P_2 -purinoceptors in a cat colon circular muscle, 112, 1237

Colon, guinea-pig, Comparison of tachykinin NK_1 and NK_2 receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150

- The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789
- Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259
- Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088

Colon, human, Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143

Colon, proximal, Nitric oxide-mediated inhibitory response of rat proximal colon: independence from changes in membrane potential, 112, 676

Colon, rat, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

- Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930

Complement, Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117

Complement activation, Pro-inflammatory activities in elapid snake venoms, 112, 723

Compound 48/80, Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35

Conductance, Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873

Congenital heart disease, Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245

Conjunctiva, Blockade by oral or parenteral RPR 100893 (a non-peptide NK_1 receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

ω -Conotoxin, Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077

- Depression of high-threshold calcium currents by activation of human D₂ (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061
- Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
- Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103
- Muscarinic (M₁) receptor-mediated inhibition of K⁺-evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L- and N-type Ca²⁺ channels, 113, 621

Contractile apparatus, Diosmin-induced increase in sensitivity to Ca²⁺ of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Contraction, Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103

Converting enzyme inhibitor, The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567

Convulsions, Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89

Coronary ligation, Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837

Coronary microcirculation, Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548

Corpus cavernosum, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81

Cortex, olfactory, NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221

Cortex, rat, Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043

- Pharmacological properties of the cloned $\alpha_{1A/D}$ -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral cortex and vas deferens, 111, 1003

Cortical neurones, Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379

Corticotropin-releasing factor, Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930

COS-M6 cells, Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Co-transmission, Blockade by 4,4'-diisothiocyanostilbene-2,2'-disulphonate (DIDS) of P_{2X}-purinoceptors in rat vas deferens, 112, 690

- Blockade of P_{2X}-purinoceptors by trypan blue in rat vas deferens, 113, 349
- Investigation of the actions of PPADS, a novel P_{2X}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913

Cotransport, Triflocin, a novel inhibitor for the Na-HCO₃ symport in the proximal tubule, 112, 465

Cough, Mogulisteine: a novel peripheral non-narcotic antitussive drug, 112, 795

- Peripheral and central sites of action of GABA_B agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344

CP-96,345, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
- (±)-CP-96,345, Differences in the effects of NK₁-receptor antagonists, (±)-CP 96,345 and CP 99,994, on agonist-induced responses in guinea-pig trachea, 112, 176
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

CP-99,994, Differences in the effects of NK₁-receptor antagonists, (±)-CP 96,345 and CP 99,994, on agonist-induced responses in guinea-pig trachea, 112, 176

D-CPPen, Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073

Cromakalim, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

- Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278
- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
- Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525

D-Current, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

CV1808, Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741

CV-3988, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123

Cyanide, Effects of metabolic blockers on Ca^{2+} -dependent currents in cultured sensory neurones from neonatal rats, 111, 57

Cyclic AMP, A_1 adenosine receptor inhibition of cyclic AMP formation and radioligand binding in the guinea-pig cerebral cortex, 113, 1501

- Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7
- Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775
- Characterization of prostanoid receptors on rat neutrophils, 113, 581
- Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866
- A comparison of A_2 adenosine receptor-induced cyclic AMP generation in cerebral cortex and relaxation of pre-contracted aorta, 111, 185
- Differential effects of P_2 -purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y} -purinoceptors, 113, 614
- The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
- Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493
- Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625
- Electrophysiological subtypes of inhibitory P_1 purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703
- Endogenous expression of histamine H_1 receptors functionally coupled to phosphoinositide hydrolysis in C6 glioma cells: regulation by cyclic AMP, 113, 1554
- Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1
- Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396
- Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
- Molecular cloning and expression of human EP_3 receptors: evidence of three variants with differing carboxyl termini, 112, 377
- Ontogenetic increase in PGE_2 and $\text{PGF}_{2\alpha}$ receptor density in brain microvessels of pigs, 112, 59
- Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271
- Pharmacological characterization of the human histamine H_2 receptor stably expressed in Chinese hamster ovary cells, 112, 847
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047
- Role of cyclic AMP-induced C1 conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Studies on the mechanism of [^3H]-noradrenaline release from SH-SY5Y cells: the role of CA^{2+} and cyclic AMP, 111, 787

Cyclic AMP accumulation, Adenosine A_{2B} -receptor-mediated cyclic AMP accumulation in primary rat astrocytes, 111, 191

Cyclic AMP-dependent protein kinase, Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625

- Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509

Cyclic AMP phosphodiesterase, Separation and characterization of a novel isoenzyme of cyclic nucleotide phosphodiesterase from rat cerebrum, 111, 389

Cyclic GMP, Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5

- Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071
- Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053
- Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867
- Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
- Loss of preconditioning in rabbits with vascular tolerance to nitroglycerin, 112, 999
- Natriuretic peptide-induced cyclic GMP accumulation in adult guinea-pig cerebellar slices, 113, 216

- Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047
- Relaxation of rat thoracic aorta induced by the Ca^{2+} -ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445

Cyclic nucleotide-gated channels, Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469

N⁶-Cyclohexyl-adenosine, Modulation of vasodilatation to levromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49

Cyclo-oxygenase, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272

- Cyclo-oxygenase and nitric oxide synthase isoforms in rat carageenan-induced pleurisy, 113, 693
- The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Relative roles of nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid in the contractile responses of rat renal arcuate arteries, 112, 369

Cyclopentyladenosine, Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A_2 -but not to an A_1 -agonist in the dog, 111, 1233

8-Cyclopentyl-1,3-dipropylxanthine, Modulation of vasodilatation to levromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49

Cyclophosphamide, Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017

- The effect of leukocyte depletion on pulmonary M_2 muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588

Cyclopiazonic acid, Effect of cyclopiazonic acid, an inhibitor of sarcoplasmic reticulum Ca^{2+} -ATPase, on the frequency-dependence of the contraction-relaxation cycle of the guinea-pig isolated atrium, 113, 1001

- Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669
- Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233
- Effects of cyclopiazonic acid on rhythmic contractions in uterine smooth muscle bundles of the rat, 112, 1132
- Inhibitory effects of nordihydroguaiaretic acid on ET_{A} -receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- Relaxation of rat thoracic aorta induced by the Ca^{2+} -ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Variable potency of nitrergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494

D-Cycloserine, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

Cyclothiazide, Cyclothiazide unmasks AMPA-evoked stimulation of [^3H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339

Cysteine, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825

L-Cysteine, Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287

Cystic fibrosis, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31

- Stimulation of chloride secretion by P_1 purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169
- Cystitis**, Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017
- Cystometry**, Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
- Cytisine**, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- Cytochrome P450**, Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548
- Cytokines**, Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Inhibition of nitric oxide synthesis in vascular smooth muscles by retinoids, 113, 1448
- The involvement of bradykinin B_1 and B_2 receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153
- Cytoskeletal proteins**, Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271
- Cytotoxicity**, Reduction by N^G -nitro-L-arginine of H_2O_2 -induced endothelial cell injury, 113, 564

D

- DAU 6215**, Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695
- DAU 6285**, Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213
- DDT₁MF-2 cells**, (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
- Defibrillation**, Antifibrillatory effects of clofilium in the rabbit isolated heart, 113, 209
- Dehydration**, Effect of dehydration and hyperosmolar hydration on lignocaine and metabolites disposition in conscious rabbits, 113, 317
- Delayed rectifier**, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411
- Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972
- Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
- Dendrotoxins**, Blockade by dendrotoxin homologues of voltage-dependent K⁺ currents in cultured sensory neurones from neonatal rats, 113, 959
- 2-Deoxyglucose**, Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57
- Dependence**, Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Depolarization**, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca²⁺ current in guinea-pig heart, 113, 451
- Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P_2 purinoceptor classification, 113, 463
- Des-Arg⁹-bradykinin**, Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- [³H]-Des-Arg⁹-kallidin, Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389
- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- Desensitization**, Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541

- Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β_1 - and β_2 -receptor-mediated responses, 112, 595
- Rapid desensitization of adrenaline- and neuropeptide Y-stimulated Ca²⁺ mobilization in HEL-cells, 112, 499
- Desipramine**, Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625
- Detrusor muscle**, Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
- Developmental change**, Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569
- Dexamethasone**, Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083
- Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents, 113, 1416
- Dexamethasone-induced translocation of lipocortin (annexin) 1 to the cell membrane of U-937 cells, 112, 347
- Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111
- Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1
- Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449
- Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480
- Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081
- A role for endogenous histamine in interleukin-1 β -induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801
- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240
- Dextrin 2-sulphate**, Infection by HIV-1 blocked by binding of dextrin 2-sulphate to the cell surface of activated human peripheral blood mononuclear cells and cultured T-cells, 113, 151
- 2',5'-Di(tert-butyl)-1,4-benzohydroquinone (BHQ), Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118
- 1,3-Di(2-tolyl)guanidine (DTG), The effects of sigma ligands and of neuropeptide Y on N-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709
- Diabetes mellitus**, Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370
- Beneficial effect of the Ca²⁺ antagonist, nifedipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887
- Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
- The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42
- Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195
- Insulin requirement for the antihyperglycaemic effect of metformin, 111, 793
- Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35
- Diacylglycerol**, The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Diaphragm**, Regulation of baseline vascular resistance in the canine

diaphragm by nitric oxide, 112, 65

Diaphragm, rat, Potentiation by tonic A_{2a} -adenosine receptor activation of CGRP-facilitated [3 H]-ACh release from rat motor nerve endings, 111, 582

Diarrhoea, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127

Diazepam, The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497

- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170

Dicentrine, The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69

DIDS, Blockade by 4,4'-diisothiocyanostilbene-2,2'-disulphonate (DIDS) of P_{2x} -purinoceptors in rat vas deferens, 112, 690

- Effects of Cl channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333

Dihydroergotamine, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105

Dihydro- β -erythroidine, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

L-threo-3,4-Dihydroxyphenylserine, The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503

Dihydropyridines, Comparison of the haemodynamic profiles of elgopidine and nicardipine in the anaesthetized dog, 111, 49

- The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- Pharmacology of high-threshold calcium currents in GH₄C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728
- Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903

Diisopropylfluorophosphate (DFP), Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695

Dilatation, neurogenic, Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630

Diltiazem, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca²⁺ current in guinea-pig heart, 113, 451

- Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989

Dimethylarginase, Metabolism of methylarginines by human vasculature; implications for the regulation of nitric oxide synthesis, 112, 43

N,N-Di-n-propyl-5-carboxamidotryptamine (DP-5-CT), Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

- Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318

Diosmin, Diosmin-induced increase in sensitivity to Ca²⁺ of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Diphenylbutylpiperidine, Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells, 111, 483

1,3-Dipropyl-cyclopentylxanthine (DPCPX), Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

Dipyridamole, Studies of the nucleoside transporter inhibitor, draflazine, in the human myocardium, 112, 137

(-)-Discretamine, (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174

Disopyramide, Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873

Disseminated intravascular coagulation (DIC), The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289

Dissociation constants, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

- pK₁ values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

Distribution, Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355

Diuresis, Central and peripheral actions of the novel κ -opioid receptor agonist, EMD 60400, 111, 837

Dizocilpine, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

- Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073
- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89
- A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

DMBAPTA, Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753

DMP728, Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

DMPP, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

DNA synthesis, Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195

DOPAC, Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

- The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417

L-DOPA, Cell inward transport of L-DOPA and 3-O-methyl-L-DOPA in rat renal tubules, 112, 611

- Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13
- Regional haemodynamic effects of dopamine and its prodrugs L-dopa and gludopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417

Dopamine, Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695

- Cell inward transport of L-DOPA and 3-O-methyl-L-DOPA in rat renal tubules, 112, 611
- Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829
- Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837
- The effects of administration of monoamine oxidase-B inhibitors on rat striatal neurone responses to dopamine, 113, 1159
- The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- Effect of type A and B monoamine oxidase selective inhibition by Ro

41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

- Pharmacology of high-threshold calcium currents in GH₄C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728
- Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
- Regional haemodynamic effects of dopamine and its prodrugs L-dopa and glucodopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417
- Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930

Dopamine, extracellular, Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073

Dopamine metabolism, Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

Dopamine receptors, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

- Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23

Dopamine receptor agonists, Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23

Dopamine receptor antagonists, Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23

Dopamine D₁ receptors, Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β ₁- and β ₂-receptor-mediated responses, 112, 595

- The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839

Dopamine D₂ autoreceptors, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

Dopamine D₂ receptors, Depression of high-threshold calcium currents by activation of human D₂ (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

- The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839

Dopamine D₃ receptors, Functional expression of human D₃ dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391

Dopamine D₄ receptors, Pharmacology of high-threshold calcium currents in GH₄C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728

Dopamine release, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

- Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Modulation of dopamine and noradrenaline release and of intracellular Ca²⁺ concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
- Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
- Role of 5-HT₁ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

Dopamine turnover, The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the N-methyl-D-aspartate (NMDA) receptor complex, 113, 729

- Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

Dopamine uptake, Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391

Dopexamine, Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199

- Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β ₁- and β ₂-receptor-mediated responses, 112, 595

L-threo-DOPS, The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α -adrenoceptors in the mouse, 111, 503

Dorsal raphe nucleus, Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429

8-OH-DPAT, The effect of 5-HT_{1A} receptor agonists on locomotor activity in the guinea-pig, 112, 861

[³H]-DPCPX, The binding of 1,3-[³H]-dipropyl-8-cyclopentylxanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249

DPMA, Ischaemia/reperfusion selectively attenuates coronary vasodilation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233

Draflazine, Studies of the nucleoside transporter inhibitor, draflazine, in the human myocardium, 112, 137

Drinking, Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328

Ductus arteriosus, Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245

Duodenum, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G-nitro-L-arginine methyl ester, 113, 1127

Duodenum, rat, Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191

DuP 996, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103

Dura mater, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

E

Econazole, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

Ectothiopate, The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747

Ecstasy, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlorimethiazole, dizocilpine and other protective compounds, 111, 131

Ecto-ATPase, Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669

- Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2X}-purinoceptor ligands, 113, 1432

Efaroxan, The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065

Eicosanoids, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926
- Modulation by endogenous prostanooids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675

Electrical field stimulation, Loss of contractile activity of endothelin-1 induced by electrical field of stimulation-generated free radicals, 113, 21

- Selective antagonism by PPADS at P_{2X}-purinoceptors in rabbit isolated blood vessels, 111, 923

Electrical stimulation, The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363

Electrolyte excretion, Natriuretic action of angiotensin(1-7), 111, 1

Electromechanical coupling, Bradykinin B₂ receptors and coupling

- mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- Electrophysiology**, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873
- Evaluation of a series of novel CCK_B antagonists using a functional assay in the rat central nervous system, 112, 666
- Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na⁺ and K⁺ transport properties in the rabbit cortical collecting duct, 111, 173
- Elevated plus-maze test**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
- Elgodipine**, Comparison of the haemodynamic profiles of elgodipine and nicardipine in the anaesthetized dog, 111, 49
- EMD 60400**, Central and peripheral actions of the novel κ-opioid receptor agonist, EMD 60400, 111, 837
- EMD 61753**, A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 61753, 113, 1317
- Emesis**, Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431
- Emopamil**, Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Enadoline**, Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002
- Enalapril**, Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837
- Endopeptidases**, Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Neutral endopeptidase (NEP) inhibition in rats with established pulmonary hypertension secondary to chronic hypoxia, 113, 1121
- Role of endopeptidase 3.4.24.16 in the catabolism of neuropeptides, *in vivo* in the vascularly perfused dog ileum, 112, 127
- Endopeptidases, neutral**, Hydrolysis of iodine labelled urodilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204
- Endothelial cells**, Agonist-induced desensitization of histamine H₁ receptor-mediated inositol phospholipid hydrolysis in human umbilical vein endothelial cells, 113, 823
- Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831
- Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y}-purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
- Inhibition of human monocyte adhesion to endothelial cells by the coumarin derivative, cloricromene, 111, 575
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α₁-adrenoceptor stimulation, 113, 1203
- Metabolism of methylarginines by human vasculature; implications for the regulation of nitric oxide synthesis, 112, 43
- Permeation properties and modulation of volume-activated Cl⁻ currents in human endothelial cells, 112, 1049
- Reduction by N^ω-nitro-L-arginine of H₂O₂-induced endothelial cell injury, 113, 564
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041
- Endothelial injury**, Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309
- Endothelins**, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257
- Characterization of endothelin (ET) receptors in the isolated gall bladder of the guinea-pig: evidence for an additional ET receptor subtype, 112, 1244
- Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- Effects of selective ET_B-receptor stimulation on arterial, 112, 888
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, platelet-activating factor and thromboxane A₂, 112, 963
- Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature, 112, 704
- Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109
- [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093
- Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041
- Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49
- Endothelin converting enzymes**, Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Endothelin ET_A receptors**, [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
- Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009
- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845
- Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
- Effects of an ET₁-receptor antagonist, FR139317, on regional haemodynamic responses to endothelin-1 and [Ala^{11,15}]Ac-endothelin-1 (6-21) in conscious rats, 112, 477
- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- The effects of the endothelin ET_A receptor antagonist, FR 139317, on infarct size in a rabbit model of acute myocardial ischaemia and reperfusion, 112, 75
- Effects of selective ET_B-receptor stimulation on arterial, 112, 888
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963
- Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336
- Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49
- Endothelin ET_A receptor antagonists**, Block of Endothelin-1-induced

release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009
- The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
- Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515
- Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336

Endothelin ET_B receptors, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845
- Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- Effects of an ET_i-receptor antagonist, FR139317, on regional haemodynamic responses to endothelin-1 and [Ala^{11,15}]Ac-endothelin-1 (6-21) in conscious rats, 112, 477
- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- Effects of selective ET_B-receptor stimulation on arterial, 112, 888
- Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336
- Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49

Endothelin ET_B receptor agonists, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

Endothelin ET_B receptor antagonists, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336

Endothelin receptors, Characterization of endothelin (ET) receptors in the isolated gall bladder of the guinea-pig: evidence for an additional ET receptor subtype, 112, 1244

- Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729
- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207

Endothelin receptor subtypes, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336

Endothelin receptor agonists, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

Endothelin receptor antagonists, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515
- Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207
- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869

Endothelin-1, Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247

- Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009
- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845
- Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- The effects of the endothelin ET_A receptor antagonist, FR 139317, on infarct size in a rabbit model of acute myocardial ischaemia and reperfusion, 112, 75
- Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
- ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515
- Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- The involvement of endothelial dysfunction, nitric oxide and prostanoids in the rat gastric microcirculatory responses to endothelin-1, 112, 267
- Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473
- Loss of contractile activity of endothelin-1 induced by electrical field of stimulation-generated free radicals, 113, 21
- Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163
- Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
- Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311
- Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207
- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869
- Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery, 113, 336

Endothelin-1, big, Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471

- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823

- ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729
- Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Endothelin-3**, Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
- Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163
- Endothelium**, Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370
- Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550
- Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801
- The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311
- Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71
- Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925
- ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729
- Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557
- Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233
- Lack of importance of NO in β -adrenoceptor-mediated relaxation of large epicardial canine coronary arteries, 111, 981
- Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77
- Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351
- Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533
- Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
- Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat, 111, 1184
- The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275
- Relaxation of human isolated mesenteric arteries by vasopressin and desmopressin, 113, 419
- Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiroxanic acid, possibly through nitric oxide formation, 111, 655
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
- Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630
- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869
- The role of myoendothelial cell contact in non-nitric oxide-, non-prostanoid-mediated endothelium-dependent relaxation of porcine coronary artery, 113, 1289
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471
- Endothelium, vascular**, Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponarrestat, 111, 42
- Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35
- Endothelium-dependent contractions**, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
- Endothelium-dependent relaxation**, The effects of aldose reductase inhibition with ponarrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Endothelium-dependent vasodilatation**, Recovery after dietary vitamin E supplementation of impaired endothelial function in vitamin E-deficient rats, 112, 515
- Endothelium-derived contracting factor**, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
- Mediation by M₃-muscarinic receptors of both endothelium-dependent contraction and relaxation to acetylcholine in the aorta of the spontaneously hypertensive rat, 112, 519
- Endothelium-derived hyperpolarizing factor**, Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831
- Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053
- Display of the characteristics of endothelium-derived hyperpolarizing factor by a cytochrome P450-derived arachidonic acid metabolite in the coronary microcirculation, 113, 1548
- Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
- The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275
- The role of myoendothelial cell contact in non-nitric oxide-, non-prostanoid-mediated endothelium-dependent relaxation of porcine coronary artery, 113, 1289
- Endothelium-derived relaxing factor**, Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolaemic rabbits, 113, 190
- Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053
- The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
- Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponarrestat, 111, 42
- Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
- Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77
- Mediation by M₃-muscarinic receptors of both endothelium-dependent contraction and relaxation to acetylcholine in the aorta of the spontaneously hypertensive rat, 112, 519
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- The role of myoendothelial cell contact in non-nitric oxide-, non-prostanoid-mediated endothelium-dependent relaxation of porcine coronary artery, 113, 1289
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567
- Endotoxin**, Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary

epithelial cells: regulation by dexamethasone, 113, 1008

- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101
- Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309

Endplate, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Energy metabolism, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

Enflurane, The effects of propofol and enflurane on single calcium channel currents of guinea-pig isolated ventricular myocytes, 111, 1147

Enkephalins, Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817

- Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Entacapone, Comparison of two new inhibitors of catechol *O*-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

Enteric nervous system, Electrophysiological subtypes of inhibitory P_1 purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703

- Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095
- L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205
- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161

Enteric reflexes, Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095

Enterotoxin, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31

Eosinophils, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

- Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205
- Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to CSa in guinea-pig skin, 113, 117
- Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35
- Long-lasting inhibitory activity of the hetazeptinic BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994
- Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
- Role of interleukin-5 in enhanced migration of eosinophils from airways of immunized guinea-pigs, 113, 749

Epibatidine, Antinociceptive and toxic effects of (+)-epibatidine oxalate attributable to nicotinic agonist activity, 113, 1487

Epidermal growth factor, Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

Epilepsy, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

- Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents, 113, 1416
- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y₂ receptors in rat hippocampal slices *in vitro*, 113, 737

Epileptiform activity, The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316

Epithelium, Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29

- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83
- Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

Epithelium, airways, Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEO-, 112, 169

- Stimulation by menthol of C1 secretion via a Ca²⁺-dependent mechanism in canine airway epithelium, 112, 571

Epithelium, ciliary, Role of cyclic AMP-induced C1 conductance in aqueous humour formation by the dog ciliary epithelium, 112, 1137

Epithelium, colonic, Calcitonin gene-related peptide receptors in human gastrointestinal epithelia, 113, 1243

- Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31

Epithelium, pulmonary, Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008

Erythrocytes, Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903

B-Escin, Diosmin-induced increase in sensitivity to Ca²⁺ of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Ethanol, Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089

Ethanol withdrawal, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017

N-Ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

Evans blue, Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

Evoked potentials, NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221

Excitation-contraction coupling, Ca²⁺ release induced by myotoxin *a*, a radio-labellable probe having novel Ca²⁺ release properties in sarcoplasmic reticulum, 113, 233

Excitatory amino acids, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

- Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188
- A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843
- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97
- Intracerebral microdialysis combined with recording of extracellular field potential: a novel method for investigation of depolarizing drugs *in vivo*, 113, 1295

Excitatory amino acid receptors, A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843

- Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-

- glutamate release from rat hippocampal synaptosomes, 113, 339
- Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364
- Excitatory postsynaptic potentials**, Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083
- Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703
- Exercise**, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231
- EXP3174**, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179
- Exploration test**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170

F

- Facilitation**, Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296
- Fatty acids**, Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509
- 8FB-PTP**, Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- Felodipine**, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017
- Fetus**, Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Fever**, Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449
- Feverfew**, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9
- Fibrates**, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551
- Fibrinogen binding**, Comparative pharmacology of analogues of S-nitroso-N-acetyl-DL-penicillamine on human platelets, 112, 1071
- Fibrinogen receptors**, Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333
- Fibrinolysis**, The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289
- Fibroblasts**, Binding and growth-inhibitory effect of heparin and oligoheparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
- Fibronectin**, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662
- Fish oil**, Effects of dietary marine oil supplementation on reactivity of human buttock subcutaneous arteries and forearm veins *in vitro*, 112, 566
- Fissistigma glaucescens**, Antimuscarinic action of liriodenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle, 113, 1464
- (-)-Discretamine, a selective α_{1D} -adrenoceptor antagonist, isolated from *Fissistigma glaucescens*, 112, 1174
- Pharmacological characteristics of liriodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275
- FK224**, 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183
- FK409**, Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137
- Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385
- Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5

- FK888**, Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183
- Flecainide**, Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873
- Flesinoxan**, Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318
- The renal functional responses to 5-HT_{1A} receptor agonist, flesinoxan, in anaesthetized, normotensive rat, 112, 214
- Flow cytometry**, Dexamethasone-induced translocation of lipocortin (annexin) 1 to the cell membrane of U-937 cells, 112, 347
- Flumazenil**, Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- (S)- α -Fluoromethylhistidine**, Role of histamine in rodent antinociception, 111, 1269
- Fluorosulphonylbenzoyladenosine**, The P_{2Z}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946
- Fluoxetine**, Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741
- Flurbiprofen**, Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983
- Fluspirilene**, Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells, 111, 483
- Formoterol**, The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687
- Forskolin**, Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625
- Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
- Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Modulation of cardiac L- Ca²⁺ channels by GTPyS in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- FPL 66096**, FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2T}-purinoceptors, 113, 1057
- FPL 66301**, Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2X}-purinoceptor ligands, 113, 1432
- FR139317**, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009
- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845
- Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251
- Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386
- Effects of an ET₁-receptor antagonist, FR139317, on regional haemodynamic responses to endothelin-1 and [Ala^{11,15}]Ac-endothelin-1 (6-21) in conscious rats, 112, 477
- The effects of the endothelin ET_A receptor antagonist, FR 139317, on infarct size in a rabbit model of acute myocardial ischaemia and reperfusion, 112, 75
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963

- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4
- [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303
- Free radicals**, The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801
- Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431
- Frequency dependent block**, Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717
- Fructose 1,6-diphosphate**, Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57
- Frusemide**, Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918
- Fura-2**, Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252
- Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989

G

- GABA**, Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188
- Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303
- Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555
- GABA receptor ligands**, Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
- GABA release**, GABA and glutamate release affected by GABA_B receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
- Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
- GABA_A receptors**, Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
- Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906
- Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555
- GABA_A receptor subunits**, Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025
- GABA_B receptors**, GABA and glutamate release affected by GABA_B receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
- Paired-pulse depression of the N-methyl-D-aspartate receptor-mediated synaptic potentials in the amygdala, 113, 1029
- Peripheral and central sites of action of GABA_B agonists to inhibit the cough reflex in the cat and guinea-pig, 113, 1344
- Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_B receptor system, 113, 1561
- GABA_B receptor antagonists**, GABA and glutamate release affected by GABA_B receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
- Galantin**, Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Gallamine**, The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-

- pigs, 112, 588
- Gall bladder**, Characterization of endothelin (ET) receptors in the isolated gall bladder of the guinea-pig: evidence for an additional ET receptor subtype, 112, 1244
- Ganglia, cardiac**, Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Ganglia, dorsal root**, Blockade by dendrotoxin homologues of voltage-dependent K⁺ currents in cultured sensory neurones from neonatal rats, 113, 959
- Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
- Ganglia, superior cervical, rat**, Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357
- Ganglion blockers**, The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Gardos channels**, Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903
- Gastric acid secretion**, Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93
- Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87
- Gastric fundus, rat**, Influence of S-nitrosothiols and nitrate tolerance in the rat gastric fundus, 111, 1280
- Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541
- Gastric fundus strips**, Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089
- Gastric mucosa**, Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
- Gastrointestinal motility**, Stimulating action of KW-5139 (Leu¹³-motilin) on gastrointestinal motility in the rabbit, 111, 288
- Gastrointestinal tract**, Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845
- Gastrointestinal transit**, Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228
- GBR 12909**, Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
- Geller-Seifter test**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
- Gemeprost**, Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369
- Gene expression, AT₁**, receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662
- General anaesthetics**, The effects of propofol and enflurane on single calcium channel currents of guinea-pig isolated ventricular myocytes, 111, 1147
- Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906
- Geniculate nucleus, dorsolateral**, The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97
- Genistein**, Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- Gentamicin**, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375
- GH₃ cells**, Lithium-induced decrease in spontaneous Ca²⁺ oscillations in single GH₃ rat pituitary cells, 112, 390
- GH₄C₁ cells**, Pharmacology of high-threshold calcium currents in GH₄C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728
- Glial cells**, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, oxamido adipate, 113, 536
- Glial fibrillary acidic protein (GFAP)**, The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997
- Glibenclamide**, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411
- Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit

- arterial myocytes, 111, 972
- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649
- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N^G -monomethyl-L-arginine but not glibenclamide, 111, 711
- Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588
- Glibenclamide analogues**, Location of the sulphonylurea receptor at the cytoplasmic face of the β -cell membrane, 113, 903
- Glioma cells**, Differential effects of P_2 -purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y} -purinoceptors, 113, 614
- Glipizide**, Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649
- Globus pallidus**, Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
- Glomerular filtration rate**, Natriuretic action of angiotensin(1-7), 111, 1
- Glucagon**, Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
- Glucocorticoids**, Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D₃, 113, 439
- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240
- Glucose**, Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
- NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
- Glucose metabolism**, Insulin requirement for the antihyperglycaemic effect of metformin, 111, 793
- Glucose utilization**, Importance of the intestine as a site of metformin-stimulated glucose utilization, 112, 671
- GluDOPA**, Regional haemodynamic effects of dopamine and its prodrugs L-dopa and glucdopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417
- Glutamate**, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
- Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247
- Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188
- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303
- Intracerebral microdialysis combined with recording of extracellular field potential: a novel method for investigation of depolarizing drugs *in vivo*, 113, 1295
- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y₂ receptors in rat hippocampal slices *in vitro*, 113, 737
- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347
- Glutamate receptors**, Modulation of dopamine and noradrenaline release and of intracellular Ca^{2+} concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- Glutamate receptor heterogeneity**, Species-dependent functional properties of non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with mammalian and avian brain mRNA, 111, 803
- Glutamate release**, Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- GABA and glutamate release affected by GABA_A receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
- Glutamate transport**, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, α amino adipate, 113, 536
- L-Glutamine**, Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- Glutamine synthetase**, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, α amino adipate, 113, 536
- γ -Glutamylcysteine synthetase**, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, α amino adipate, 113, 536
- Glutathione**, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825
- Glyburide**, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411
- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levocromakalim, 113, 889
- Glycerol**, Regional haemodynamic effects of dopamine and its prodrugs L-dopa and glucdopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- Glyceryl trinitrate**, Nitric oxide formation from glyceryl trinitrate by rabbit aortic strip: detection by rabbit taenia coli concurrent with vasorelaxation, 113, 1508
- Glycine**, Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- A novel antagonist, phenylbenzene α phosphono- α -amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord, 113, 165
- Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951
- Glycine receptors**, The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the *N*-methyl-D-aspartate (NMDA) receptor complex, 113, 729
- Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- Glycine receptor antagonists**, A novel antagonist, phenylbenzene α phosphono- α -amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord, 113, 165
- Gomisin C**, Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
- G-proteins**, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852
- Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
- Characterization of the binding of the first selective radiolabelled histamine H₁-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355
- The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
- Effect of a peptide inhibitor of protein kinase C on G-protein-mediated increase in myofilament Ca^{2+} -sensitivity in rabbit arterial skinned muscle, 111, 311
- Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
- The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussis toxin, 112, 709
- 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
- Modulation of cardiac L- Ca^{2+} channels by GTP γ S in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
- Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377
- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55
- Sensitivity of G-protein involved in endothelin-1-induced Ca^{2+} influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989

- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240
- G-protein coupling**, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295
- [³H]-GR65630, Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541
- GR79236**, Characterization of the adenosine receptors mediating hypothermia in the conscious mouse, 113, 1386
- Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₃ receptor, 113, 741
- GR82334**, Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985
- GR113808**, Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1
- Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213
- GR113808: a novel, selective antagonist with high affinity at the 5-HT₄ receptor, 111, 332
- GR117289**, Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137
- Gramine**, Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951
- Granisetron**, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968
- Granule cells, cerebellar**, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
- Grooming**, The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839
- p-Guanidinobenzoic acid**, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na⁺ and K⁺ transport properties in the rabbit cortical collecting duct, 111, 173
- Guanine nucleotides**, Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355
- Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71
- Guanylin**, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31
- Guanylyl cyclase**, Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569
- GYKI**, Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- GYKI 52466**, Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- GYKI 53655**, A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843

H

- H7**, Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237
- Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604
- HA-966**, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Haemodynamics**, Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
- Comparison of the haemodynamic profiles of elgiodipine and nicardipine in the anaesthetized dog, 111, 49
- Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111
- Effects of an ET₁-receptor antagonist, FR139317, on regional haemodynamic responses to endothelin-1 and [Ala^{11,15}]Ac-endothelin-1 (6-21) in conscious rats, 112, 477
- Effects of bosentan (Ro 47-0203), an ET_A-, ET_B-receptor antagonist, on regional haemodynamic responses to endothelins in conscious rats, 112, 823
- Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- Effects of chronic infusions of α-trinisolol on regional and cardiac haemodynamics in conscious rats, 113, 129
- Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649
- Effects of N^G-nitro-L-arginine methyl ester on regional haemodynamic responses to MgSO₄ in conscious rats, 111, 325
- Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
- Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂ but not to an A₁-agonist in the dog, 111, 1233
- Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209
- Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347
- Regional haemodynamic effects of dopamine and its prodrugs L-dopa and glucopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589
- The renal functional responses to 5-HT_{1A} receptor agonist, flesinoxan, in anaesthetized, normotensive rat, 112, 214
- Haemolytic activity**, Pro-inflammatory activities in elapid snake venoms, 112, 723
- Haemorrhage**, Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117
- Haloperidol**, The effects of sigma ligands and of neuropeptide Y on N-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709
- Heart**, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca²⁺ current in guinea-pig heart, 113, 451
 - Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529
- Heart, isolated**, Antifibrillatory effects of clofium in the rabbit isolated heart, 113, 209
- Heart, rabbit**, The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69
 - Loss of preconditioning in rabbits with vascular tolerance to nitroglycerin, 112, 999
 - Potentiation by ACE inhibitors of the dilator response to bradykinin in the coronary microcirculation: interaction at the receptor level, 111, 238
- Heart, rat**, α₁-Adrenoceptors in the conduction system of rat hearts, 111, 465
 - Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775
 - Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963
 - Functional studies on α₁-adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539
 - Radioligand binding studies of α₁-adrenoceptor subtypes in rat heart, 111, 533
 - Role of K⁺ channels in the vasodilator response to bradykinin in the rat heart, 113, 954
- Heart contractility**, Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271
- Heart failure**, Effects of the calcium channel blockers, diltiazem and

Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395

Heart rate, Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

- Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199
- Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064

HeLa cells, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598

Hemidiaphragm, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Heparin, Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

- Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
- Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29
- The effect of allosteric antagonists in modulating muscarinic M_2 -receptor function in guinea-pig isolated trachea, 112, 901

Heptanitrosyl-tri- μ 3-thioxotetraferrate (1-), Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Hernandezine, Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767

12-HETE, Effect of cytoplasmic pH on Ca^{2+} -stimulated eicosanoid biosynthesis in human platelets, 113, 926

Hexahydrosiladiphenol, Muscarinic (M_1) receptor-mediated inhibition of K^+ -evoked [3H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of *L*- and *N*-type Ca^{2+} channels, 113, 621

Hexamethonium, The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

Hexacyclium/sila-hexacyclium derivatives, Binding and functional properties of hexacyclium and sila-hexacyclium derivatives to muscarinic receptor subtypes, 112, 505

High pressure neurological syndrome, Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951

Hind limbs, Importance of the intestine as a site of metformin-stimulated glucose utilization, 112, 671

Hippocampal neurones, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Hippocampal slices, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y_2 receptors in rat hippocampal slices *in vitro*, 113, 737

Hippocampus, Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083

- Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318
- Differential sensitivity of 3H -agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400
- The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316
- Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commisural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521
- Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695
- Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226
- Modulation of dopamine and noradrenaline release and of intracellular Ca^{2+} concentration by presynaptic glutamate receptors in hippocampus, 113, 1439

- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y_2 receptors in rat hippocampal slices *in vitro*, 113, 737
- **Histamine**, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
- Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H_3 receptors in the trachea, 111, 445
- Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93
- Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159
- Differential effects of Na^+, K^+ -ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87
- Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918
- Investigation of the endogenous chemoattractants involved in ^{111}In -eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35
- Involvement of apamin-sensitive K^+ channels in antigen-induced spasms of guinea-pig isolated trachea, 112, 958
- Novel histamine H_3 receptor antagonists: affinities in an H_3 receptor binding assay and potencies in two functional H_3 receptor models, 112, 1043
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- A role for endogenous histamine in interleukin-8-induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801
- Role of histamine in rodent antinociception, 111, 1269
- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041
- Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101
- **Histamine H₁ receptors**, Agonist-induced desensitization of histamine H₁ receptor-mediated inositol phospholipid hydrolysis in human umbilical vein endothelial cells, 113, 823
- Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
- Endogenous expression of histamine H₁ receptors functionally coupled to phosphoinositide hydrolysis in C6 glioma cells: regulation by cyclic AMP, 113, 1554
- Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262
- **Histamine H₂ receptors**, Pharmacological characterization of the human histamine H₂ receptor stably expressed in Chinese hamster ovary cells, 112, 847
- **Histamine H₃ receptors**, Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [^{125}I]-iodophenpropit, to rat brain, 113, 355
- Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H_3 receptors in the trachea, 111, 445
- **Histamine H₃ receptor subtypes**, Novel histamine H₃ receptor antagonists: affinities in an H_3 receptor binding assay and potencies in two functional H_3 receptor models, 112, 1043
- **Histamine H₃ receptor antagonists**, Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [^{125}I]-iodophenpropit, to rat brain, 113, 355
- **Histamine secretion**, IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179
- **L-Histidine**, Role of histamine in rodent antinociception, 111, 1269
- **HIV-1**, Infection by HIV-1 blocked by binding of dextrin 2-sulphate to the cell surface of activated human peripheral blood mononuclear cells and cultured T-cells, 113, 151
- **HL-60 cells**, Dual effects of tetrandrine on cytosolic calcium in human

leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767

Hoe-140, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683

- Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004
- Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167
- Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717
- Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81
- Reduction by Hoe 140, the B₂ kinin receptor antagonist, of antigen-induced nasal blockage, 111, 969
- Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036
- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Homeostasis, Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57

HPLC, Hydrolysis of iodine labelled urodilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204

- Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625
- Pharmacokinetics, plasma protein binding and urinary excretion of N^o-nitro-L-arginine in rats, 111, 394

5-HT, Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083

- Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711
- Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370
- Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105
- Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1
- Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240
- Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143
- Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
- Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541
- Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318
- Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400
- Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695
- The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789
- Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
- Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213
- GR113808: a novel, selective antagonist with high affinity at the 5-HT₄ receptor, 111, 332
- The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13
- 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
- Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
- Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431

- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358
- Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095
- *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939
- A pharmacological analysis of receptors mediating the excitatory response to 5-hydroxytryptamine in the guinea-pig isolated trachea, 112, 763
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093
- Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline mesenteric circulation, 111, 264
- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968
- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471
- Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673

5-HT autoreceptors, Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

5-HT receptors, Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095

- Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939
- A pharmacological analysis of receptors mediating the excitatory response to 5-hydroxytryptamine in the guinea-pig isolated trachea, 112, 763

5-HT receptor agonists, Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358

5-HT receptor antagonists, A pharmacological analysis of receptors mediating the excitatory response to 5-hydroxytryptamine in the guinea-pig isolated trachea, 112, 763

[³H]-5-HT release, Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

5-HT reuptake blockade, Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

5-HT turnover, Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

5-HT₁ mechanisms, Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358

5-HT_{1A} receptors, Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083

- Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
- Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318
- Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400
- Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

5-HT_{1A} receptor agonists, The effect of 5-HT_{1A} receptor agonists on locomotor activity in the guinea-pig, 112, 861

- The renal functional responses to 5-HT_{1A} receptor agonist, flesinoxan,

in anaesthetized, normotensive rat, 112, 214

5-HT_{1D} receptors, The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13

- Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939

5-HT₂ receptors, Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240

- On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77
- Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline mesenteric circulation, 111, 264

5-HT₂ receptor antagonists, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272

5-HT_{2A} receptors, *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797

5-HT_{2B} receptors, Further characterization of 5-hydroxytryptamine receptors (putative 5-HT_{2B}) in rat stomach fundus longitudinal muscle, 112, 323

- *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797

5-HT_{2C} receptors, *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797

5-HT₃ receptors, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

- Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240
- Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541
- Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741
- Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695
- 5-HT₃ receptors do not mediate vagally-induced relaxation or contraction of the isolated stomach of the guinea-pig, 111, 346
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745
- Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485
- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

5-HT₃ receptor antagonists, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527

- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

5-HT₄ receptors, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

- Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1
- Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143
- The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789
- Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213
- GR113808: a novel, selective antagonist with high affinity at the 5-HT₄ receptor, 111, 332
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77

5-HT₄ receptor antagonists, The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789

- GR113808: a novel, selective antagonist with high affinity at the 5-HT₄ receptor, 111, 332

Hyaluronan, Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Hydrogen peroxide, Reduction by *N*^G-nitro-L-arginine of H₂O₂-induced endothelial cell injury, 113, 564

Hydroquinone, Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287

8-Hydroxy-2-(di-n-propylamino)tetralin (8-OH-DPAT), Adaptive changes in 5-HT_{1A} receptor-mediated hippocampal inhibition in the alert rat produced by repeated 8-OH-DPAT treatment, 112, 1083

- Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
- Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318
- Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400

5-Hydroxydecanoate, Antifibrillatory effects of clofibrate in the rabbit isolated heart, 113, 209

5-Hydroxyindole, Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541

Hydroxyl radicals, Loss of contractile activity of endothelin-1 induced by electrical field of stimulation-generated free radicals, 113, 21

Hyperaemia, cutaneous, Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681

Hyperaemia, gastric, Role of bradykinin in the hyperaemia following acid challenge of the rat gastric mucosa, 113, 1036

Hyperalgesia, Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735

- The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- A pharmacological profile of the novel, peripherally-selective κ -opioid receptor agonist, EMD 61753, 113, 1317
- Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_A receptor system, 113, 1561

Hypercholesterolaemia, Abolition of flow-dependent EDRF release before that evoked by agonists in hypercholesterolaemic rabbits, 113, 190

- Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035

Hyperglycaemia, The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801

Hyperkalaemia, The dose-related hyper- and hypokalaemic effects of salbutamol and its arrhythmic potential, 111, 73

Hyperosmolality, Effect of dehydration and hyperosmolar hydration on lignocaine and metabolites disposition in conscious rabbits, 113, 317

- Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604

Hyperpolarization, Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557

- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525

Hypersensitivity, Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230

- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Hypertension, The effects of perfusion rate and *N*^G-nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13

- Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067
- Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925
- Role of Ca²⁺-dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022

Hypertension, genetic, Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111

Hypertension, portal, Study on the vascular reactivity and α -adrenoceptors of portal hypertensive rats, 111, 439

Hypertension, pulmonary, Neutral endopeptidase (NEP) inhibition in rats with established pulmonary hypertension secondary to chronic hypoxia, 113, 1121

Hypertension, renal, Attenuated renal response to moxonidine and rilmenidine in one kidney-one clip hypertensive rats, 112, 200

Hypertensive rat models, Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137

- Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717
- Role of Ca²⁺-dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022

Hypokalaemia, The dose-related hyper-and-hypokalaemic effects of salbutamol and its arrhythmogenic potential, 111, 73

Hypotension, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A₂ receptor, 113, 741

Hypothalamus, Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

- Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619, 113, 659
- Benzodiazepine/cholecystokinin interactions at functional CCK receptors in rat brain, 112, 429
- Evaluation of a series of novel CCK_B antagonists using a functional assay in the rat central nervous system, 112, 666
- Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347

Hypothermia, Characterization of the adenosine receptors mediating hypothermia in the conscious mouse, 113, 1386

- Covariation of α₂-adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855
- The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352

Hypoxia, Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N^G-monomethyl-L-arginine but not glibenclamide, 111, 711

- Modulation of vasodilatation to levocromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Neutral endopeptidase (NEP) inhibition in rats with established pulmonary hypertension secondary to chronic hypoxia, 113, 1121
- Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311

I

IB4, Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123

Iberiotoxin, Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958

Ibdilast, Possible role of cyclic AMP phosphodiesterases in the actions of ibdilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081

ICAM-1, Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912

Icatibant, Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004

- Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167
- Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453

ICI 118551, Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199

- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589

ICI 197067, Central and peripheral actions of the novel κ-opioid receptor agonist, EMD 60400, 111, 837

- A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 61753, 113, 1317
- A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 60400, 111, 837
- A pharmacological profile of the novel, peripherally-selective κ-opioid receptor agonist, EMD 61753, 113, 1317

ICI 215001, β₁- and β₂-adrenoceptor antagonist activities of ICI-215001, a putative β₃-adrenoceptor agonist, 112, 55

Idazoxan, The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997

- Modulation by central postsynaptic α₂-adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140
- pK_i values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

IgE, Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

IgE antibodies, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

IgE receptors, IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179

IgG1 antibodies, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

Ileocolonic junction, Effect of potassium channel blockade and α₂-adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341

Ileum, Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₂ receptor, 112, 461

Ileum, dog, Role of endopeptidase 3.4.24.16 in the catabolism of neurotensin, *in vivo* in the vascularly perfused dog ileum, 112, 127

Ileum, guinea-pig, β₁- and β₂-adrenoceptor antagonist activities of ICI-215001, a putative β₃-adrenoceptor agonist, 112, 55

- Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240
- Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77
- Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043
- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161

Imidazoles, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

Imidazoline binding sites, Attenuated renal response to moxonidine and rilmenidine in one kidney-one clip hypertensive rats, 112, 200

- The effects of chronic imidazoline drug treatment on glial fibrillary acidic protein concentrations in rat brain, 111, 997
- The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065
- Sodium excretion following central administration of an I₁ imidazoline receptor agonist, moxonidine, 112, 1089

Imipramine, Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741

- Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625

Immunoblockade, Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703

Immunomodulation, Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240

Immunosuppression, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Indo-1, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261

- Comparison of the effects of caffeine and other methylxanthines on $[Ca^{2+}]_i$ in rat ventricular myocytes, 111, 455
- Indomethacin**, Effects of intrinsic prostaglandins on the spontaneous contractile and electrical activity of the proximal renal pelvis of the guinea-pig, 113, 431
- Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
- Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83
- Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983
- Indorenate**, Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616
- Inflammation, airway**, The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004
- Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423
- Cyclo-oxygenase and nitric oxide synthase isoforms in rat carrageenin-induced pleurisy, 113, 693
- Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513
- Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292
- Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7
- Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Long-lasting inhibitory activity of the tetrazepinic BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994
- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
- A role for endogenous histamine in interleukin-8-induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Inflammation, allergic**, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480
- Inflammation, cutaneous**, Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Inflammation, intestinal**, Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309
- Inflammation, joint**, The role of bradykinin B₁ receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940
- Inflammation, neurogenic**, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920
 - Interleukin-18 enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
 - Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
- Influenza A virus**, ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Inhibitory junction potentials**, Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
- Inhibitory postsynaptic potentials**, Y₂-receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883
- Inositol phosphates**, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
 - Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
 - Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598
 - Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614
 - Differential regulation of inositol 1,4,5-trisphosphate by co-existing P₂-purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
 - Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299
 - The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
 - 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
 - Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
 - Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231
 - Ontogenetic increase in PGE₂ and PGF_{2α} receptor density in brain microvessels of pigs, 112, 59
 - Pharmacological characterization of the human histamine H₂ receptor stably expressed in Chinese hamster ovary cells, 112, 847
- Inositol phospholipid turnover**, A₁ adenosine receptor inhibition of cyclic AMP formation and radioligand binding in the guinea-pig cerebral cortex, 113, 1501
 - Agonist-induced desensitization of histamine H₁ receptor-mediated inositol phospholipid hydrolysis in human umbilical vein endothelial cells, 113, 823
 - Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
 - Endogenous expression of histamine H₁ receptors functionally coupled to phosphoinositide hydrolysis in C6 glioma cells: regulation by cyclic AMP, 113, 1554
 - Participation of cytoskeleton in the effect of antilaminin IgG on cardiac cholinoreceptors, 111, 271
- Inositol 1,4,5-trisphosphate**, Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
 - Lithium-induced decrease in spontaneous Ca²⁺ oscillations in single GH₃ rat pituitary cells, 112, 390
- Inotropic agents**, Cardiovascular actions of dopexamine in anaesthetized and conscious dogs, 111, 199
 - Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539
 - A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757
- Insulin**, Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
 - The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065
 - Insulin requirement for the antihyperglycaemic effect of metformin, 111, 793
 - Location of the sulphonylurea receptor at the cytoplasmic face of the B-cell membrane, 113, 903
 - NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
 - Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35
- Integrins**, Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Integrin receptors**, Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type

plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Intercellular adhesion molecule-1 (ICAM-1), Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912

- Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117

Interferon- γ , Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720

- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Interleukin-1, The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809

Interleukin-1 receptor antagonists, The involvement of bradykinin B₁ and B₂ receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63

Interleukin-1 β , Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1

- Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449
- Inhibition of nitric oxide synthesis in vascular smooth muscles by retinoids, 113, 1448
- Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
- Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389

Interleukin-2, Inhibition by SK&F 96365 of Ca²⁺ current, IL-2 production and activation in T lymphocytes, 113, 861

Interleukin-5, Role of interleukin-5 in enhanced migration of eosinophils from airways of immunized guinea-pigs, 113, 749

Interleukin-8, A role for endogenous histamine in interleukin-8-induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801

Intestinal motility, Benzodiazepine-induced intestinal motor disturbances in rats: mediation by w₂ (BZ₂) sites on capsaicin-sensitive afferent neurones, 113, 268

- L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205

Intestinal oxygen consumption, L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205

Intestinal secretion, Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228

Intestine, Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240

- Differential effects of luminal L-arginine and N^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593
- Importance of the intestine as a site of metformin-stimulated glucose utilization, 112, 671
- Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

Intracellular recording, Species-dependent functional properties of non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with mammalian and avian brain mRNA, 111, 803

Intracerebroventricular injection, Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

Inward rectification, Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873

- Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118

Inward rectifier, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527

[¹²⁵I]-**Iodoaminopentideline,** Pharmacological characterization of the human histamine H₂ receptor stably expressed in Chinese hamster ovary cells, 112, 847

[¹²⁵I]-**Iodomelatonin binding,** Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

[¹²⁵I]-**Iodophenpropit,** Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355

Ion channels, All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972

- The P_{2U}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

Ion channel blockers, Block of human voltage-sensitive Na⁺ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600

Ion currents, Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in NIE-115 neuroblastoma cells, 112, 541

Ion transport, Calcitonin gene-related peptide receptors in human gastrointestinal epithelia, 113, 1243

- Stimulation by menthol of Cl⁻ secretion via a Ca²⁺-dependent mechanism in canine airway epithelium, 112, 571

Ionophoresis, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

Ipratropium bromide, Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918

Iris sphincter, Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342

- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Iris sphincter, rabbit, Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179

IRL 1620, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009
- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

Irreversible antagonism, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9

Ischaemia, Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233

- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

Ischaemia, cardiac, Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B₂ receptors, 113, 1167

- Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083
- Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123
- Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081

Islets of Langerhans, The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065

3-Isobutyl-1-methyl-xanthine, Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259

Isocarbacyclin, Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635

Isoflurane, Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906

Isoprenaline, Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β_1 - and β_2 -receptor-mediated responses, 112, 595

- Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625
- Effects of muscarinic M_2 and M_3 receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- Lack of importance of NO in β -adrenoceptor-mediated relaxation of large epicardial canine coronary arteries, 111, 981
- Modulation of cardiac L -Ca²⁺ channels by GTPyS in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- Isosorbide**, Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137
- Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385
- Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127
- The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328
- Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5
- [³H]-Isradipine binding, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017

J

Jaw-opening reflex, Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140

Jejunum, Adrenoceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576

- Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127
- JO-1784**, The effects of sigma ligands and of neuropeptide Y on N -methyl-D-aspartate-induced neuronal activation of CA_3 , dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709

K

Kainic acid, Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215

- Species-dependent functional properties of non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with mammalian and avian brain mRNA, 111, 803

Kallikrein, The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567

KC 128, Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889

KF 17837, Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659

- Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43

KH 1060, Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D₃, 113, 439

Kidney, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Cell inward transport of L-DOPA and 3-O-methyl-L-DOPA in rat renal tubules, 112, 611

- Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

- Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously

- hypertensive rats, 111, 111

- Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269
- Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1, 113, 931
- The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881
- [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303

Kidney, rat, Natriuretic action of angiotensin(1-7), 111, 1

- Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296
- The renal handling of dopamine originating from L-DOPA and γ -glutamyl-L-DOPA, 112, 417

Kindling, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97**Kinins**, Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508**Kinin receptors**, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81**Kininase II**, The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567**KIZD-06**, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81**KRN2391**, Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278**KW-5139 (Leu¹³-motilin)**, Stimulating action of KW-5139 (Leu¹³-motilin) on gastrointestinal motility in the rabbit, 111, 288**Kyotorphin**, Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817**L****L-362,855**, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628**L-652,731**, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123**L-687,414**, The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the N-methyl-D-aspartate (NMDA) receptor complex, 113, 729**Labyrinthectomy**, A dose-response analysis of the beneficial effects of the ACTH-(4-9) analogue, Org 2766, on behavioural recovery following unilateral labyrinthectomy in guinea-pig, 111, 358**Lactate**, Attenuation by chloramphenicol administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188**Laser-Doppler flowmetry**, Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111**Laudanosine**, Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377**Laxatives**, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127**Nitric oxide** as a mediator of the laxative action of magnesium sulphate, 113, 228**Lesopitron (E-4424)**, Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425**Leucocytes**, Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912**Drug modulation** of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111**Pharmacological analysis** of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123

- Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480
- Leucocytes, polymorphonuclear**, Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Leukotrienes**, Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205
- Metabolic disposition of leukotriene B₄ (LTB₄) and oxidation-resistant analogues of LTB₄ in conscious rabbits, 112, 654
- Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Leukaemia**, Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767
- Leukotrienes**, Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292
- Effect of *in vivo* desensitization to leukotriene B₄ on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
- Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958
- Metabolic disposition of leukotriene B₄ (LTB₄) and oxidation-resistant analogues of LTB₄ in conscious rabbits, 112, 654
- Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101
- Leukotriene analogues**, Metabolic disposition of leukotriene B₄ (LTB₄) and oxidation-resistant analogues of LTB₄ in conscious rabbits, 112, 654
- Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35
- Leukotriene receptors**, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852
- Levcromakalim**, Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levcromakalim, in vascular smooth muscle, 112, 1223
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Modulation of vasodilatation to levcromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levcromakalim, 113, 889
- Levosimendan**, A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757
- Lifarizine**, Block of human voltage-sensitive Na⁺ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600
- Ligand binding**, Characterization of the binding of the first selective radiolabelled histamine H₃-receptor antagonist, [¹²⁵I]-iodophenpropit, to rat brain, 113, 355
- Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389
- Lignocaine**, Block of human voltage-sensitive Na⁺ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600
- Effect of dehydration and hyperosmolal hydration on lignocaine and metabolites disposition in conscious rabbits, 113, 317
- Linopirdine**, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103
- Lipids, plasma**, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551
- Lipocortins**, Dexamethasone-induced translocation of lipocortin (annexin) 1 to the cell membrane of U-937 cells, 112, 347
- A role for endogenous histamine in interleukin-8-induced neutrophil infiltration into mouse air-pouch: investigation of the modulatory action of systemic and local dexamethasone, 112, 801
- Lipopolysaccharides**, Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942
- Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487
- Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111
- Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289
- Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- Long-lasting inhibitory activity of the tetrazepinic BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328
- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
- Lipoxygenase**, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852
- The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360
- Relative roles of nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid in the contractile responses of rat renal arcuate arteries, 112, 369
- Lipoxygenase inhibitors**, Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205
- Liriodenine**, Antimuscarinic action of liriodenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle, 113, 1464
- Pharmacological characteristics of liriodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275
- Lisinopril**, Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555
- Lithium**, Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
- Lithium-induced decrease in spontaneous Ca²⁺ oscillations in single GH₃ rat pituitary cells, 112, 390
- Local anaesthetics**, Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Local anaesthetic actions**, Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179
- Locomotor activity**, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073
- The effect of 5-HT_{1A} receptor agonists on locomotor activity in the guinea-pig, 112, 861
- Regulation of nicotinic receptors in rat brain following quasi-

irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917

Long-term potentiation, Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521

- Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695

Losartan, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

- Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277
- Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589

LP-805, The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173

Lung, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825

- Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230

Lung, guinea-pig, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H₃ receptors in the trachea, 111, 445

Lung, human, Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228

- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists cilepilol, bisoprolol and propranolol on human bronchi, 113, 1043

Lung, rabbit, Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973

Lung, rat, Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

Lymphocytes, Inhibition by SK&F 96365 of Ca²⁺ current, IL-2 production and activation in T lymphocytes, 113, 861

- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240

Lymphocytes, human leukaemic, The P_{2Z}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

M

McNeil A 343, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

Macrophages, Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487

- Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664

Magnesium, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

- Mg²⁺-dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Magnesium sulphate, Effects of N^G-nitro-L-arginine methyl ester on regional haemodynamic responses to MgSO₄ in conscious rats, 111, 325

- Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228

Manganese, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

MAP4, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

Mast cell degranulation, Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958

Mast cells, IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179

- Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342

Mastoparan, Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299

MCCG, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

M cells, A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593

M-current, Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160

- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55

MDL 16,455A, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262

Mecamylamine, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397

Medetomidine, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

- Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951

Megakaryocytes, Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245

Megakaryocytes, rat, Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca²⁺ oscillation, 112, 349

Melanotrophs, Block by capsaicin of voltage-gated K⁺ currents in melanotrophs of the rat pituitary, 112, 616

Melatonin, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Melatonin receptors, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Membrane hyperpolarization, Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165

Membrane ion current, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Membrane potential, Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889
- Role of Ca²⁺-dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022

Membrane translocation, Dexamethasone-induced translocation of lipocortin (annexin) 1 to the cell membrane of U-937 cells, 112, 347

Menthol, Stimulation by menthol of C1 secretion via a Ca²⁺-dependent mechanism in canine airway epithelium, 112, 571

Mepacrine, Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745

Mephenesin, Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951

irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917

Long-term potentiation, Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521

- Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695

Losartan, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

- Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277
- Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589

LP-805, The effects of a novel vasodilator, LP-805, on cytosolic Ca²⁺ concentrations and on tension in rabbit isolated femoral arteries, 113, 1173

Lung, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825

- Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230

Lung, guinea-pig, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H₃ receptors in the trachea, 111, 445

Lung, human, Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228

- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists cilepilol, bisoprolol and propranolol on human bronchi, 113, 1043

Lung, rabbit, Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973

Lung, rat, Developmental changes in ANP-stimulated guanylyl cyclase activity enhanced by ATP in rat lung membrane fractions, 113, 569

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

Lymphocytes, Inhibition by SK&F 96365 of Ca²⁺ current, IL-2 production and activation in T lymphocytes, 113, 861

- Sensitization by dexamethasone of lymphocyte cyclic AMP formation: evidence for increased function of the adenylyl cyclase catalyst, 113, 240

Lymphocytes, human leukaemic, The P_{2Z}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

M

McNeil A 343, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

Macrophages, Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487

- Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664

Magnesium, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

- Mg²⁺-dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Magnesium sulphate, Effects of N^G-nitro-L-arginine methyl ester on regional haemodynamic responses to MgSO₄ in conscious rats, 111, 325

- Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228

Manganese, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

MAP4, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

Mast cell degranulation, Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958

Mast cells, IgE-receptor activated chloride uptake in relation to histamine secretion from rat mast cells, 111, 1179

- Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342

Mastoparan, Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299

MCCG, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

M cells, A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593

M-current, Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160

- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55

MDL 16,455A, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262

Mecamylamine, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406

- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397

Medetomidine, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

- Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951

Megakaryocytes, Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245

Megakaryocytes, rat, Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca²⁺ oscillation, 112, 349

Melanotrophs, Block by capsaicin of voltage-gated K⁺ currents in melanotrophs of the rat pituitary, 112, 616

Melatonin, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Melatonin receptors, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Membrane hyperpolarization, Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165

Membrane ion current, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Membrane potential, Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889
- Role of Ca²⁺-dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022

Membrane translocation, Dexamethasone-induced translocation of lipocortin (annexin) 1 to the cell membrane of U-937 cells, 112, 347

Menthol, Stimulation by menthol of C1 secretion via a Ca²⁺-dependent mechanism in canine airway epithelium, 112, 571

Mepacrine, Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745

Mephenesin, Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951

[³H]-Mepyramine, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262

Mesangial cells, Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7

Mesenteric arterial bed, Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat, 111, 1184

Mesenteric arterial bed, perfused, The effects of perfusion rate and *N*^G-nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13

Mesenteric circulation, Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline mesenteric circulation, 111, 264

- The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275

Metabolic inhibition, Effects of metabolic blockers on Ca²⁺-dependent currents in cultured sensory neurones from neonatal rats, 111, 57

Metabotropic glutamate receptors, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Modulation of cyclic AMP formation by putative metabotropic receptor agonists, 111, 364
- Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231

Metergoline, The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13

Metformin, Importance of the intestine as a site of metformin-stimulated glucose utilization, 112, 671

- Insulin requirement for the antihyperglycaemic effect of metformin, 111, 793

Methoctramine, Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Methoxamine, Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649

- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057

o-Methoxy-sila-hexocyclium, Binding and functional properties of hexocyclium and sila-hexocyclium derivatives to muscarinic receptor subtypes, 112, 505

5-Methoxytryptamine, Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213

3-Methoxytyramine, Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391

α-Methyl amino acids, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

Methylarginines, Metabolism of methylarginines by human vasculature; implications for the regulation of nitric oxide synthesis, 112, 43

MethylDOPA, Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

3-O-MethylDOPA, Cell inward transport of L-DOPA and 3-O-methyl-L-DOPA in rat renal tubules, 112, 611

N-Methyl-D-aspartate, Intracerebral microdialysis combined with recording of extracellular field potential: a novel method for investigation of depolarizing drugs *in vivo*, 113, 1295

N-Methyl-D-aspartate receptors, Paired-pulse depression of the N-methyl-D-aspartate receptor-mediated synaptic potentials in the amygdala, 113, 1029

2-Methyl-5-hydroxytryptamine, Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

1-Methyl-4-phenylpyridinium ion (MPP⁺), Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837

5-Methyl-urapidil, Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318

- Pharmacological evidence of distinct α₁-adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- Radioligand binding studies of α₁-adrenoceptor subtypes in rat heart, 111, 533

Methylene blue, Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328

α,β-Methylene-ATP, Blockade by 4,4'-diisothiocyanostilbene-2,2'-disulphonate (DIDS) of P_{2X}-purinoceptors in rat vas deferens, 112, 690

- Blockade of P_{2X}-purinoceptors by trypan blue in rat vas deferens, 113, 349
- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂ purinoceptor classification, 113, 463
- Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357
- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2T}-purinoceptors, 113, 1057
- Selective antagonism by PPADS at P_{2X}-purinoceptors in rabbit isolated blood vessels, 111, 923
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α,β-methylene adenosine 5'-triphosphate, 112, 282
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015

[³H]-α,β-Methylene-ATP, Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151

3,4-Methylenedioxymethamphetamine, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

Methylhistamine, Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H₃ receptors in the trachea, 111, 445

- Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043
- Role of histamine in rodent antinociception, 111, 1269

2-Methylthio-ATP, Characterization of the effects of 2-methylthio-ATP and 2-chloro-ATP on brain capillary endothelial cells: similarities to ADP and differences from ATP, 112, 775

- Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂ purinoceptor classification, 113, 463
- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2T}-purinoceptors, 113, 1057
- The P_{2X}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946
- Selective antagonism by PPADS at P_{2X}-purinoceptors in rabbit isolated blood vessels, 111, 923
- Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015

Methylxanthines, Adenosine A_{2B}-receptor-mediated cyclic AMP accumulation in primary rat astrocytes, 111, 191

- Comparison of the effects of caffeine and other methylxanthines on [Ca²⁺]_i in rat ventricular myocytes, 111, 455

Methysergide, Comparison of antinociception induced by supraspinally administered L-arginine and kyotorphin, 112, 817

Metoprine, Role of histamine in rodent antinociception, 111, 1269

Metoprolol, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

Mexiletine, Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673

MgADP, Interaction of tolbutamide and cytosolic nucleotides in

controlling the ATP-sensitive K^+ channel in mouse β -cells, 111, 302

MgGDP, Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K^+ channel in mouse β -cells, 111, 302

Miconazole, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

Microcirculation, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105

- Effects of selective ET_1 -receptor stimulation on arterial, 112, 888

Microdialysis, Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188

- Comparison of two new inhibitors of catechol O -methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13

- Effect of acute administration of the 5-HT_{1A} receptor ligand, lesopitron, on rat cortical 5-HT and dopamine turnover, 113, 425

- Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837

- Intracerebral microdialysis combined with recording of extracellular field potential: a novel method for investigation of depolarizing drugs *in vivo*, 113, 1295

- Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625

Microglia, Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942

- Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29

Microsphere-embolism, Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226

Microvascular leakage, Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649

Microvasculature, Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522

Microvessels, Ontogenetic increase in PGE₂ and PGF_{2 α} receptor density in brain microvessels of pigs, 112, 59

Micturition, Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111

Migraine, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

Milrinone, A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757

Miniature endplate currents, Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Miniature endplate potentials, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103

Misoprostol, Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

MK-801, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

- Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247

- Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073

[³H]-MK-801, Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89

Mobilization, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151

Mogulisteine, Mogulisteine: a novel peripheral non-narcotic antitussive drug, 112, 795

Monensin, Contribution of intra- and extracellular Ca^{2+} to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795

Monkey, Cardiovascular effects of GR117289, a novel angiotensin AT₁ receptor antagonist, 111, 137

- Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213

- A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145

Monoamine oxidase inhibitors, The effects of administration of monoamine oxidase-B inhibitors on rat striatal neurone responses to dopamine, 113, 1159

- Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

- Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485

Monoamines, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

Monoclonal antibodies, Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123

Monocyte adhesion, Inhibition of human monocyte adhesion to endothelial cells by the coumarin derivative, cloricromene, 111, 575

***N*^G-Monomethyl-L-arginine (L-NMMA)**, Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609

- Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557

- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147

- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by *N*^G-monomethyl-L-arginine but not glibenclamide, 111, 711

- Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309

Morphine, Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002

- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215

- Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140

- Modulation of morphine antinociception in the mouse by endogenous nitric oxide, 113, 1372

Mossy fibres, Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521

Motilin, Stimulating action of KW-5139 (Leu¹³-motilin) on gastrointestinal motility in the rabbit, 111, 288

Motor endplate, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Motor impairment, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

Motor nerve, Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Motor nerve terminals, Potentiation by tonic A_{2a}-adenosine receptor activation of CGRP-facilitated [³H]-ACh release from rat motor nerve endings, 111, 582

Moxonidine, Attenuated renal response to moxonidine and rilmenidine in one kidney-one clip hypertensive rats, 112, 200

- Sodium excretion following central administration of an I₁ imidazoline receptor agonist, moxonidine, 112, 1089

mRNA, Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396

mRNA detection, Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49

mRNA translation, Species-dependent functional properties of non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with mammalian and avian brain mRNA, 111, 803

Mucociliary transport, Stimulation by menthol of C1 secretion via a Ca^{2+} -dependent mechanism in canine airway epithelium, 112, 571

Mucosa, Differential effects of luminal L-arginine and *N*^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304

Mucosa, airways, β -Adrenoceptors mediate inhibition of [³H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221

Mucosal injury, intestinal, Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute

endotoxaemia, 113, 1131

Mucosal transport, Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240

Mucus secretion, 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

Muscarinic receptors, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

- The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- Mediation by M₃-muscarinic receptors of both endothelium-dependent contraction and relaxation to acetylcholine in the aorta of the spontaneously hypertensive rat, 112, 519
- Muscarinic (M₁) receptor-mediated inhibition of K⁺-evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L- and N-type Ca²⁺ channels, 113, 621
- Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Muscarinic receptor subtypes, Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775

- Binding and functional properties of hexocyclium and sila-hexocyclium derivatives to muscarinic receptor subtypes, 112, 505
- Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179
- Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Muscarinic receptor agonists, Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179

Muscarinic receptor antagonists, Antimuscarinic action of liriiodenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle, 113, 1464

- Binding and functional properties of hexocyclium and sila-hexocyclium derivatives to muscarinic receptor subtypes, 112, 505
- Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179
- Pharmacological characteristics of liriiodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275

Muscarinic M₁ receptors, Antagonistic properties of McNeil-A-343 at 5-HT₄ and 5-HT₃ receptors, 113, 711

- The effect of allosteric antagonists in modulating muscarinic M₂-receptor function in guinea-pig isolated trachea, 112, 901

Muscarinic M₂ receptors, Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Muscarinic M₃ receptors, Calcium-dependence of histamine- and carbachol-induced inositol phosphate formation in human U373 MG astrocytoma cells: comparison with HeLa cells and brain slices, 111, 598

Muscarinic M₄ receptors, Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529

Muscle relaxants, Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

- Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
- *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Mustard oil, Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181

Mutagenesis, Mutations in transmembrane segment VII of the AT₁ receptor differentiate between closely related insurmountable and competitive angiotensin antagonists, 113, 331

Myaesthesia gravis, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Myenteric plexus, Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703

- Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197

Myocardial contractility, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

Myocardial infarction, Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5

- Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123
- The effects of the endothelin ET_A receptor antagonist, FR 139317, on infarct size in a rabbit model of acute myocardial ischaemia and reperfusion, 112, 75
- Role of endogenous endothelin in myocardial and coronary endothelial injury after ischaemia and reperfusion in rats: studies with bosentan, a mixed ET_A-ET_B antagonist, 113, 869

Myocardial ischaemia, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

- Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963
- Loss of preconditioning in rabbits with vascular tolerance to nitroglycerin, 112, 999
- A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757

Myocardium, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527

Myocardium, human, Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137

Myocytes, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527

Myocytes, arterial, Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972

Myocytes, cardiac, The electrophysiological effects of antiarrhythmic potential of a secoaporphine, N-allylsecoboldine, 113, 221

Myocytes, ventricular, Activation of chloride current by P₂-purinoceptors in rat ventricular myocytes, 111, 1355

- Comparison of the effects of caffeine and other methylxanthines on [Ca²⁺]_i in rat ventricular myocytes, 111, 455
- Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873
- The effects of propofol and enflurane on single calcium channel currents of guinea-pig isolated ventricular myocytes, 111, 1147
- Pharmacological profile of the ATP-mediated increase in L-calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982

Myograph, Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579

Myosin phosphorylation, Effect of a peptide inhibitor of protein kinase C on G-protein-mediated increase in myofilament Ca²⁺-sensitivity in rabbit arterial skinned muscle, 111, 311

Myotoxin a, Ca²⁺ release induced by myotoxin a, a radio-labellable probe having novel Ca²⁺ release properties in sarcoplasmic reticulum, 113, 233

Myotubes, Induction of Na⁺/K⁺-ATPase activity by long-term stimulation of nicotinic acetylcholine receptors in C2C12 myotubes, 111, 459

N

N1E-115 mouse neuroblastoma cells, Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541

Na⁺/K⁺ pumps, Induction of Na⁺/K⁺-ATPase activity by long-term stimulation of nicotinic acetylcholine receptors in C2C12 myotubes, 111, 459

NADH-fluorescence, A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757

NADPH oxidase, Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945

NADPH-diaphorase, Noradrenergic-nitrenergic interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403

Nafamostat, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na^+ and K^+ transport properties in the rabbit cortical collecting duct, 111, 173

Naftidrofuryl, Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226

Naloxone, Moguisteine: a novel peripheral non-narcotic antitussive drug, 112, 795

Naltrindole, Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltrindole, 111, 956

NAN-190, Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695

Natriuresis, Sodium excretion following central administration of an I₁ imidazoline receptor agonist, moxonidine, 112, 1089

Natriuretic peptides, Natriuretic peptide-induced cyclic GMP accumulation in adult guinea-pig cerebellar slices, 113, 216

Natriuretic peptide receptors, Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1, 113, 931

- Natriuretic peptide-induced cyclic GMP accumulation in adult guinea-pig cerebellar slices, 113, 216

Natural killer cells, The effects of β -adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium, 113, 1311

Natural products, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9

Naxagolide [(+)-PHNO], Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

NBQX, A comparison of intravenous NBQX and GYKI 53655 as AMPA antagonists in the rat spinal cord, 112, 843

- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

Nedocromil sodium, Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918

Neomycin, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Neostigmine, Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695

Neovascularization, The effect of the ET_A receptor antagonist, FR 139317, on [¹²⁵I]-ET-1 binding to the atherosclerotic human coronary artery, 112, 386

- Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894

Nerve conduction velocity, Beneficial effect of the Ca^{2+} antagonist, nifedipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887

Nerve depolarization, The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α , β -methylene adenosine 5'-triphosphate, 112, 282

Nerves, sensory, Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425

Neuroblastoma cells, Studies on the mechanism of [³H]-noradrenaline release from SH-SY5Y cells: the role of Ca^{2+} and cyclic AMP, 111, 787

Neurokinins, The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77

- Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Neurokinin A, Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

- Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103

Neurokinin B, Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

- Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Neurokinin receptors, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Neurokinin receptor antagonists, Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570

- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Neurokinin NK₁ receptors, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183
- Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Neurokinin NK₁ receptor subtypes, Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Neurokinin NK₁ receptor antagonists, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
- Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Neurokinin NK₂ receptors, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570
- Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

Neurokinin NK₃ receptors, Presence of NK₃-sensitive neurones in different proportions in the medial habenula of guinea-pig, rat and gerbil, 112, 717

Neuroleptics, Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells, 111, 483

- Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769

Neuromodulators, Contribution of intra- and extracellular Ca^{2+} to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795

- Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Neuromuscular junction, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Neuromuscular junction, frog, Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753

Neuromuscular synapses, Potentiation by endogenously released ATP of spontaneous transmitter secretion at developing neuromuscular synapses in *Xenopus* cell cultures, 111, 880

Neuromuscular transmission, Characterization of the 5-HT receptor potentiating neuromuscular cholinergic transmission in strips of human isolated detrusor muscle, 113, 1

- Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375
- Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Neurones, cultured, Effects of metabolic blockers on Ca^{2+} -dependent currents in cultured sensory neurones from neonatal rats, 111, 57

Neurones, sensory, Blockade by dendrotoxin homologues of voltage-dependent K^+ currents in cultured sensory neurones from neonatal rats, 113, 959

- Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283

Neurones, submucous, Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Neuropathy, Beneficial effect of the Ca^{2+} antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887

Neuropeptide K, Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250

Neuropeptide receptor antagonists, The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

Neuropeptide Y, The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA_3 dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709

- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123
- Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y_2 prejunctional receptor, 113, 479
- Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y_2 receptors in rat hippocampal slices *in vitro*, 113, 737
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
- Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231
- Presence of neuropeptide Y in the rat seminal vesicle and its effects on noradrenaline- and nerve-induced contractions, 113, 877
- Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Neuropeptide Y analogues, Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Neuropeptide Y receptors, Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129

- Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y_2 prejunctional receptor, 113, 479
- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y_2 receptors in rat hippocampal slices *in vitro*, 113, 737
- Rapid desensitization of adrenaline- and neuropeptide Y-stimulated Ca^{2+} mobilization in HEL-cells, 112, 499

- Y_2 -receptor-mediated selective inhibition of slow, inhibitory postsynaptic potential in submucous neurones of guinea-pig caecum, 113, 883

Neuropeptide Y receptor agonists, Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y_2 prejunctional receptor, 113, 479

Neuroprotection, Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat, 112, 188

- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

Neurosecretion, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103

Neurotensin, Role of endopeptidase 3.4.24.16 in the catabolism of neurotensin, *in vivo* in the vascularly perfused dog ileum, 112, 127

Neurotensin receptor antagonists, The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352

Neurotensin receptor subtypes, The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352

Neurotoxicity, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

Neurotransmission, Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703

Neurotransmitters, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

- pK_1 values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

Neurotransmitter release, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y_2 receptors in rat hippocampal slices *in vitro*, 113, 737
- Reduction by intracellular calcium chelation of acetylcholine secretion without occluding the effects of adenosine at frog motor nerve endings, 111, 753

Neutral endopeptidase, Hydrolysis of iodine labelled urotilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204

Neutropenia, Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

Neutrophil chemotactic factor, Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480

Neutrophils, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B_4 and platelet-activating factor in human neutrophils, 111, 852

- Characterization of prostanoid receptors on rat neutrophils, 113, 581
- Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513
- Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123
- Effect of *in vivo* desensitization to leukotriene B_4 on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
- Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945
- Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

NG108-15 cells, Depression of high-threshold calcium currents by activation of human D_2 (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

- Functional expression of human D_3 dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391
- Routes of delivery of muscarinic acetylcholine receptors to the plasma membrane in NG108-15 cells, 111, 1023
- Nicardipine**, Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
- Comparison of the haemodynamic profiles of elgodipine and nicardipine in the anaesthetized dog, 111, 49
- Nickel**, Sensitivity of G-protein involved in endothelin-1-induced Ca^{2+} influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989
- Nicotine**, Antinociceptive and toxic effects of (+)-epibatidine oxalate attributable to nicotinic agonist activity, 113, 1487
- Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073
- Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414
- Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine and other nicotinic antagonists administered *in vitro*, 111, 406
- Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695
- Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- Regulation of nicotinic receptors in rat brain following quasi-irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917
- Nicotinic receptors**, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642
- Regulation of nicotinic receptors in rat brain following quasi-irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917
- The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898
- Nicotinic receptor agonists**, Antinociceptive and toxic effects of (+)-epibatidine oxalate attributable to nicotinic agonist activity, 113, 1487
- Nicotinic receptor channels**, Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Nifedipine**, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca^{2+} current in guinea-pig heart, 113, 451
- Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278
- Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- Muscarinic (M_1) receptor-mediated inhibition of K^+ -evoked [3 H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L - and N -type Ca^{2+} channels, 113, 621
- Neuropeptide Y in rat detrusor and its effect on nerve-mediated and acetylcholine-evoked contractions, 113, 95
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093
- Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- Niflumic acid**, Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977
- (+)-Niguldipine**, Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- (-)-Niguldipine**, Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- Nimodipine**, Beneficial effect of the Ca^{2+} antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887
- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831
- Nisoldipine**, Depression of high-threshold calcium currents by activation of human D_2 (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061
- Nitrate**, Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
- Nitrate tolerance**, Influence of *S*-nitrosothiols and nitrate tolerance in the rat gastric fundus, 111, 1280
- Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533
- Nitrendipine**, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017
- Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
- Nitrenergic stimulation**, Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- Nitrenergic transmission**, Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089
- Nitric oxide**, Antianginal effects of FK409, a new spontaneous NO releaser, 113, 1137
- Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247
- Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385
- Attenuation of the antiarrhythmic effects of ischaemic preconditioning by blockade of bradykinin B_2 receptors, 113, 1167
- Attenuation by dexamethasone of endotoxin protection against ischaemia-induced ventricular arrhythmias, 113, 1083
- Block of Endothelin-1-induced release of thromboxane A_2 from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_1 receptor antagonist, BQ-788, 113, 1257
- Close correlation of the cardioprotective effect of FK409, a spontaneous NO releaser, with an increase in plasma cyclic GMP level, 113, 5
- Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071
- Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
- Contribution of ATP and nitric oxide to NANC inhibitory transmission in rat pyloric sphincter, 113, 681
- Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831
- Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550
- Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Cyclo-oxygenase and nitric oxide synthase isoforms in rat carrageenin-induced pleurisy, 113, 693
- Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825
- Differential effects of luminal L-arginine and *N*^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487
- Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of *N*^G-nitro-L-arginine methyl ester, 113, 1127
- Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257
- The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801
- Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 496
- Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521
- The effects of aldose reductase inhibition with ponalrestat on changes

in vascular function in streptozotocin diabetic rats, 113, 576

- Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089
- Effects of *N*^G-nitro-L-arginine methyl ester on regional haemodynamic responses to MgSO₄ in conscious rats, 111, 325
- The effects of perfusion rate and *N*^G-nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
- Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
- Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
- Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
- Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443
- Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557
- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396
- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by *N*^G-monomethyl-L-arginine but not glibenclamide, 111, 711
- Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961
- Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Interdependence of contractile responses of rat small mesenteric arteries on nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid, 112, 360
- The involvement of endothelial dysfunction, nitric oxide and prostanoids in the rat gastric microcirculatory responses to endothelin-1, 112, 267
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- Lack of importance of NO in β -adrenoceptor-mediated relaxation of large epicardial canine coronary arteries, 111, 981
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77
- Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328
- Metabolism of methylarginines by human vasculature; implications for the regulation of nitric oxide synthesis, 112, 43
- Modulation of morphine antinociception in the mouse by endogenous nitric oxide, 113, 1372
- Modulation of vasodilatation to levocromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205
- Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425
- Nitric oxide formation from glyceryl trinitrate by rabbit aortic strip: detection by rabbit taenia coli concurrent with vasorelaxation, 113, 1508
- Nitric oxide-mediated inhibitory response of rat proximal colon: independence from changes in membrane potential, 112, 676
- Nitric oxide as a mediator of the laxative action of magnesium sulphate, 113, 228
- Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197
- Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83
- NO-dependent and -independent elevation of plasma levels of insulin and glucose in rats by L-arginine, 113, 345
- Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287
- Noradrenergic-nitroergic interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403
- Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334
- Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153
- Pharmacokinetics, plasma protein binding and urinary excretion of *N*^o-nitro-L-arginine in rats, 111, 394
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87
- Prevention by dexamethasone of the marked antiarrhythmic effects of preconditioning induced 20 h after rapid cardiac pacing, 113, 1081
- Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- The relative importance of nitric oxide and nitric oxide-independent mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275
- Relative roles of nitric oxide and cyclo-oxygenase and lipoxygenase products of arachidonic acid in the contractile responses of rat renal arcuate arteries, 112, 369
- Relaxation of human isolated mesenteric arteries by vasopressin and desmopressin, 113, 419
- Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
- Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Role of R-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309
- The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

Nitric oxide donors, photosensitive, Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Nitric oxide gas, Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 1508

113, 496

Nitric oxide inhibitors, Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83

Nitric oxide production *in vivo*, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825

Nitric oxide release, Effect of potassium channel blockade and α_2 -adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341

Nitric oxide synthase, Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282

- The effects of perfusion rate and N^G -nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
- Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine, 112, 720
- Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1
- Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396
- Inhibition of nitric oxide synthesis in vascular smooth muscles by retinoids, 113, 1448
- Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Nitric oxide synthase, inducible, Effect of anti-fungal imidazoles in mRNA levels and enzyme activity of inducible nitric oxide synthase, 111, 1257

- Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 496
- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

Nitric oxide synthase inhibition, Effect of nitric oxide synthase inhibition on long-term potentiation at associational-commissural and mossy fibre synapses on CA3 pyramidal neurones, 111, 521

- The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289

Nitric oxide synthase inhibitors, Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282

- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567
- Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309

Nitrite, Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550

- Discrimination between citrulline and arginine transport in activated murine macrophages: inefficient synthesis of NO from recycling of citrulline to arginine, 112, 487
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961

Nitro blue tetrazolium, Noradrenergic-nitric interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403

7-Nitro indazole, Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282

N^G -Nitro-L-arginine (L-NOARG), Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370

- Differential effects of luminal L-arginine and N^G -nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837
- Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443
- Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557
- Noradrenergic-nitric interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403
- Pharmacokinetics, plasma protein binding and urinary excretion of N^G -nitro-L-arginine in rats, 111, 394
- Reduction by N^G -nitro-L-arginine of H_2O_2 -induced endothelial cell injury, 113, 564
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- Requirement for endothelium-derived nitric oxide in vasodilatation produced by stimulation of cholinergic nerves in rat hindquarters, 112, 630

N^G -Nitro-L-arginine benzyl ester, Reduction by N^G -nitro-L-arginine of H_2O_2 -induced endothelial cell injury, 113, 564

N^G -Nitro-L-arginine methyl ester (L-NAME), Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247

- Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257
- Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
- Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159
- Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of N^G -nitro-L-arginine methyl ester, 113, 1127
- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
- Effects of N^G -nitro-L-arginine methyl ester on regional haemodynamic responses to MgSO₄ in conscious rats, 111, 325
- The effects of perfusion rate and N^G -nitro-L-arginine methyl ester on cirazoline- and KCl-induced responses in the perfused mesenteric arterial bed of rats, 111, 13
- Evidence for reduction of bradykinin-induced bronchoconstriction in guinea-pigs by release of nitric oxide, 113, 1147
- Inhibition of nitric oxide biosynthesis and carotid arteriovenous anastomotic shunting in the pig, 111, 961
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328
- Modulation of vasodilatation to levocromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205
- Nitric oxide-dependent release of vasodilator quantities of calcitonin gene-related peptide from capsaicin-sensitive nerves in rabbit skin, 111, 425
- Reduction by N^G -nitro-L-arginine of H_2O_2 -induced endothelial cell injury, 113, 564
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- The relative importance of nitric oxide and nitric oxide-independent

- mechanisms in acetylcholine-evoked dilatation of the rat mesenteric bed, 113, 1275
- Role of *R*-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309
- The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Nitroglycerin**, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825
- Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278
- Loss of preconditioning in rabbits with vascular tolerance to nitroglycerin, 112, 999
- L-NAME, nitric oxide and jejunal motility, blood flow and oxygen uptake in dogs, 111, 205
- Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334
- Nitroprusside**, Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- S-Nitroso-*N*-acetyl-DL-penicillamine analogues**, Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071
- S-Nitroso-L-cysteine**, Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287
- Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088
- S-Nitrosoglutathione**, Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287
- S-Nitrosothiols**, Influence of *S*-nitrosothiols and nitrate tolerance in the rat gastric fundus, 111, 1280
- Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287
- Nitrovasodilators**, Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533
- Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- NMDA**, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349
- The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709
- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89
- Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951
- NMDA receptors**, The anticonvulsant and behavioural profile of L-687,414, a partial agonist acting at the glycine modulatory site on the *N*-methyl-D-aspartate (NMDA) receptor complex, 113, 729
- Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97
- Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002
- Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231
- The pharmacology of the nicotinic antagonist, chlorisondamine, investigated in rat brain and autonomic ganglion, 111, 397
- NMDA receptor antagonists**, Anti-ischaemic efficacy of a nitric oxide synthase inhibitor and a *N*-methyl-D-aspartate receptor antagonist in models of transient and permanent focal cerebral ischaemia, 113, 247
- Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073
- NMDA antagonists increase recovery of evoked potentials from slices of rat olfactory cortex after anoxia, 111, 1221
- Possible involvement of NMDA receptor-mediated transmission in barbiturate physical dependence, 111, 89
- Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951
- Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347
- NMR spectroscopy**, The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881
- Nociception, formalin-induced**, Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
- Nociceptive modulation**, Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
- Nodose ganglion**, Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741
- Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745
- Nomifensine**, Regional effects of amphetamine, cocaine, nomifensine and GBR 12909 on the dynamics of dopamine release and metabolism in the rat brain, 113, 1391
- Non-adrenergic non-cholinergic nerves**, Influence of *S*-nitrosothiols and nitrate tolerance in the rat gastric fundus, 111, 1280
- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287
- Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541
- Non-adrenergic non-cholinergic neurotransmission**, Contribution of ATP and nitric oxide to NANC inhibitory transmission in rat pyloric sphincter, 113, 681
- Effect of potassium channel blockade and α_2 -adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341
- Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
- Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473
- Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y₂ prejunctional receptor, 113, 479
- Nitric oxide-mediated inhibitory response of rat proximal colon: independence from changes in membrane potential, 112, 676
- Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197
- Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088
- Non-selective cationic channels**, Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29
- Pharmacological profile of the ATP-mediated increase in *L*-type calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Non-steroidal anti-inflammatory drugs**, The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113
- Noradrenaline**, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105
- Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

- Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067
- Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443
- *In vitro* studies of release of adenine nucleotides and adenosine from rat vascular endothelium in response to α_1 -adrenoceptor stimulation, 113, 1203
- Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951
- Modulation by the endothelium of sympathetic vasoconstriction in an *in vitro* preparation of the rat tail artery, 111, 351
- Presence of neuropeptide Y in the rat seminal vesicle and its effects on noradrenaline- and nerve-induced contractions, 113, 877
- Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat, 111, 1184
- Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439
- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363
- Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673
- Noradrenaline-ATP cotransmission**, A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- Noradrenaline-induced contractions**, Heterogeneity of α_1 -adrenoceptor subtypes involved in adrenergic contractions of dog blood vessels, 112, 1167
- Noradrenaline precursors**, The noradrenaline precursor *L-threo*-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α_1 -adrenoceptors in the mouse, 111, 503
- Noradrenaline release**, Contribution of intra- and extracellular Ca²⁺ to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
 - Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625
 - Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867
 - Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
 - Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
 - Modulation of dopamine and noradrenaline release and of intracellular Ca²⁺ concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
 - Muscarinic (M₁) receptor-mediated inhibition of K⁺-evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of *L*- and *N*-type Ca²⁺ channels, 113, 621
 - Noradrenergic-nitroergic interactions in the rat anococcygeus muscle: evidence for postjunctional modulation by nitric oxide, 112, 403
 - Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043
- [³H]-Noradrenaline release**, Modulation of 5-HT release in the guinea-pig brain following long-term administration of antidepressant drugs, 113, 485
 - Studies on the mechanism of [³H]-noradrenaline release from SH-SY5Y cells: the role of CA²⁺ and cyclic AMP, 111, 787
- Noradrenergic mechanisms**, Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109
- Nordihydroguaiaretic acid**, Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- NPC 567**, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683
- NS 1619**, Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619, 113, 659
 - Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Nucleoside transporters**, Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137
- Nucleoside uptake inhibition**, Studies of the nucleoside transporter inhibitor, drafazine, in the human myocardium, 112, 137
- Nucleotides**, Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219
- Nucleotide receptors**, Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y}-purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
 - A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST₄₅, 111, 1135
 - Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455
- Nucleus accumbens**, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

O

- Obesity**, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231
- Oedema**, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513
 - Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111
 - Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205
 - Effect of duration of ischaemia on reduction of myocardial infarct size by inhibition of neutrophil accumulation using an anti-CD18 monoclonal antibody, 111, 1123
 - Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649
 - Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453
 - Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
 - Role of adhesion glycoproteins CD18 and intercellular adhesion molecule-1 in complement-mediated reactions of rabbit skin, 111, 117
 - Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811
 - Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230
 - Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
 - Olfaction**, Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469
 - Olfactory bulb**, Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775
 - Olfactory tubercle**, Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968
 - Oligo-heparin**, Binding and growth-inhibitory effect of heparin and oligo-heparin (2 kDa) in Balb/c 3T3 cells: lack of effect on PDGF- or serum-induced inositol lipid turnover, 113, 254
 - Omeprazole**, Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87
 - Ondansetron**, Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
 - Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968
 - Open channel block**, Block by capsaicin of voltage-gated K⁺ currents in melanotrophs of the rat pituitary, 112, 616
 - Opioids**, Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents,

113, 1416

- Effect of activity at metabotropic, as well as ionotropic (NMDA), glutamate receptors on morphine dependence, 113, 1215
- Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789
- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

κ -Opioid agonists, Central and peripheral actions of the novel κ -opioid receptor agonist, EMD 60400, 111, 837

- A pharmacological profile of the novel, peripherally-selective κ -opioid receptor agonist, EMD 61753, 113, 1317

Opioid receptors, Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789

- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Opioid receptor agonists, Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002

Opioid δ -receptors, Antidepressant-like effects of CCK_B antagonists in mice: antagonism by naltrexone, 111, 956

- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Opioid δ -receptor agonists, Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077

Opioid κ -receptors, Hyperpolarization by opioids acting on μ -receptors of a sub-population of rat periaqueductal gray neurones *in vitro*, 113, 121

Opioid μ -receptors, Dexamethasone-induced selective inhibition of the central μ opioid receptor: functional *in vivo* and *in vitro* evidence in rodents, 113, 1416

- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Org 2766, A dose-response analysis of the beneficial effects of the ACTH-(4-9) analogue, Org 2766, on behavioural recovery following unilateral labyrinthectomy in guinea-pig, 111, 358

Osteoarthritis, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

- Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Ouabain, Contribution of intra- and extracellular Ca²⁺ to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795

- Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87
- Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067

β -Oxidation, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551

Oxidized ATP, The P_{2 U} -purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946

Oxytocin, Effects of cyclopiazonic acid on rhythmic contractions in uterine smooth muscle bundles of the rat, 112, 1132

P

pA₂ values, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219

- pK₁ values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat

isolated vas deferens, 111, 227

Pain, Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004

- Effect of topical administration of L-arginine on formalin-induced nociception in the mouse: a dual role of peripherally formed NO in pain modulation, 112, 547
- Role of histamine in rodent antinociception, 111, 1269

Pain, biliary, Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547

Pain, inflammatory, Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002

Pain reflexes, Blood pressure reflexes following activation of capsaicin-sensitive afferent neurones in the biliopancreatic duct of rats, 111, 547

Paired-pulse depression, Paired-pulse depression of the N-methyl-D-aspartate receptor-mediated synaptic potentials in the amygdala, 113, 1029

Palmitoyl-CoA hydrolase, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551

Palmitoyl-CoA synthetase, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551

Pancreas, Down-regulation of bombesin binding to guinea-pig pancreatic acinar cells during homologous desensitization, 112, 1037

Pancreatic β -cells, Block of ATP-sensitive K⁺ channels in isolated mouse pancreatic β -cells by 2,3-butanedione monoxime, 112, 143

- Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K⁺ channel in mouse β -cells, 111, 302
- Location of the sulphonylurea receptor at the cytoplasmic face of the β -cell membrane, 113, 903
- Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903

Pancuronium, Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283

Papaverine, The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769

- Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489

Papillary muscle, Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca²⁺ current in guinea-pig heart, 113, 451

- Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Parasympathetic nerves, The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588

- Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895

Parasympathetic neurones, Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663

Parasympathetic neurotransmission, Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y₂ prejunctional receptor, 113, 479

Paraventricular nucleus, Prevention by NMDA receptor antagonists of the centrally-evoked increases of cardiac inotropic responses in rabbits, 111, 1347

Pargyline, Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

Parietal cells, Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87

Passive cutaneous anaphylactic (PCA) reaction, Differential effects

of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

Patch clamp, Block of ATP-sensitive K^+ channels in isolated mouse pancreatic β -cells by 2,3-butanedione monoxime, 112, 143

- Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741
- Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
- Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663
- Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745
- Permeation properties and modulation of volume-activated Cl⁻ currents in human endothelial cells, 112, 1049

Paw test, Peripheral and central adrenoceptor modulation of the behavioural effects of clozapine in the paw test, 112, 769

PC12 cells, Fluspirilene block of *N*-type calcium current in NGF-differentiated PC12 cells, 111, 483

- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992

PD 145065, Characterization of ET_B receptors mediating contractions induced by endothelin-1 or IRL 1620 in guinea-pig isolated airways: effects of BQ-123, FR139317 or PD 145065, 111, 1009

- Reversal of established responses to endothelin-1 *in vivo* and *in vitro* by the endothelin receptor antagonists, BQ-123 and PD 145065, 112, 207

PD 151242, [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303

- [¹²⁵I]-PD151242: a selective radioligand for human ET_A receptors, 111, 4

Penile erection, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81

Penitrem A, Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538

(+)-Pentazocine, The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709

Pentobarbitone, A study of the mechanism of MDMA ('Ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds, 111, 131

Peptides, Role of endopeptidase 3.4.24.16 in the catabolism of neuropeptidin, *in vivo* in the vascularly perfused dog ileum, 112, 127

Peptidases, Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310

- Role of endopeptidase 3.4.24.16 in the catabolism of neuropeptidin, *in vivo* in the vascularly perfused dog ileum, 112, 127

Peptide histidine glycine (PHI-Gly), Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541

Peptide histidine isoleucine (PHI), Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541

Peptide histidine valine [PHV(1-42)], Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541

Periaqueductal gray, Hyperpolarization by opioids acting on μ -receptors of a sub-population of rat periaqueductal gray neurones *in vitro*, 113, 121

- Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Perindopril, Effects of bradykinin B₂ receptor antagonism on the hypotensive effects of ACE inhibition, 113, 717

- Lack of correlation of hypotensive effects with prevention of cardiac hypertrophy by perindopril after ligation of rat coronary artery, 112, 837

Peripheral nerve regeneration, Beneficial effect of the Ca²⁺ antagonist, nimodipine, on existing diabetic neuropathy in the BB/Wor rat, 111, 887

Peripheral nerve stimulation, Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951

Peristalsis, Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095

Permeability, vascular, Characterization of receptors mediating vascular responses to endothelin-1 in the conscious rat, 113, 845

- The involvement of endothelial dysfunction, nitric oxide and prostanooids in the rat gastric microcirculatory responses to endothelin-1, 112, 267

Permeabilization, Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525

Peroxisomal proliferation, Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates, 112, 551

Pertussis toxin, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852

- Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252
- Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299
- The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussis toxin, 112, 709
- Potentiation by adenosine of ATP-evoked dopamine release via a pertussis toxin-sensitive mechanism in rat phaeochromocytoma PC12 cells, 112, 992
- Sensitivity of G-protein involved in endothelin-1-induced Ca²⁺ influx to pertussis toxin in porcine endothelial cells *in situ*, 111, 989

Pervanadate, Regulation of vascular and gastric smooth muscle contractility by pervanadate, 113, 403

Peyer's patches, A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593

PF 5901, Effect of a 5-lipoxygenase inhibitor and leukotriene antagonist (PF 5901) on antigen-induced airway responses in neonatally immunized rabbits, 112, 292

PF 10040, Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7

pH, Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926

- Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

pH, intracellular, Triflocin, a novel inhibitor for the Na-HCO₃ symport in the proximal tubule, 112, 465

Phaeochromocytoma cells, Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296

Pharmacokinetics, Measurement and pharmacokinetic analysis of imipramine and its metabolite by brain microdialysis, 112, 625

- Pharmacokinetics, plasma protein binding and urinary excretion of *N*^o-nitro-L-arginine in rats, 111, 394

Pharmacomechanical coupling, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607

Phenelzine, Effects of antidepressants on the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 741

Phenoxybenzamine, Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Phentolamine, Comparison of antinociception induced by supraspinally administered L-arginine and kytorphin, 112, 817

1-Phenylbiguanide, Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

Phenylephrine, Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233

2-Phenylethylamine, The effects of administration of monoamine oxidase-B inhibitors on rat striatal neurone responses to dopamine, 113, 1159

Phoneutria nigriventer venom, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81

Phorbol esters, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

- Bradykinin-stimulated phosphoinositide metabolism in cultured canine

- tracheal smooth muscle cells, 111, 21
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
- Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
- 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55
- Phosphatidylinositol hydrolysis**, Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163
- Phosphodietyl 03**, Role of endopeptidase 3.4.24.16 in the catabolism of neurotensin, *in vivo* in the vascularly perfused dog ileum, 112, 127
- Phosphodiesterase**, The identification of apparently novel cyclic AMP and cyclic GMP phosphodiesterase activities in guinea-pig tracheal smooth muscle, 113, 3
- Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
- Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198
- Phosphodiesterase, cyclic AMP-specific**, Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
- Phosphodiesterase inhibitors**, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423
- Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259
- Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198
- A quantitative comparison of functional and anti-ischaemic effects of the phosphodiesterase-inhibitors, amrinone, milrinone and levosimendan in rabbit isolated hearts, 112, 757
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Separation and characterization of a novel isoenzyme of cyclic nucleotide phosphodiesterase from rat cerebrum, 111, 389
- Phosphodiesterase isoenzymes**, The identification of apparently novel cyclic AMP and cyclic GMP phosphodiesterase activities in guinea-pig tracheal smooth muscle, 113, 3
- Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Separation and characterization of a novel isoenzyme of cyclic nucleotide phosphodiesterase from rat cerebrum, 111, 389
- Phosphoinositide turnover**, Modulatory effects of NMDA on phosphoinositide responses evoked by the metabotropic glutamate receptor agonist 1S,3R-ACPD in neonatal rat cerebral cortex, 112, 231
- Phospholipase A₂**, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852
- Phospholipase C**, Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614
- Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y}-purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
- Dual effects of mastoparan on intracellular free Ca²⁺ concentrations in human astrocytoma cells, 112, 299
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041
- Phospholipase C inhibitors**, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
- Phospholipase D**, Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311
- Phospholipase D inhibitors**, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435
- Phosphoramidon**, Different endothelin receptors involved in endothelin-1 and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137
- Hydrolysis of iodine labelled urodilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204
- Photoreleased nucleotides**, Modulation of cardiac L-type Ca²⁺ channels by GTP_S in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250
- Phrenic nerve**, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642
- Physostigmine**, Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695
- Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19
- Picotamide**, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81
- Picumast**, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262
- Pilocarpine**, The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- Pinacidil**, Modulation of vasodilatation to levocromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Pineal gland**, Presence of P₂-purinoceptors in the rat pineal gland, 112, 107
- Pirenzepine**, Muscarinic (M₁) receptor-mediated inhibition of K⁺-evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of L- and N-type Ca²⁺ channels, 113, 621
- Two populations of muscarinic binding sites in the chick heart distinguished by affinities for ligands and selective inactivation, 113, 1529
- Pituitary**, Block by capsaicin of voltage-gated K⁺ currents in melanotrophs of the rat pituitary, 112, 616
- Pharmacology of high-threshold calcium currents in GH₃C₁ pituitary cells and their regulation by activation of human D₂ and D₄ dopamine receptors, 112, 728
- Pituitary, rat**, Lithium-induced decrease in spontaneous Ca²⁺ oscillations in single GH₃ rat pituitary cells, 112, 390
- Pituitary adenylate cyclase-activating polypeptide (1-27) (PACAP27)**, Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589
- Pituitary cell lines**, Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
- pK₁ values**, A new method for estimating dissociation constants of competitive and non-competitive antagonists with no prior knowledge of agonist concentrations, 111, 219
- pK₁ values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227
- PK11195**, Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257
- Placenta, human**, Inhibition by fatty acids of cyclic AMP-dependent protein kinase activity in brush border membranes isolated from human placental vesicles, 111, 509
- Plasma extravasation**, The role of bradykinin B₁ receptors in the maintenance of intra-articular plasma extravasation in chronic antigen-induced arthritis, 113, 940
- Plasma protein extravasation**, Blockade by oral or parenteral RPR

100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

- Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017
- Mediation by bradykinin of rat paw oedema induced by collagenase from *Clostridium histolyticum*, 112, 453

Plasminogen activator, Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Plasminogen activator inhibitor (PAI), The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289

Platelets, Antiplatelet activities of FK409, a new spontaneous NO releaser, 113, 385

- Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81
- Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071
- Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926
- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334

Platelet-activating factor, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Effect of PF 10040 on PAF-induced airway responses in neonatally immunized rabbits, 111, 7
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963
- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Investigation of the endogenous chemoattractants involved in ¹¹¹In-eosinophil accumulation in passive cutaneous anaphylactic reactions in the guinea-pig, 113, 35
- Long-lasting inhibitory activity of the heterazepinic BN 50730 on exudation and cellular alterations evoked by PAF and LPS, 113, 994
- Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Role of R-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Strain-dependency of leukotriene C₄ generation from isolated lungs of immunized mice, 112, 1230
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

Platelet-activating factor receptors, Autocrine enhancement of leukotriene synthesis by endogenous leukotriene B₄ and platelet-activating factor in human neutrophils, 111, 852

Platelet-activating factor receptor antagonists, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

Platelet activation, Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894

Platelet aggregation, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

- Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272
- Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071
- Effects of the new A₂ adenosine receptor antagonist 8FB-PTP, an 8 substituted pyrazolo-triazolo-pyrimidine, on *in vitro* functional models, 112, 659
- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2U}-purinoceptors, 113, 1057

Pleurisy, carrageein-induced, Cyclo-oxygenase and nitric oxide synthase isoforms in rat carrageein-induced pleurisy, 113, 693

Polyamines, Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355

Poly-L-arginine, Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29

- The effect of allosteric antagonists in modulating muscarinic M₂-receptor function in guinea-pig isolated trachea, 112, 901
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895

Polymerase chain reaction, [¹²⁵I]-PD151242: a selective ligand for endothelin ET_A receptors in human kidney which localizes to renal vasculature, 113, 1303

- Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396

Ponalrestat, The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576

- Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42

Porphyromonas gingivalis, Anti-inflammatory and analgesic activity of the bradykinin antagonist, icatibant (Hoe 140), against an extract from *Porphyromonas gingivalis*, 112, 1004

Postpartum haemorrhage, Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

Postsynaptic potentials, Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303

Postsynaptic potentials, spontaneous, Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695

Potassium, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Potassium channels, Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619, 113, 659

- Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
- Augmentation by intracellular ATP of the delayed rectifier current independently of the glibenclamide-sensitive K-current in rabbit arterial myocytes, 111, 972
- Block of the delayed rectifier current (I_K) by the 5-HT₃ antagonists ondansetron and granisetron in feline ventricular myocytes, 113, 527
- Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942
- Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29
- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- Effect of potassium channel blockade and α_2 -adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341
- Effects of disopyramide and flecainide on the kinetics of inward rectifier potassium channels in rabbit heart muscle, 111, 873
- Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649
- Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324
- Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N^G-monomethyl-L-arginine but not glibenclamide, 111, 711
- Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958
- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55
- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889
- Role of K⁺ channels in the vasodilator response to bradykinin in the

- rat heart, 113, 954
- Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903
- Potassium channels, ATP-sensitive**, The effects of a novel vasodilator, LP-805, on cytosolic Ca^{2+} concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Location of the sulphonylurea receptor at the cytoplasmic face of the β -cell membrane, 113, 903
- Potassium channels, calcium sensitive**, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607
- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Role of Ca^{2+} -dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022
- Potassium channel antagonists**, Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
- Potassium channel openers**, Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619, 113, 659
- Differential vasodilator properties of KRN2391, cromakalim, nitroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278
- Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687
- Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca^{2+} mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588
- Modulation of vasodilatation to levromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889
- Potassium chloride**, Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471
- Potassium conductance**, Adenosine-induced hyperpolarization of the membrane voltage in rat mesangial cells in primary culture, 113, 7
- Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703
- Hyperpolarization by opioids acting on μ -receptors of a subpopulation of rat periaqueductal gray neurones *in vitro*, 113, 121
- Potassium currents**, Blockade by dendrotoxin homologues of voltage-dependent K⁺ currents in cultured sensory neurones from neonatal rats, 113, 959
- The electrophysiological effects of antiarrhythmic potential of a secoaporphine, *N*-allylsecoboldine, 113, 221
- Potassium currents, slow outward**, Block by capsaicin of voltage-gated K⁺ currents in melanotrophs of the rat pituitary, 112, 616
- PPADS**, Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151
- Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614
- Selective antagonism by PPADS at P_{2X}-purinoceptors in rabbit isolated blood vessels, 111, 923
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015
- Prazosin**, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105
- Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717
- Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- p K_1 values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- [³H]-Prazosin, Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart, 111, 533
- Pregnancy**, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Prejunctional inhibition**, Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Prejunctional modulation**, Effect of potassium channel blockade and α_2 -adrenoceptor activation on the release of nitric oxide from non-adrenergic non-cholinergic nerves, 112, 341
- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123
- Preload**, Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334
- Presynaptic inhibition**, Inhibition by opioids acting on μ -receptors of GABAergic and glutamatergic postsynaptic potentials in single rat periaqueductal gray neurones *in vitro*, 113, 303
- Neuropeptide Y inhibits potassium-stimulated glutamate release through Y₂ receptors in rat hippocampal slices *in vitro*, 113, 737
- Presynaptic receptors**, Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- GABA and glutamate release affected by GABA_B receptor antagonists with similar potency: no evidence for pharmacologically different presynaptic receptors, 113, 1515
- Modulation of dopamine and noradrenaline release and of intracellular Ca^{2+} concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733
- Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043
- Priapism**, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81
- Primary afferents**, Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- Primary afferent depolarization**, The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497
- Primed burst stimulation (PB)**, Effects of DAU 6215, a novel 5-hydroxytryptamine₃ (5-HT₃) antagonist on electrophysiological properties of the rat hippocampus, 112, 695
- Procainamide**, The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69
- Procaine**, Ca^{2+} release induced by myotoxin *a*, a radio-labellable probe having novel Ca^{2+} release properties in sarcoplasmic reticulum, 113, 233
- Procaterol**, Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781
- Prodrug**, Regional haemodynamic effects of dopamine and its prodrugs L-dopa and gludopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117
- Proinflammatory action**, Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
- Pro-inflammatory activities in elapid snake venoms, 112, 723
- Prokallikreins**, Pharmacological characterization of rabbit corpus cavernosum relaxation mediated by the tissue kallikrein-kinin system, 113, 81
- Propafenone**, Voltage- and time-dependent inhibitory effects on rat aortic and porcine coronary artery contraction induced by propafenone and quinidine, 113, 1281
- Propofol**, The effects of propofol and enflurane on single calcium

channel currents of guinea-pig isolated ventricular myocytes, 111, 1147

Propranolol, Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

- Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911
- Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

Prostacyclin, Characterization of prostanoid receptors on rat neutrophils, 113, 581

- Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635
- Effects of selective ET_B -receptor stimulation on arterial, 112, 888
- Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
- Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035

Prostaglandins, β -Adrenoceptors mediate inhibition of [3 H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221

- Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E_2 to mice, 112, 735
- Characterization of prostanoid receptors on rat neutrophils, 113, 581
- The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103
- The effect of non-steroidal anti-inflammatory drugs on the accumulation and release of interleukin-1-like activity by peritoneal macrophages from the mouse, 113, 809
- Effects of intrinsic prostaglandins on the spontaneous contractile and electrical activity of the proximal renal pelvis of the guinea-pig, 113, 431
- Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008
- Interleukin-1 β enhances capsaicin-induced neurogenic vasodilatation in the rat skin, 111, 681
- The involvement of bradykinin B_1 and B_2 receptor mechanisms in cytokine-induced mechanical hyperalgesia in the rat, 113, 63
- The involvement of endothelial dysfunction, nitric oxide and prostanoids in the rat gastric microcirculatory responses to endothelin-1, 112, 267
- Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522
- Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
- Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377
- Ontogenetic increase in PGE₂ and PGF_{2 α} receptor density in brain microvessels of pigs, 112, 59
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Prostaglandin E₂, Increase of extracellular brain calcium involved in interleukin-1 β -induced pyresis in the rabbit: antagonism by dexamethasone, 112, 449

- Interactions between indomethacin, noradrenaline and vasodilators in the fetal rabbit ductus arteriosus, 111, 1245
- Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369
- Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983

Prostaglandin EP-receptors, Characterization of the prostaglandin E₂ sensitive (EP)-receptor in the rat isolated trachea, 112, 133

Prostaglandin I₂, Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77

Prostanoid receptors, Characterization of prostanoid receptors on rat neutrophils, 113, 581

- Ontogenetic increase in PGE₂ and PGF_{2 α} receptor density in brain microvessels of pigs, 112, 59
- Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

Prostanoid DP receptors, Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

Prostanoid EP receptors, Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377

Prostanoid EP receptor agonists, Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735

Prostanoid EP₃ receptors, Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

Prostanoid ET receptors, Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature, 112, 704

Prostanoid IP receptor agonists, Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735

Prostanoid TP receptor antagonists, Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

Protein binding, Pharmacokinetics, plasma protein binding and urinary excretion of N^o-nitro-L-arginine in rats, 111, 394

Protein extravasation, Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963

Protein kinases, Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867

- Phorbol ester-induced M-current suppression in bull-frog sympathetic ganglion cells: insensitivity to kinase inhibitors, 113, 55
- Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455

Protein kinase C, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

- Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975
- Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
- Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237
- Effect of a peptide inhibitor of protein kinase C on G-protein-mediated increase in myofilament Ca²⁺-sensitivity in rabbit arterial skinned muscle, 111, 311
- Effects of protein kinase C activators upon the late stages of the ACTH secretory pathway of AtT-20 cells, 113, 171
- 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
- Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Mechanical and biochemical responses to endothelin-1 and endothelin-3 in bovine bronchial smooth muscle, 111, 1163
- Possible roles of protein kinases in neutrophil chemoattractant factor production by leucocytes in allergic inflammation in rats, 113, 1480
- Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311
- Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455

Protein kinase C inhibitors, Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975

- Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237
- Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604

Proton-coupled transport, Substrate specificity of the di/tripeptide transporter in human intestinal epithelia (Caco-2): identification of substrates that undergo H⁺-coupled absorption, 113, 1050

Proximal tubules, The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881

- Triflocin, a novel inhibitor for the Na-HCO₃ symport in the proximal

tubule, 112, 465

Pseudomonas aeruginosa, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493

Pulse wave velocity, Increased arterial distensibility induced by the angiotensin-converting enzyme inhibitor, lisinopril, in normotensive rats, 111, 555

Purinergic mechanisms, Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473

- Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109

Purinergic transmission, Blockade by 4,4'-diisothiocyanatostilbene-2,2'-disulphonate (DIDS) of P_{2x} -purinoceptors in rat vas deferens, 112, 690

- Blockade of P_{2x} -purinoceptors by trypan blue in rat vas deferens, 113, 349
- Investigation of the actions of PPADS, a novel P_{2x} -purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913
- Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085

Purines, Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123

Purinoceptors, The binding of 1,3-[³H]-dipropyl-8-cyclopentylxanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249

- Biological actions of purines on rat megakaryocytes: potentiation by adenine of the purinoceptor-operated cytoplasmic Ca^{2+} oscillation, 112, 349
- Characterization of the effects of 2-methylthio-ATP and 2-chloro-ATP on brain capillary endothelial cells: similarities to ADP and differences from ATP, 112, 775
- Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y} -purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723
- Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71
- Electrophysiological subtypes of inhibitory P₁ purinoceptors on myenteric neurones of guinea-pig small bowel, 113, 703
- A nucleotide receptor that mobilizes Ca^{2+} in the mouse submandibular salivary cell line ST₄₈₅, 111, 1135
- Stimulation of phospholipase C in cultured microvascular endothelial cells from human frontal lobe by histamine, endothelin and purinoceptor agonist, 111, 1041
- Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245

Purinoceptor P₁ subtypes, Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29

- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-, 112, 169
- Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117

Purinoceptor P₂ subtypes, Activation of chloride current by P₂-purinoceptors in rat ventricular myocytes, 111, 1355

- Blockade by 4,4'-diisothiocyanatostilbene-2,2'-disulphonate (DIDS) of P_{2x} -purinoceptors in rat vas deferens, 112, 690
- Blockade of P_{2x} -purinoceptors by trypan blue in rat vas deferens, 113, 349
- Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853
- Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942
- Characterization and transduction mechanisms of purinoceptors in activated rat microglia, 113, 29
- Comparative studies on the affinities of ATP derivatives for P_{2x} -purinoceptors in rat urinary bladder, 112, 1151
- Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191
- Contractions mediated by α_1 -adrenoceptors and P₂-purinoceptors in a cat colon circular muscle, 112, 1237

- Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614
- Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219
- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669
- Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂-purinoceptor classification, 113, 463
- Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357
- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2T}-purinoceptors, 113, 1057
- Investigation of the actions of PPADS, a novel P_{2x}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913
- Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones, 113, 1085
- The P_{2Z}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP, 112, 946
- Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2x}-purinoceptor ligands, 113, 1432
- Pharmacological profile of the ATP-mediated increase in L- calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Presence of P₂-purinoceptors in the rat pineal gland, 112, 107
- Selective antagonism by PPADS at P_{2x}-purinoceptors in rabbit isolated blood vessels, 111, 923
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117
- The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α,β -methylene adenosine 5'-triphosphate, 112, 282
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2x}-purinoceptors, 113, 1015

Purinoceptor P₂ subtype antagonists, Blockade by 4,4'-diisothiocyanatostilbene-2,2'-disulphonate (DIDS) of P_{2x}-purinoceptors in rat vas deferens, 112, 690

- Blockade of P_{2x} -purinoceptors by trypan blue in rat vas deferens, 113, 349

Purinoceptor antagonists, Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71

Purkinje fibres, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

- Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Putamen, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

Putrescine, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Pyloric sphincter, Contribution of ATP and nitric oxide to NANC inhibitory transmission in rat pyloric sphincter, 113, 681

Pyocyanin, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493

Pyramidal neurones, Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160

Pyridoxal-phosphate-6-azophenyl-2'-4'-disulphonic acid (PPADS), Investigation of the actions of PPADS, a novel P_{2x}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913

Pyridoxal-phosphate-6-azophenyl-2',5'-disulphonic acid (iso-PPADS), The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α,β -methylene adenosine 5'-triphosphate, 112, 282

Pyrimidinoceptors, Contractile effects of uridine 5'-triphosphate in the

rat duodenum, 113, 1191

– Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357

Pyrogallol, Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431

– Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287

Pyruvate, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289

Q

Quantal release, Effect of the putative cognitive enhancer, linopirdine (DuP 996), on quantal parameters of acetylcholine release at the frog neuromuscular junction, 111, 1103

– Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Quantum content, The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747

Quinacrine, Mepacrine-induced inhibition of the inward current mediated by 5-HT₃ receptors in rat nodose ganglion neurones, 112, 745

Quinidine, The electrophysiological effects of antiarrhythmic potential of a secoaporphine, *N*-allylsecoboldine, 113, 221

– The electrophysiological effects of dicentrine on the conduction system of rabbit heart, 113, 69

– Voltage- and time-dependent inhibitory effects on rat aortic and porcine coronary artery contraction induced by propafenone and quinidine, 113, 1281

Quinpirole, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

– Depression of high-threshold calcium currents by activation of human D₂ (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

– Functional expression of human D₃ dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391

R

Radioligand binding, The binding of 1,3-[³H]-dipropyl-8-cyclopentyl-xanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249

– Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

– Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151

– Competitive and non-competitive effects of 5-hydroxyindole on 5-HT₃ receptors in N1E-115 neuroblastoma cells, 112, 541

– Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263

Ramipril, Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Raphe nucleus, Differential sensitivity of ³H-agonist binding to pre- and postsynaptic 5-HT_{1A} receptors in bovine brain, 113, 1400

Rauwolscine, A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65

RBL-2H3 cells, Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118

Reactive blue 2, Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151

– Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2X}-purinoceptors, 113, 614

– Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815

– Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245

Receptors, Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508

Receptors, recombinant, Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251

– Comparative pharmacology of recombinant rat AT_{1A}, AT_{1B} and human AT₁ receptors expressed by transfected COS-M6 cells, 112, 277

Receptor binding, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

Receptor modulation, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Receptor-operated ion channels, Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321

Receptor recycling, Routes of delivery of muscarinic acetylcholine receptors to the plasma membrane in NG108-15 cells, 111, 1023

Receptor regulation, Regulation of nicotinic receptors in rat brain following quasi-irreversible nicotinic blockade by chlorisondamine and chronic treatment with nicotine, 113, 917

Reflex tachycardia, Comparison of the haemodynamic profiles of elogidipine and nicardipine in the anaesthetized dog, 111, 49

Relaxation, Cyclic GMP-independent relaxation and hyperpolarization with acetylcholine in guinea-pig coronary artery, 111, 1053

– Peptide histidine isoleucine-like immunoreactivity release from the rat gastric fundus, 113, 541

– Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525

– Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Renal autoregulation, Effect of nitrendipine on autoregulation of perfusion in the cortex and papilla of kidneys from Wistar and stroke prone spontaneously hypertensive rats, 111, 111

Renal failure, Regional haemodynamic effects of dopamine and its prodrugs L-dopa and gludopa in the rat and in the glycerol-treated rat as a model for acute renal failure, 111, 1117

Renal function, The renal functional responses to 5-HT_{1A} receptor agonist, flesinoxan, in anaesthetized, normotensive rat, 112, 214

Renal haemodynamics, ET_A receptor-mediated responses to endothelin-1 and big endothelin-1 in the rat kidney, 111, 729

Renal hypertrophy, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

Renal mesangial cells, Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1

– Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455

Renal pelvis, Effects of intrinsic prostaglandins on the spontaneous contractile and electrical activity of the proximal renal pelvis of the guinea-pig, 113, 431

Renal pelvis, guinea-pig, Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687

Renal tubules, α_{1b} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819

Renin-angiotensin systems, Facilitation by procatenol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781

– A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145

Reperfusion, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007

– Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233

Reserpine, Modulation by central postsynaptic α_2 -adrenoceptors of the jaw-opening reflex induced by orofacial stimulation in rats, 111, 1140

Respiratory distress syndrome, Effect of nitric oxide gas on the generation of nitric oxide by isolated blood vessels: implications for inhalation therapy, 113, 496

Respiratory system, Detection of nitric oxide in exhaled air during administration of nitroglycerin *in vivo*, 111, 825

Respiratory tract virus, ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188

Retina, Enhancement of retinal acetylcholine release by DAMGO: possibly a direct opioid receptor-mediated excitatory effect, 113, 789

– Modulation by endogenous ATP of the light-evoked release of ACh

from retinal cholinergic neurones, 113, 1085

Retinoic acid, Inhibition of nitric oxide synthesis in vascular smooth muscles by retinoids, 113, 1448

Retractor penis, bovine, Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of *S*-nitrosothiols, 111, 1287

Reversed passive Arthus reaction, Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

Rheumatoid arthritis, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

Rhythmic contractions, Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233

Ricinoleic acid, Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of *N*^G-nitro-L-arginine methyl ester, 113, 1127

Rilmenidine, Attenuated renal response to moxonidine and rilmenidine in one kidney-one clip hypertensive rats, 112, 200

Riluzole, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261

RINm5F cells, The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065

Ro 5-4864, Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Ro 19-6327, Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

Ro 31-8220, Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237

Ro 40-5967, Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395

Ro 41-1049, Effect of type A and B monoamine oxidase selective inhibition by Ro 41-1049 and Ro 19-6327 on dopamine outflow in rat kidney slices, 113, 1269

Ro 46-2005, Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251

Rolipram, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

- Effects of phosphodiesterase isoenzyme inhibitors on cutaneous inflammation in the guinea-pig, 112, 332
- Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259
- Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Roussin's Black Salt, Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

RP 67580, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141
- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

RP 73401, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

RPR 100893, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

RS 23597-190, Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213

RU 24213, The effects of clozapine on behavioural responses to the selective 'D₁-like' dopamine receptor agonist, A 68930, and to the selective 'D₂-like' agonist, RU 24213, 113, 839

Rubidium efflux, Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889

Ruthenium red, The effect of allosteric antagonists in modulating muscarinic M₂-receptor function in guinea-pig isolated trachea, 112, 901

- The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103

RX 821002, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

[³H]-RX 821002, The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I₁- and I₂-sites, 112, 1065

Ryanodine, The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103

- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- Release of Ca²⁺ from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca²⁺ entry, 113, 457

S

S-1452, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

S-16257, Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Salbutamol, Anti-inflammatory and bronchodilator properties of RP 73401, a novel and selective phosphodiesterase type IV inhibitor, 113, 1423

- The dose-related hyper-and-hypokalaemic effects of salbutamol and its arrhythmogenic potential, 111, 73
- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328
- Regional haemodynamic responses to acetylcholine, methoxamine, salbutamol and bradykinin during lipopolysaccharide infusion in conscious rats, 112, 1057
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Salivation, Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411

Salmeterol, Effect of salmeterol on human nasal epithelial cell ciliary beating: inhibition of the ciliotoxin, pyocyanin, 112, 493

- The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687

Salt intake, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

[¹²⁵I]-Sar¹Ile⁸ binding, BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179

Sarafotoxins, Characterization of endothelin (ET) receptors in the isolated gall bladder of the guinea-pig: evidence for an additional ET receptor subtype, 112, 1244

- Different endothelin receptors involved in endothelin-1- and sarafotoxin S6B-induced contractions of the human isolated coronary artery, 113, 1471
- Endothelin receptors mediating functional responses in human small arteries and veins, 111, 609
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat

kidney, 111, 515

- Inhibitory effects of nordihydroguaiaretic acid on ET_A -receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561

Sarcoplasmic reticulum, Activation of the sheep cardiac sarcoplasmic reticulum Ca^{2+} -release channel by analogues of sulmazole, 111, 1212

- Ca^{2+} release induced by myotoxin *a*, a radio-labellable probe having novel Ca^{2+} release properties in sarcoplasmic reticulum, 113, 233
- Effect of cyclopiazonic acid, an inhibitor of sarcoplasmic reticulum Ca^{2+} -ATPase, on the frequency-dependence of the contraction-relaxation cycle of the guinea-pig isolated atrium, 113, 1001
- Effects of bipyridylum compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle, 112, 1216
- Effects of cyclopiazonic acid on phenylephrine-induced contractions in the rabbit ear artery, 111, 233

SB 200646, Further characterization of 5-hydroxytryptamine receptors (putative 5-HT_{2B}) in rat stomach fundus longitudinal muscle, 112, 323

SB 200646A, *In vivo* properties of SB 200646A, a 5-HT_{2C/2B} receptor antagonist, 111, 797

SB 204070, The effects of SB 204070, a highly potent and selective 5-HT₄ receptor antagonist, on guinea-pig distal colon, 112, 789

- Further characterization of 5-hydroxytryptamine receptors (putative 5-HT_{2B}) in rat stomach fundus longitudinal muscle, 112, 323

SB 209670, Characterization of three non-peptide endothelin receptor ligands using human cloned ET_A and ET_B receptors, 112, 1251

Schizophrenia, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

Sciatic nerve, frog, Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342

- Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179

SDZ EAA 494, Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists, 111, 1073

Secretion, Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471

Secretomotor neurones, Characterization of 5-hydroxytryptamine receptors mediating mucosal secretion in guinea-pig ileum, 111, 1240

Sedation, Covariation of α_2 -adrenoceptor density and function following irreversible antagonism with EEDQ, 112, 855

Seglitide, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

Seizures, The differential effects of felodipine and nitrendipine on cerebral dihydropyridine binding *ex vivo* and the ethanol withdrawal syndrome in mice, 112, 1017

- Effects of the non-NMDA antagonists NBQX and the 2,3-benzodiazepine GYKI 52466 on different seizure types in mice: comparison with diazepam and interactions with flumazenil, 113, 1349

P-Selectin, Comparative pharmacology of analogues of *S*-nitroso-*N*-acetyl-DL-penicillamine on human platelets, 112, 1071

Seminal vesicle, Presence of neuropeptide Y in the rat seminal vesicle and its effects on noradrenaline- and nerve-induced contractions, 113, 877

Senktide, On the nature of the 5-HT receptor subtype inhibiting acetylcholine release in the guinea-pig ileum, 113, 77

- Presence of NK₃-sensitive neurones in different proportions in the medial habenula of guinea-pig, rat and gerbil, 112, 717

Sensory nerves, Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588

- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

Septic shock, The effect of nitric oxide synthase inhibition on the plasma fibrinolytic system in septic shock in rats, 112, 289

Septum, lateral, Differential effects of centrally-active antihypertensives on 5-HT_{1A} receptors in rat dorso-lateral septum, rat hippocampus and guinea-pig hippocampus, 111, 318

Sequestration, Routes of delivery of muscarinic acetylcholine receptors to the plasma membrane in NG108-15 cells, 111, 1023

D-Serine, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

- Differential modulation of κ and μ opioid antinociception by the glycine/NMDA receptor agonist D-serine, 112, 1002

Serine protease inhibitors, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na^+ and K^+ transport properties in the rabbit cortical collecting duct, 111, 173

Sesquiterpene lactones, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9

Shock, Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153

- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

SH-SY5Y cells, Block of human voltage-sensitive Na^+ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600

- Muscarinic (M₁) receptor-mediated inhibition of K^+ -evoked [³H]-noradrenaline release from human neuroblastoma (SH-SY5Y) cells via inhibition of *L*- and *N*-type Ca^{2+} channels, 113, 621

Sibutramine, The effects of BTS 54 505, a metabolite of sibutramine, on monoamine and excitatory amino acid-evoked responses in the rat dorsolateral geniculate nucleus *in vivo*, 111, 97

Sigma ligands, Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930

Sigma receptors, The effects of sigma ligands and of neuropeptide Y on *N*-methyl-D-aspartate-induced neuronal activation of CA₃ dorsal hippocampus neurones are differentially affected by pertussin toxin, 112, 709

Signal transduction, Effect of cytoplasmic pH on Ca^{2+} -stimulated eicosanoid biosynthesis in human platelets, 113, 926

- Induction of nitric oxide synthase activity in phagocytic cells inhibited by tricyclodecan-9-yl-xanthogenate (D609), 113, 664
- Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389

Siguazodan, Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Silicone collar, Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

SIN-1, Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867

Single channel currents, Local anaesthetic blockade of neuronal nicotinic ACh receptor-channels in rat parasympathetic ganglion cells, 111, 663

Sinoatrial nodes, α_1 -Adrenoceptors in the conduction system of rat hearts, 111, 465

- Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37
- Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064

SITS, Effects of Cl channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333

SK&F 94120, Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259

SK&F 95654, Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

SK&F 96231, Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198

SK&F 96365, Inhibition by SK&F 96365 of Ca^{2+} current, IL-2 production and activation in T lymphocytes, 113, 861

- Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263
- Variable potency of nitrergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494

SK&F 103829, Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline

mesenteric circulation, 111, 264

Skeletal muscle, Ca^{2+} release induced by myotoxin α , a radio-labellable probe having novel Ca^{2+} release properties in sarcoplasmic reticulum, 113, 233

- Classical and atypical binding sites for β -adrenoceptor ligands and activation of adenylyl cyclase in bovine skeletal muscle and adipose tissue membranes, 111, 866
- Effects of bipyridylum compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle, 112, 1216
- Effects of selective ET_B -receptor stimulation on arterial, 112, 888
- Pharmacological relevance of peripheral type benzodiazepine receptors on motor nerve and skeletal muscle, 112, 257

Skeletal muscle, mouse, The origin of the effects of an anticholinesterase on the latencies of action potentials in mouse skeletal muscles, 111, 747

Skin atrophy, Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D_3 , 113, 439

Skin, guinea-pig, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

- Effect of *in vivo* desensitization to leukotriene B_4 on eosinophil infiltration in response to C5a in guinea-pig skin, 113, 117
- Role of CD18 in the accumulation of eosinophils and neutrophils and local oedema formation in inflammatory reactions in guinea-pig skin, 111, 811

Skin, murine, Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D_3 , 113, 439

Skinned fibres, Diosmin-induced increase in sensitivity to Ca^{2+} of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Small intestine, Adrenoceptors mediating relaxation to catecholamines in rat isolated jejunum, 112, 576

- Investigation of the role of 5-HT₃ and 5-HT₄ receptors in ascending and descending reflexes to the circular muscle of guinea-pig small intestine, 112, 1095
- Molecular cloning and expression of human EP₃ receptors: evidence of three variants with differing carboxyl termini, 112, 377

Smooth muscle, The binding of 1,3-[³H]-dipropyl-8-cyclopentyl-xanthine to adenosine A₁ receptors in rat smooth muscle preparations, 113, 1249

- Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143
- Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853
- Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
- Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195
- Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87
- Pilocarpine-induced relaxation of rat tail artery by a non-cholinergic mechanism and in the absence of an intact endothelium, 112, 525
- Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154
- Role of Ca^{2+} -dependent K-channels in the membrane potential and contractility of aorta from spontaneously hypertensive rats, 113, 1022
- Role of phosphodiesterases III and IV in the modulation of vascular cyclic AMP content by the NO/cyclic GMP pathway, 113, 445
- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161
- Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9

Smooth muscle, airways, Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29

- Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237
- The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- ET_B but not ET_A receptor-mediated contractions to endothelin-1 attenuated by respiratory tract viral infection in mouse airways, 112, 1188
- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641
- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895
- Production of PGE₂ by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors, 111, 983
- Regional and species differences in glyburide-sensitive K⁺ channels in airway smooth muscles as estimated from actions of KC 128 and levromakalim, 113, 889

Smooth muscle, arterial, Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925

- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Smooth muscle, bronchial, Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103

- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328

Smooth muscle, colon, Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259

Smooth muscle, gastric, Regulation of vascular and gastric smooth muscle contractility by pervanadate, 113, 403

Smooth muscle, ileum, Specific inhibition of Ca-activated K channels in red cells by selected dihydropyridine derivatives, 111, 903

Smooth muscle, intestinal, Mediation of most atypical effects by species homologues of the β_3 -adrenoceptor, 112, 911

Smooth muscle, tracheal, Effects of muscarinic M₂ and M₃ receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea *in vitro*, 112, 179

- The identification of apparently novel cyclic AMP and cyclic GMP phosphodiesterase activities in guinea-pig tracheal smooth muscle, 113, 3
- Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081
- Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749

Smooth muscle, urinary bladder, Subtypes of purinoceptors in rat and dog urinary bladder smooth muscles, 112, 117

Smooth muscle, uterine, Effects of cyclopiazonic acid on rhythmic contractions in uterine smooth muscle bundles of the rat, 112, 1132

Smooth muscle, vascular, Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977

- Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
- Atypical β -adrenoceptors in the rat isolated common carotid artery, 113, 699
- BMS-180560, an insurmountable inhibitor of angiotensin II-stimulated responses: comparison with losartan and EXP3174, 113, 179
- Contribution of both nitric oxide and a change in membrane potential to acetylcholine-induced relaxation in the rat small mesenteric artery, 112, 831
- Differences in sensitivity of rat mesenteric small arteries to agonists when studied as ring preparations or as cannulated preparations, 112, 579
- Diosmin-induced increase in sensitivity to Ca^{2+} of the smooth muscle

- contractile apparatus in the rat isolated femoral vein, 111, 978
- The effect of hyperglycaemia on function of rat isolated mesenteric resistance artery, 113, 801
- Effect of a peptide inhibitor of protein kinase C on G-protein-mediated increase in myofilament Ca^{2+} -sensitivity in rabbit arterial skinned muscle, 111, 311
- Effects of Cl^- channel blockers on Ca^{2+} -activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333
- Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635
- Effects of a newly synthesized K^+ channel opener, Y-26763, on noradrenaline-induced Ca^{2+} mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- The effects of a novel vasodilator, LP-805, on cytosolic Ca^{2+} concentrations and on tension in rabbit isolated femoral arteries, 113, 1173
- Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067
- Endothelial function in the isolated perfused mesentery and aorta of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42
- Evidence for a differential location of vasoconstrictor endothelin receptors in the vasculature, 112, 704
- Importance of inositol (1,4,5)-trisphosphate, intracellular Ca^{2+} release and myofilament Ca^{2+} sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525
- Induction of nitric oxide synthase in cultured vascular smooth muscle cells: the role of cyclic AMP, 112, 396
- Inhibition of nitric oxide synthesis in vascular smooth muscles by retinoids, 113, 1448
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209
- Mechanism of the cardiovascular activity of laudanosine: comparison with papaverine and other benzylisoquinolines, 113, 1377
- Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35
- Regulation of vascular and gastric smooth muscle contractility by pervaanol, 113, 403
- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- Smooth muscle cells**, Antimuscarinic action of liriodenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle, 113, 1464
- Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- Smooth muscle cells, intestinal**, All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972
- Smooth muscle cells, tracheal**, Bradykinin-stimulated phosphoinositide metabolism in cultured canine tracheal smooth muscle cells, 111, 21
- The effect of cyclic AMP elevating agents on bradykinin- and carbachol-induced signal transduction in canine cultured tracheal smooth muscle cells, 112, 781
- 5-Hydroxytryptamine receptor-mediated phosphoinositide hydrolysis in canine cultured tracheal smooth muscle cells, 111, 777
- Up-regulation of [³H]-des-Arg¹⁰-kallidin binding to the bradykinin B₁ receptor by interleukin-1 β in isolated smooth muscle cells: correlation with B₁ agonist-induced PGI₂ production, 113, 389
- Smooth muscle cells, vascular**, Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195
- Regulation by hypoxia of endothelin-1-stimulated phospholipase D activity in sheep pulmonary artery cultured smooth muscle cells, 112, 311
- Smooth muscle contraction**, Differences in sensitivity to the specific protein kinase C inhibitor Ro31-8220 between small and large bronchioles of the rat, 113, 1237
- Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628
- Predominance of endothelin_A (ET_A) receptors in ovine airway smooth muscle and their mediation of ET-1-induced contraction, 112, 749
- Smooth muscle relaxation**, Antagonism by lipophilic quaternary ions of the K^+ channel opener, levromakalim, in vascular smooth muscle, 112, 1223
- β_1 - and β_2 -adrenoceptor antagonist activities of ICI-215001, a putative β_3 -adrenoceptor agonist, 112, 55
- Characterization of the prostaglandin E₂ sensitive (EP)-receptor in the rat isolated trachea, 112, 133
- Smooth muscle relaxing effects of NO, nitrosothiols and a nerve-induced relaxing factor released in guinea-pig colon, 113, 1088
- Snake venoms**, Pro-inflammatory activities in elapid snake venoms, 112, 723
- SNP**, Pulmonary effects of type V cyclic GMP specific phosphodiesterase inhibition in the anaesthetized guinea-pig, 111, 1198
- Social interaction test**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
- Sodium**, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640
- Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189
- Sodium absorption**, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31
- Sodium bicarbonate**, Triflocin, a novel inhibitor for the $\text{Na}-\text{HCO}_3$ symport in the proximal tubule, 112, 465
- Sodium/calcium exchange**, Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111, 1067
- Sodium channels**, Block of human voltage-sensitive Na^+ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600
- Extracellular pancuronium affects sodium current in chick embryo sensory neurones, 111, 283
- Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractures in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195
- Sodium conductance**, Mechanisms of the hyperkalaemia caused by nafamostat mesilate: effects of its two metabolites on Na^+ and K^+ transport properties in the rabbit cortical collecting duct, 111, 173
- Sodium currents**, Block of human voltage-sensitive Na^+ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600
- The electrophysiological effects of antiarrhythmic potential of a secoaporphine, N-allylsecoboldine, 113, 221
- Sodium-hydrogen exchange**, Adenosine-sensitive α_1 -adrenoceptor effects on reperfused ischaemic hearts: comparison with phorbol ester, 112, 1007
- Sodium, intracellular**, Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289
- Sodium ions**, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262
- Sodium metabisulphite**, Effect of sodium metabisulphite on bronchial blood flow in conscious sheep: pharmacological modulation, 111, 918
- Sodium nitroprusside**, Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Effects of ethanol and other aliphatic alcohols on NO-mediated relaxations in rat anococcygeus muscles and gastric fundus strips, 111, 1089
- Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894
- The interaction of α -human atrial natriuretic peptide (ANP) with salbutamol, sodium nitroprusside and isosorbide dinitrate in human bronchial smooth muscle, 113, 1328
- Potentiation by forskolin of both SNP- and ANP-stimulated cyclic GMP accumulation in porcine isolated palmar lateral vein, 112, 1146
- Regulation of baseline vascular resistance in the canine diaphragm by nitric oxide, 112, 65
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047
- Variable potency of nitrergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- Sodium pump**, Effect of subthreshold ouabain on the tone of guinea-pig aortic strips following repeated noradrenaline stimulation, 111,

1067

Sodium reabsorption, α_{1b} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819

Sodium transport, The influence of acetazolamide and amlodipine on the intracellular sodium content of rat proximal tubular cells, 112, 881

Somatostatin receptors, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

Spermidine, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Spermidine dialdehyde, The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Spermine, Inhibition by spermine of the induction of nitric oxide synthase in J774.2 macrophages: requirement of a serum factor, 112, 355

- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757

Spinal cord, Actions of two new antagonists showing selectivity for different sub-types of metabotropic glutamate receptor in the neonatal rat spinal cord, 112, 809

- Characterization of EP-receptor subtypes involved in allodynia and hyperalgesia induced by intrathecal administration of prostaglandin E₂ to mice, 112, 735
- The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497
- Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141
- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
- Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358
- Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310
- Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951
- Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_A receptor system, 113, 1561

Spinal cord, rat, A novel antagonist, phenylbenzene ω -phosphono- α -amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord, 113, 165

Spleen, Modulation by endogenous prostanoids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675

Spontaneously hypertensive rats, Endothelium-derived relaxing factor released by 5-HT: distinct from nitric oxide in basilar arteries of normotensive and hypertensive rats, 113, 324

- Enhancement of arterial relaxation by long-term atenolol treatment in spontaneously hypertensive rats, 112, 925
- Mediation by M₃-muscarinic receptors of both endothelium-dependent contraction and relaxation to acetylcholine in the aorta of the spontaneously hypertensive rat, 112, 519

SQ-28,603, Hydrolysis of iodine labelled urodiatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204

SR 47436, A pharmacodynamic study of SR 47436, a selective AT₁ receptor antagonist, on blood pressure in conscious cynomolgus monkeys, 111, 145

SR 48692, The nonpeptide neurotensin antagonist, SR 48692, used as a tool to reveal putative neurotensin receptor subtypes, 112, 352

SR 48968, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

SR 140333, Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635

- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522

SRIF, Mediation by SRIF₁ receptors of the contractile action of somatostatin in rat isolated distal colon; studies using some novel SRIF analogues, 113, 628

Staphylococcus aureus α -toxins, Importance of inositol (1,4,5)-trisphosphate, intracellular Ca²⁺ release and myofilament Ca²⁺ sensitization in 5-hydroxytryptamine-evoked contraction of rabbit mesenteric artery, 111, 525

Staurosporine, An endothelium-dependent contraction in canine mesenteric artery caused by caffeine, 111, 435

- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561

Stereoselective actions, Stereoselective and non-stereoselective actions of isoflurane on the GABA_A receptor, 112, 906

Stereotypy, Anticonvulsant effects of the glycine/NMDA receptor ligands D-cycloserine and D-serine but not R-(+)-HA-966 in amygdala-kindled rats, 112, 97

Steroids, Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone, 113, 1008

Stomach, guinea-pig, 5-HT₃ receptors do not mediate vagally-induced relaxation or contraction of the isolated stomach of the guinea-pig, 111, 346

- Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197

Stomach, mouse, Differential effects of Na⁺,K⁺-ATPase inhibition by ouabain on acid secretory responses to histamine and bethanechol in the mouse isolated stomach, 112, 87

Stomach, rodent, Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93

Stomach fundus, rat, Further characterization of 5-hydroxytryptamine receptors (putative 5-HT_{2B}) in rat stomach fundus longitudinal muscle, 112, 323

Streptomycin, Decrease in calcium currents induced by aminoglycoside antibiotics in frog motor nerve endings, 113, 375

Streptozotocin, Attenuated 5-hydroxytryptamine receptor-mediated responses in aortae from streptozotocin-induced diabetic rats, 111, 370

- Effect of lithium on plasma glucose, insulin and glucagon in normal and streptozotocin-diabetic rats: role of glucagon in the hyperglycaemic response, 111, 861
- The effects of aldose reductase inhibition with ponalrestat on changes in vascular function in streptozotocin diabetic rats, 113, 576
- Endothelial function in the isolated perfused mesentery and aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat, 111, 42
- Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195
- Prevention by insulin treatment of endothelial dysfunction but not enhanced noradrenaline-induced contractility in mesenteric resistance arteries from streptozotocin-induced diabetic rats, 111, 35

Stress, Reversal of CRF- and dopamine-induced stimulation of colonic motility by CCK and igmesine (JO 1784) in the rat, 111, 930

Striatum, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

- Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695
- Comparison of two new inhibitors of catechol O-methylation on striatal dopamine metabolism: a microdialysis study in rats, 112, 13
- Effect of L-arginine/nitric oxide pathway on MPP⁺-induced cell injury in the striatum of rats, 111, 837
- The effects of administration of monoamine oxidase-B inhibitors on rat striatal neurone responses to dopamine, 113, 1159
- Effects of naftidrofuryl oxalate on microsphere embolism-induced decrease in regional blood flow of rat brain, 112, 226
- Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, ctaminoadipate, 113, 536
- Inhibition by KF17837 of adenosine A_{2A} receptor-mediated modulation of striatal GABA and ACh release, 113, 43
- Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23
- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

Striatum, rat, Antagonism by (R)- and (S)-trihexyphenidyl of

muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775

Stroke-prone rats, Replacement of salt by a novel potassium- and magnesium-enriched salt alternative improves the cardiovascular effects of ramipril, 111, 1189

Stroke-prone spontaneously hypertensive rats, Improvement of cardiovascular effects of metoprolol by replacement of common salt with a potassium- and magnesium-enriched salt alternative, 112, 640

Structure-activity relationships, Activation of the sheep cardiac sarcoplasmic reticulum Ca^{2+} -release channel by analogues of sulmazole, 111, 1212

- Binding and functional properties of hexocyclium and sila-hexocyclium derivatives to muscarinic receptor subtypes, 112, 505

Strychnine, Inhibitory effect of strychnine on acetylcholine receptor activation in bovine adrenal medullary chromaffin cells, 113, 471

- A novel antagonist, phenylbenzene ω -phosphono- α -amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord, 113, 165

Strychnine-insensitive glycine sites, Possible NMDA antagonist properties of drugs that affect high pressure neurological syndrome, 111, 951

Submandibular gland, mouse, A nucleotide receptor that mobilizes Ca^{2+} in the mouse submandibular salivary cell line ST₄₈₅, 111, 1135

Substance P, Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

- Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- Effect of the tachykinin NK₁ receptor antagonists, RP 67580 and SR 140333, on electrically-evoked substance P release from rat spinal cord, 113, 635
- Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103
- Inhibition of excitatory non-adrenergic non-cholinergic bronchoconstriction in guinea-pig airways *in vitro* by activation of an atypical 5-HT receptor, 111, 1095
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Inhibition of substance P-induced microvascular leakage by inhaled methoxamine in rat airways, 113, 649
- Involvement of neurokinins in the non-cholinergic response to activation of 5-HT₃ and 5-HT₄ receptors in guinea-pig ileum, 111, 419
- Mechanism underlying substance P-induced relaxation in dog isolated superficial temporal arteries, 111, 77
- Pharmacological reactivity of human epicardial coronary arteries: characterization of relaxation responses to endothelium-derived relaxing factor, 113, 1099
- Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA_B receptor system, 113, 1561
- Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739
- Typical and atypical NK₁ tachykinin receptor characteristics in the rabbit isolated iris sphincter, 112, 985

Substance P release, Lack of effect of microinjection of noradrenaline or medetomidine on stimulus-evoked release of substance P in the spinal cord of the cat: a study with antibody microprobes, 112, 951

Substantia nigra, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831

Sucrose, Factors underlying the increased sensitivity to field stimulation of urinary bladder strips from streptozotocin-induced diabetic rats, 113, 195

Sulmazole analogues, Activation of the sheep cardiac sarcoplasmic reticulum Ca^{2+} -release channel by analogues of sulmazole, 111, 1212

Sulphated polysaccharides, Infection by HIV-1 blocked by binding of dextrin 2-sulphate to the cell surface of activated human peripheral blood mononuclear cells and cultured T-cells, 113, 151

Sulphonylureas, Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649

- Mg^{2+} -dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632

Sulphonylurea receptors, Location of the sulphonylurea receptor at the cytoplasmic face of the β -cell membrane, 113, 903

- Mg^{2+} -dependent inhibition of K_{ATP} by sulphonylureas in CRI-G1 insulin-secreting cells, 111, 632

8-Sulphophenyltheophylline (8-SPT), Action of adenosine receptor antagonists on the cardiovascular response to defence area stimulation in the rat, 113, 159

- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2U}-purinoceptors, 113, 1057

Sulphydryl groups, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9

Sulpiride, Depression of high-threshold calcium currents by activation of human D₂ (short) dopamine receptors expressed in differentiated NG108-15 cells, 111, 1061

- Functional expression of human D₃ dopamine receptors in differentiated neuroblastoma X glioma NG108-15 cells, 111, 391

Sulprostone, Potent contractile actions of prostanoid EP₃-receptor agonists on human isolated pulmonary artery, 113, 369

Sumatriptan, Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

- The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13
- Mediation by 5-HT_{1D} receptors of 5-hydroxytryptamine-induced contractions of rabbit middle and posterior cerebral arteries, 112, 939

Suncus murinus, Induction of emesis in *Suncus murinus* by pyrogallol, a generator of free radicals, 111, 431

Superoxide anion, Inhibition by gomisin C (a lignan from *Schizandra chinensis*) of the respiratory burst of rat neutrophils, 113, 945

- Non-adrenergic, non-cholinergic relaxation of the bovine retractor penis muscle: role of S-nitrosothiols, 111, 1287

Superoxide dismutase, Reduced endothelium-dependent relaxation at enhanced NO release in hearts of hypercholesterolaemic rabbits, 111, 1035

Suramin, Blockade of P_{2X}-purinoceptors by trypan blue in rat vas deferens, 113, 349

- Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853
- Characterization and possible function of adenosine 5'-triphosphate receptors in activated rat microglia, 111, 942
- Comparative studies on the affinities of ATP derivatives for P_{2X}-purinoceptors in rat urinary bladder, 112, 1151
- Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191
- Differential effects of P₂-purinoceptor antagonists on phospholipase C- and adenylyl cyclase-coupled P_{2Y}-purinoceptors, 113, 614
- Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219
- Electrophysiological evidence for different release mechanism of ATP and NO as inhibitory NANC transmitters in guinea-pig colon, 112, 1077
- Evidence for P₂-purinoceptor-mediated inhibition of noradrenaline release in rat brain cortex, 113, 815
- FPL 66096: a novel, highly potent and selective antagonist at human platelet P_{2U}-purinoceptors, 113, 1057
- Pharmacological analysis of ecto-ATPase inhibition: evidence for combined enzyme inhibition and receptor antagonism in P_{2X}-purinoceptor ligands, 113, 1432
- Pharmacological profile of the ATP-mediated increase in L- calcium current amplitude and activation of a non-specific cationic current in rat ventricular cells, 113, 982
- Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455
- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65
- Suramin and reactive blue 2 are antagonists for a newly identified purinoceptor on rat megakaryocyte, 111, 245
- The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α , β -methylene adenosine 5'-triphosphate, 112, 282

Sympathetic co-transmission, Modulatory effects of endothelin-1 on purinergic and adrenergic components of sympathetically-mediated contractile activity of rabbit saphenous artery, 112, 1109

Sympathetic ganglia, The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898

Sympathetic nerves, Facilitation by procaterol, a β -adrenoceptor agonist, of noradrenaline release in the pithed rat independently of angiotensin II formation, 113, 781

- Mechanisms of sympathetic enhancement and inhibition of parasympathetically induced salivary secretion in anaesthetized dogs, 112, 411
- Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

Sympathetic nerve activity, Comparison of the effects of IVth ventricular administration of some tryptamine analogues with those of 8-OH-DPAT on autonomic outflow in the anaesthetized cat, 111, 616

Sympathetic nerve terminals, Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975

Sympathetic nervous system, Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321

- Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- The role of nitric oxide, adrenergic activation and kinin-degradation in blood pressure homeostasis following an acute kinin-induced hypotension, 113, 1567

Sympathetic neurones, The sensitivity of nicotinic synapses in bullfrog sympathetic ganglia to α -bungarotoxin and neuronal-bungarotoxin, 113, 898

Sympathetic neurotransmission, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151

- Evoked noradrenaline release in the rabbit ear artery: enhancement by purines, attenuation by neuropeptide Y and lack of effect of calcitonin gene-related peptide, 112, 123
- Inhibitory 5-hydroxytryptamine receptors involved in pressor effects obtained by stimulation of sympathetic outflow from spinal cord in pithed rats, 113, 1358
- Investigation of the actions of PPADS, a novel P_{2X}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913
- Modulation by neuropeptide Y of parasympathetic nerve-evoked nasal vasodilatation via Y₂ prejunctional receptor, 113, 479
- Pharmacological characterization and anatomical localization of prejunctional β -adrenoceptors in the rat kidney, 111, 1296
- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363

Sympathomimetic activity, Pharmacological actions of the selective and non-selective β -adrenoceptor antagonists celiprolol, bisoprolol and propranolol on human bronchi, 113, 1043

Synaptic plasticity, Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339

Synaptosomes, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

- Cyclothiazide unmasks AMPA-evoked stimulation of [³H]-L-glutamate release from rat hippocampal synaptosomes, 113, 339
- Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Modulation of dopamine and noradrenaline release and of intracellular Ca²⁺ concentration by presynaptic glutamate receptors in hippocampus, 113, 1439
- Regulation of aromatic L-amino acid decarboxylase in rat striatal synaptosomes: effects of dopamine receptor agonists and antagonists, 112, 23

Synovium, AT₁ receptor characteristics of angiotensin analogue binding in human synovium, 112, 435

SZL-49, α_{1b} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819

T

Tachykinins, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

- Characterization of the tachykinin NK₂ receptor in the human bronchus: influence of amastatin-sensitive metabolic pathways, 111, 570
- Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
- Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141
- Facilitatory effects of selective agonists for tachykinin receptors on cholinergic neurotransmission: evidence for species differences, 111, 103
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310
- Role of intrathecal tachykinins for micturition in unanaesthetized rats with and without bladder outlet obstruction, 113, 111
- Role of kinins in anaphylactic-induced bronchoconstriction mediated by tachykinins in guinea-pigs, 113, 508
- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161
- Tachykinin NK₁ receptor subtypes in the rat urinary bladder, 111, 739

Tachykinin receptors, Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240

- Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759
- Evidence for a capsaicin-sensitive, tachykinin-mediated, component in the NANC contraction of the rat urinary bladder to nerve stimulation, 112, 1123
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183
- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161

Tachykinin receptor antagonists, Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250

- Cardiovascular and behavioural effects of centrally administered tachykinins in the rat: characterization of receptors with selective antagonists, 112, 240
- Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342
- Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

Tachykinin NK₁ receptors, Cardiovascular and behavioural effects of centrally administered neuropeptide K in the rat: receptor characterization, 112, 250

- Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150
- Differences in the distribution and characteristics of tachykinin NK₁ binding sites between human and guinea pig lung, 113, 1407
- Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141

Tachykinin NK₁ receptor antagonists, Effects of RP 67580, a tachykinin NK₁ receptor antagonist, on a primary afferent-evoked response of ventral roots in the neonatal rat spinal cord, 113, 1141

- Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161

Tachykinin NK₁ receptor subtypes, Differences in the effects of NK₁-receptor antagonists, (\pm)-CP 96,345 and CP 99,994, on agonist-induced responses in guinea-pig trachea, 112, 176

Tachykinin NK₂ receptors, Comparison of tachykinin NK₁ and NK₂ receptors in the circular muscle of the guinea-pig ileum and proximal colon, 112, 150

Tachykinin NK₂ receptor antagonists, Tachykinin NK₁ and NK₂ receptor antagonists and atropine-resistant ascending excitatory reflex to the circular muscle of the guinea-pig ileum, 112, 161

Tacrine, Blockade of nicotinic responses by physostigmine, tacrine and other cholinesterase inhibitors in rat striatum, 111, 695

- Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Taenia caeci, Receptor for catecholamines responding to catechol which potentiates voltage-dependent calcium current in single cells from guinea-pig taenia caeci, 111, 1154

Taenia caeci, guinea-pig, Bradykinin B₂ receptors and coupling mechanisms in the smooth muscle of the guinea-pig taenia caeci, 113, 607

- Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Taurodeoxycholate, Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle, 112, 1209

[³H]TBPS binding, Effects of GABA and various allosteric ligands on TBPS binding to cloned rat GABA_A receptor subtypes, 112, 1025

T-cells, human cultured, Infection by HIV-1 blocked by binding of dextrin 2-sulphate to the cell surface of activated human peripheral blood mononuclear cells and cultured T-cells, 113, 151

TCV-116, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662

Temelastine, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262

Temperature, Cooling effects on nitric oxide production by rabbit ear and femoral arteries during cholinergic stimulation, 113, 550

Tenoxicam, Effects of tenoxicam and aspirin on the metabolism of proteoglycans and hyaluronan in normal and osteoarthritic human articular cartilage, 113, 1113

Terfenadine, Modulation of antagonist binding to histamine H₁-receptors by sodium ions and by 2-amino-2-hydroxymethyl-propan-1,3-diol HCl, 111, 1262

Tetanic contracture, Prevention by specific chemical classes of α₁-adrenoceptor antagonists of veratrine-contractures in rat left atria independently of α₁-adrenoceptor blockade, 112, 195

Tetrandrine, Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767

Tetraphenylboron, Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levcromakalim, in vascular smooth muscle, 112, 1223

Tetraphenylphosphonium, Antagonism by lipophilic quaternary ions of the K⁺ channel opener, levcromakalim, in vascular smooth muscle, 112, 1223

Tetrodotoxin, Block of human voltage-sensitive Na⁺ currents in differentiated SH-SY5Y cells by lifarizine, 113, 600

- Differential effects of ω-conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
- Effects of caffeine on intracellular sodium activity in cardiac Purkinje fibres: relation to force, 113, 289
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895

Thapsigargin, Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118

- Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, 113, 767
- The effect of calcium removal on the suppression by adenosine of epileptiform activity in the hippocampus: demonstration of desensitization, 112, 316

Theophylline, Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423

- The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769

Thermodynamics, Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor, 111, 295

Thermogenesis, Potentiation of the anti-obesity effect of the selective β₃-adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231

Thermolysin, Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137

Thiol inhibitors, Photon pharmacology of an iron-sulphur cluster nitrosyl compound acting on smooth muscle, 113, 87

Thioperamide, Novel histamine H₃ receptor antagonists: affinities in an H₃ receptor binding assay and potencies in two functional H₃ receptor models, 112, 1043

- Role of histamine in rodent antinociception, 111, 1269

Thiorphan, Generation by the phosphoramidon-sensitive peptidases, endopeptidase-24.11 and thermolysin, of endothelin-1 and C-terminal fragment from big endothelin-1, 113, 137

- Involvement of enzymatic degradation in the inactivation of tachykinin neurotransmitters in neonatal rat spinal cord, 113, 310

L-threo-DOPS, The noradrenaline precursor L-threo-3,4-dihydroxyphenylserine exhibits antinociceptive activity via central α-adrenoceptors in the mouse, 111, 503

Thrombin, Evidence that nitric oxide is an endogenous antiangiogenic mediator, 111, 894

- Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

Thrombocytopenia, Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111

Thrombolysis, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272

- Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Thrombosis, Antithrombotic actions of argatroban in rat models of venous, 'mixed' and arterial thrombosis, and its effects on the tail transection bleeding time, 113, 1209

- Prevention of thrombosis and rethrombosis and enhancement of the thrombolytic actions of recombinant tissue-type plasminogen activator in the canine heart by DMP728, a glycoprotein IIb/IIIa antagonist, 113, 1333

Thromboxane, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

- Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272
- Effect of cytoplasmic pH on Ca²⁺-stimulated eicosanoid biosynthesis in human platelets, 113, 926
- Endothelin-1-induced myocardial ischaemia and oedema in the rat: involvement of the ET_A receptor, 112, 963
- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Modulation by endogenous prostanooids of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675

Thromboxane A₂, Block of Endothelin-1-induced release of thromboxane A₂ from the guinea-pig lung and nitric oxide from the rabbit kidney by a selective ET_B receptor antagonist, BQ-788, 113, 1257

- Effects of isocarbacyclin, a stable prostacyclin analogue, on monkey isolated cerebral and peripheral arteries, 112, 635
- Pharmacological reactivity of human epicardial coronary arteries: phasic and tonic responses to vasoconstrictor agents differentiated by nifedipine, 113, 1093

Thromboxane A₂ receptors, Contribution of thromboxane A₂ to the antigen-induced immediate asthmatic response mediated by IgG1 antibody by augmentation of bronchial responsiveness in guinea-pigs, 111, 339

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Pharmacological characterization of cinnamophilin, a novel dual inhibitor of thromboxane synthase and thromboxane A₂ receptor, 111, 906

Thromboxane A₂ receptor antagonists, Pharmacological characterization of cinnamophilin, a novel dual inhibitor of thromboxane synthase and thromboxane A₂ receptor, 111, 906

Thromboxane generation, Possible role of cyclic AMP phosphodiesterases in the actions of ibudilast on eosinophil thromboxane generation and airways smooth muscle tone, 111, 1081

Thromboxane receptor antagonists, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

Thromboxane synthase inhibitors, Pharmacological characterization of cinnamophilin, a novel dual inhibitor of thromboxane synthase and thromboxane A₂ receptor, 111, 906

Thromboxane synthesis inhibitors, Binding kinetics and antiplatelet activities of picotamide, a thromboxane A₂ receptor antagonist, 112, 81

[³H]-Thymidine, Characterization of the angiotensin II AT₁ receptor subtype involved in DNA synthesis in cultured vascular smooth muscle cells, 112, 1195

Tissue plasminogen activator, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272

Tizanidine, The effects of central myorelaxants on synaptically-evoked primary afferent depolarization in the immature rat spinal cord *in vitro*, 111, 497

TMB-8, The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103

Tolbutamide, Actions of cromakalim on outward currents of CA1 neurones in hippocampal slices, 113, 411

- Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K⁺ channel in mouse B-cells, 111, 302

Tolerance, Parallel tolerance between platelet cyclic GMP and preload effects of nitroglycerin in anaesthetized mini-pigs, 113, 334

Torpedo, Tacrine-induced increase in the release of spontaneous high quantal content events in *Torpedo* electric organ, 112, 19

Toxicity, Inhibition of the glutamate transporter and glial enzymes in rat striatum by the gliotoxin, oxamino adipate, 113, 536

Trachea, B₂-Adrenoceptors mediate inhibition of [³H]-acetylcholine release from the isolated rat and guinea-pig trachea: role of the airway mucosa and prostaglandins, 113, 1221

- Contractile properties of synthetic cationic polypeptides in guinea-pig isolated trachea, 111, 29
- Des-Arg⁹-bradykinin-induced increases in intracellular calcium ion concentration in single bovine tracheal smooth muscle cells, 112, 934
- Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461
- Involvement of apamin-sensitive K⁺ channels in antigen-induced spasm of guinea-pig isolated trachea, 112, 958
- 'Sensory-efferent' neural control of mucus secretion: characterization using tachykinin receptor antagonists in ferret trachea *in vitro*, 113, 1183

Trachea, guinea-pig, B₁- and B₂-adrenoceptor antagonist activities of ICI-215001, a putative B₃-adrenoceptor agonist, 112, 55

- Characterization of the histamine receptors in the guinea-pig lung: evidence for relaxant histamine H₁ receptors in the trachea, 111, 445
- Differences in the effects of NK₁-receptor antagonists, (±)-CP 96,345 and CP 99,994, on agonist-induced responses in guinea-pig trachea, 112, 176
- The effect of allosteric antagonists in modulating muscarinic M₂-receptor function in guinea-pig isolated trachea, 112, 901
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- The interaction between salmeterol and B₂-adrenoceptor agonists with higher efficacy on guinea-pig trachea and human bronchus *in vitro*, 113, 687
- Nitric oxide pathway-mediated relaxant effect of bradykinin in the guinea-pig isolated trachea, 111, 83
- A pharmacological analysis of receptors mediating the excitatory response to 5-hydroxytryptamine in the guinea-pig isolated trachea, 112, 763
- Pharmacological characteristics of liriodenine, isolated from *Fissistigma glaucescens*, a novel muscarinic receptor antagonist in guinea-pigs, 113, 275
- Poly-L-arginine-mediated release of acetylcholine from parasympathetic nerves in rat and guinea-pig airways, 112, 895
- Relaxation of guinea-pig trachea by cyclic AMP phosphodiesterase inhibitors and their enhancement by sodium nitroprusside, 111, 1047

Trachea, rat, Characterization of the prostaglandin E₂ sensitive (EP)-receptor in the rat isolated trachea, 112, 133

- Inhibitory effects of nordihydroguaiaretic acid on ET_A-receptor-mediated contractions to endothelin-1 in rat trachea, 111, 561

Transduction, Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469

Transfection, Coupling of a transfected human brain A₁ adenosine receptor in CHO-K1 cells to calcium mobilisation via a pertussis toxin-sensitive mechanism, 111, 1252

Transforming growth factor- β 1, AT₁ receptor-mediated stimulation by angiotensin II of rat aortic fibronectin gene expression *in vivo*, 113, 662

Transgenic mice, Chloride secretion in response to guanylin in colonic epithelia from normal and transgenic cystic fibrosis mice, 112, 31

Transmitter release, Blockade of nicotinic receptor-mediated release of dopamine from striatal synaptosomes by chlorisondamine administered *in vivo*, 111, 414

TRH, Lithium-induced decrease in spontaneous Ca²⁺ oscillations in single GH₃ rat pituitary cells, 112, 390

Trichloroethanol, Trichloroethanol potentiation of γ -aminobutyric acid-activated chloride current in mouse hippocampal neurones, 113, 555

Triflocin, Triflocin, a novel inhibitor for the Na-HCO₃ symport in the proximal tubule, 112, 465

Trifluoperazine, The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103

- Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469

Trigeminal ganglion, Blockade by oral or parenteral RPR 100893 (a non-peptide NK₁ receptor antagonist) of neurogenic plasma protein extravasation within guinea-pig dura mater and conjunctiva, 112, 920

- The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

Trihexyphenidyl enantiomers, Antagonism by (R)- and (S)-trihexyphenidyl of muscarinic stimulation of adenylyl cyclase in rat olfactory bulb and inhibition in striatum and heart, 113, 775

α -Trinositol, Effects of chronic infusions of α -trinositol on regional and cardiac haemodynamics in conscious rats, 113, 129

Tropisetron, Characterization of the 5-hydroxytryptamine receptor type involved in inhibition of spontaneous activity of human isolated colonic circular muscle, 113, 143

- Role of 5-HT₃ receptors in basal and K⁺-evoked dopamine release from rat olfactory tubercle and striatal slices, 113, 968

Trypan blue, Blockade of P_{2X}-purinoceptors by trypan blue in rat vas deferens, 113, 349

(+)-**Tubocurarine**, *In vitro* potency and mode of action of ANQ9040: a novel fast acting muscle relaxant, 113, 642

Tumour necrosis factor- α (TNF- α), Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat, 113, 912

- Drug modulation of antigen-induced paw oedema in guinea-pigs: effects of lipopolysaccharide, tumour necrosis factor and leucocyte depletion, 112, 111
- Expression of nitric oxide synthase in rat glomerular mesangial cells mediated by cyclic AMP, 112, 1
- The mechanism of the inhibitory effect of polyamines on the induction of nitric oxide synthase: role of aldehyde metabolites, 113, 757
- Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328
- Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153
- Study of the effector mechanism involved in the production of haemorrhagic necrosis of the small intestine in rat passive anaphylaxis, 112, 1101

Tumours, Bradykinin binding sites in healthy and carcinomatous human lung, 111, 1228

Tyramine, Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443

Tyrosine kinase, Possible roles of protein kinases in neutrophil chemotactic factor production by leucocytes in allergic inflammation in rats, 113, 1480

- Regulation of vascular and gastric smooth muscle contractility by pervanadate, 113, 403

Tyrosine kinase receptors, Involvement of tyrosine kinase in the induction of cyclo-oxygenase and nitric oxide synthase by endotoxin in cultured cells, 113, 1522

Tyrosine phosphatase, Regulation of vascular and gastric smooth muscle contractility by pervanadate, 113, 403

U

U-46619, Inhibition by salbutamol of the proliferation of human airway smooth muscle cells grown in culture, 111, 641

- Variable potency of nitrenergic-nitrovasodilator relaxations of the mouse anococcygeus against different forms of induced tone, 113, 1494
- Vasoconstrictor responses after neo-intima formation and endothelial removal in the rabbit carotid artery, 112, 471

UK-68,798, Antifibrillatory effects of clofilium in the rabbit isolated heart, 113, 209

UK-74,505, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

UL-FS 49, Electrophysiological effects of S 16257, a novel sino-atrial node modulator, on rabbit and guinea-pig cardiac preparations: comparison with UL-FS 49, 112, 37

Ureter, guinea-pig, Effect of cromakalim and glibenclamide on spontaneous and evoked motility of the guinea-pig isolated renal pelvis and ureter, 111, 687

- Inhibitory transmitter action of calcitonin gene-related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K channels, 113, 588

Urethra, human, Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723

Urinary bladder, Involvement of ET_A receptors in the facilitation by endothelin-1 of non-adrenergic non-cholinergic transmission in the rat urinary bladder, 111, 473

Urinary bladder, monkey, Evidence for an inhibitory 5-HT₄ receptor in urinary bladder of *Rhesus* and *Cynomolgus* monkeys, 111, 213

Urinary bladder, rat, Characterization of the capsaicin-sensitive component of cyclophosphamide-induced inflammation in the rat urinary bladder, 111, 1017

Urodilatin, Hydrolysis of iodine labelled urodilatin and ANP by recombinant neutral endopeptidase EC. 3.4.24.11, 113, 204

Ussing chamber, A distinctive electrophysiological signature from the Peyer's patches of rabbit intestine, 113, 593

Uterus, The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103

UTP, Characterization of P₂-purinoceptors in the smooth muscle of the rat tail artery: a comparison between contractile and electrophysiological responses, 113, 853

- Contractile effects of uridine 5'-triphosphate in the rat duodenum, 113, 1191
- Evidence from desensitization studies for distinct receptors for ATP and UTP on the rat superior cervical ganglion, 112, 357
- Stimulation by extracellular ATP and UTP of the mitogen-activated protein kinase cascade and proliferation of rat renal mesangial cells, 113, 1455
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X}-purinoceptors, 113, 1015

UTP, extracellular, A nucleotide receptor that mobilizes Ca²⁺ in the mouse submandibular salivary cell line ST₈₈₅, 111, 1135

UTP receptors, Differential regulation of inositol 1,4,5-trisphosphate by co-existing P_{2Y}-purinoceptors and nucleotide receptors on bovine aortic endothelial cells, 111, 723

V

Vagus nerve, Comparative analysis of the vagal stimulation of gastric acid secretion in rodent isolated stomach preparations, 112, 93

- The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- 5-HT₃ receptors do not mediate vagally-induced relaxation or contraction of the isolated stomach of the guinea-pig, 111, 346

Vagus nerve, rat, Effects of divalent cations on the potency of ATP and related agonists in the rat isolated vagus nerve: implications for P₂ purinoceptor classification, 113, 463

- The use of antagonists to characterize the receptors mediating depolarization of the rat isolated vagus nerve by α,β -methylene adenosine 5'-triphosphate, 112, 282

Vanadate, Regulation of vascular and gastric smooth muscle contractility by pervanadate, 113, 403

Vascular/cardiac selectivity, Comparison of the haemodynamic pro-

- files of elgiodipine and nicardipine in the anaesthetized dog, 111, 49

Vascular contraction, Sensitivity to protein kinase C inhibitors of nicardipine-insensitive component of high K⁺ contracture in rat and guinea-pig aorta, 112, 604

Vascular injury, Induction of kinin B₁ receptor-dependent vasoconstriction following balloon catheter injury to the rabbit carotid artery, 111, 1029

Vascular permeability, Cutaneous permeability responses to bradykinin and histamine in the guinea-pig: possible differences in their mechanism of action, 111, 159

- Effect of PAF on rat lung vascular permeability: role of platelets and polymorphonuclear leucocytes, 111, 1111
- Interactions of constitutive nitric oxide with PAF and thromboxane on rat intestinal vascular integrity in acute endotoxaemia, 113, 1131
- Studies on the mechanisms involved in the inflammatory response in a reversed passive Arthus reaction in guinea-pig skin: contribution of neutrophils and endogenous mediators, 113, 1363

Vascular reactivity, Effects of dietary marine oil supplementation on reactivity of human buttock subcutaneous arteries and forearm veins *in vitro*, 112, 566

- Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock, 113, 1153

Vascular resistance, Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233

Vascular system, rat, Characterization of NK₁ and NK₂ tachykinin receptors in guinea-pig and rat bronchopulmonary and vascular systems, 111, 759

Vascular tone, Effects of glibenclamide on systemic and splanchnic haemodynamics in conscious rats, 112, 649

- Effects of P₁ and P_{2Y} purinoceptor antagonists on endothelium-dependent and -independent relaxations of rat mesenteric artery to GTP and guanosine, 112, 71

Vascuлатure, human, Metabolism of methylarginines by human vascuлатure; implications for the regulation of nitric oxide synthesis, 112, 43

Vascuлатure, intestinal, Time-dependent enhancement or inhibition of endotoxin-induced vascular injury in rat intestine by nitric oxide synthase inhibitors, 111, 1309

Vascuлатure, renal, rat, Effects of chronic intravenous infusions of dopexamine and isoprenaline to rats on D₁-, β_1 - and β_2 -receptor-mediated responses, 112, 595

Vas deferens, Differential effects of suramin on P₂-purinoceptors mediating contraction of the guinea-pig vas deferens and urinary bladder, 112, 219

- Effect of 7-nitro indazole in neurotransmission in the rat vas deferens: mechanisms unrelated to inhibition of nitric oxide synthase, 113, 282
- Effects of cyclopiazonic acid on contractility and ecto-ATPase activity in guinea-pig urinary bladder and vas deferens, 113, 669
- Functional effects and ligand binding of chimeric galanin-neuropeptide Y (NPY) peptides on NPY and galanin receptor types, 111, 1129
- Investigation of the actions of PPADS, a novel P_{2X}-purinoceptor antagonist, in the guinea-pig isolated vas deferens, 111, 913

Vas deferens, guinea-pig, Biochemical machinery involved in the release of ATP from sympathetic nerve terminals, 111, 975

- Contribution of intra- and extracellular Ca²⁺ to noradrenaline exocytosis induced by ouabain and monensin from guinea-pig vas deferens, 113, 795
- Investigation of the specificity of FK 888 as a tachykinin NK₁ receptor antagonist, 111, 1342
- Non-specific actions of the non-peptide tachykinin receptor antagonists, CP-96,345, RP 67580 and SR 48968, on neurotransmission, 111, 179

Vas deferens, rat, α_1 -Adrenoceptors and calcium sources in adrenergic neurogenic contractions of rat vas deferens, 111, 151

- Blockade by 4,4'-diisothiocyanostilbene-2,2'-disulphonate (DIDS) of P_{2X}-purinoceptors in rat vas deferens, 112, 690
- Blockade of P_{2X}-purinoceptors by trypan blue in rat vas deferens, 113, 349
- Demonstration of the neurotransmitter role of calcitonin gene-related peptides (CGRP) by immunoblockade with anti-CGRP monoclonal antibodies, 111, 703
- Pharmacological properties of the cloned α_{1AD} -adrenoceptor subtype are consistent with the α_{1A} -adrenoceptor characterized in rat cerebral

cortex and vas deferens, 111, 1003

- pK_1 values of prazosin and idazoxan for receptors stimulated by neuronally released transmitter in the epididymal portion of rat isolated vas deferens, 111, 227

Vasoactive intestinal peptides, The influence of the trigeminal ganglion on carotid blood flow in anaesthetized guinea-pigs, 112, 262

- Nitric oxide, and not vasoactive intestinal peptide, as the main neurotransmitter of vagally induced relaxation of the guinea-pig stomach, 113, 1197
- Regional haemodynamic responses to pituitary adenylate cyclase-activating polypeptide and vasoactive intestinal polypeptide in conscious rats, 111, 589

Vasoconstriction, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105

- Differential effects of ω -conotoxin GVIA and tetrodotoxin on vasoconstrictions evoked by electrical stimulation and nicotinic receptor stimulation in canine isolated, perfused splenic arteries, 111, 1321
- Enhancement of noradrenergic constriction of large coronary arteries by inhibition of nitric oxide synthesis in anaesthetized dogs, 112, 443
- Modulation by endogenous prostaglandins of the vasoconstrictor activity of endothelin-1 in the canine isolated, perfused spleen, 113, 675
- Modulatory effect of neuropeptide Y on acetylcholine-induced oedema and vasoconstriction in isolated perfused lungs of rabbit, 113, 973
- Potentiation of responses to sympathetic nerve stimulation and vasoconstrictor agents by SK&F 103829 in the feline mesenteric circulation, 111, 264
- Sympathetic neurotransmission in the tail artery of aging rats, 113, 363
- The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X} -purinoreceptors, 113, 1015

Vasoconstriction, neurogenic, Effects of cyclic AMP and analogues on neurogenic transmission in the rat tail artery, 111, 625

- Effects of cyclic GMP and analogues on neurogenic transmission in the rat tail artery, 112, 867

Vasoconstriction, renal, Evidence from receptor antagonists of an important role for ET_B receptor-mediated vasoconstrictor effects of endothelin-1 in the rat kidney, 111, 515

Vasodilatation, Cutaneous vasodilatation induced by nitric oxide-evoked stimulation of afferent nerves in the rat, 112, 1181

- Effects of chronic infusions of α -trinol on regional and cardiac haemodynamics in conscious rats, 113, 129
- Effects of N^G -nitro-L-arginine methyl ester on regional haemodynamic responses to $MgSO_4$ in conscious rats, 111, 325
- Evidence for differential roles of nitric oxide (NO) and hyperpolarization in endothelium-dependent relaxation of pig isolated coronary artery, 112, 557
- The 5-HT₁-like receptor mediating the increase in canine external carotid blood flow: close resemblance to the 5-HT_{1D} subtype, 113, 13
- Inhibition by SR140333 of NK₁ tachykinin receptor-evoked, nitric oxide-dependent vasodilatation in the hamster cheek pouch microvasculature *in vivo*, 113, 522
- Ischaemia/reperfusion selectively attenuates coronary vasodilatation to an adenosine A₂- but not to an A₁-agonist in the dog, 111, 1233
- Modulation of vasodilatation to levromakalim by adenosine analogues in the rabbit ear: an explanation for hypoxic augmentation, 112, 49
- Nitric oxide formation from glyceryl trinitrate by rabbit aortic strip: detection by rabbit taenia coli concurrent with vasorelaxation, 113, 1508
- Recovery after dietary vitamin E supplementation of impaired endothelial function in vitamin E-deficient rats, 112, 515
- Role of K⁺ channels in the vasodilator response to bradykinin in the rat heart, 113, 954
- The transhepatic action of ATP on the hepatic arterial and portal venous vascular beds of the rabbit: the role of nitric oxide, 113, 987
- Vasoconstrictor and vasodilator responses to various agonists in the rat perfused mesenteric arterial bed: selective inhibition by PPADS of contractions mediated via P_{2X} -purinoreceptors, 113, 1015

Vasopressin receptors, Relaxation of human isolated mesenteric arteries by vasopressin and desmopressin, 113, 419

Vasorelaxation, Inhibition of hypoxia-induced relaxation of rabbit isolated coronary arteries by N^G -monomethyl-L-arginine but not glibenclamide, 111, 711

- Relationship between cytosolic calcium concentration and force in the papaverine-induced relaxation of medial strips of pig coronary artery, 111, 489
- Relaxation of rat thoracic aorta induced by the Ca²⁺-ATPase inhibitor, cyclopiazonic acid, possibly through nitric oxide formation, 111, 655
- Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor, 111, 652
- Vasorelaxant effect of mexiletine in mesenteric resistance arteries of rats, 111, 673

Veins, Agonist-induced desensitization of histamine H₁ receptor-mediated inositol phospholipid hydrolysis in human umbilical vein endothelial cells, 113, 823

- Effects of selective ET_B -receptor stimulation on arterial, 112, 888
- Endothelin receptors mediating functional responses in human small arteries and veins, 111, 351
- Nitrovasodilator-induced relaxation and tolerance development in porcine vena cordis magna: dependence on intact endothelium, 112, 533

Veins, femoral, Diosmin-induced increase in sensitivity to Ca²⁺ of the smooth muscle contractile apparatus in the rat isolated femoral vein, 111, 978

Veins, femoral, rabbit, Differential vasodilator properties of KRN2391, cromakalim, nitrroglycerin and nifedipine in rabbit isolated femoral artery and vein, 111, 278

Veins, forearm, Effects of dietary marine oil supplementation on reactivity of human buttock subcutaneous arteries and forearm veins *in vitro*, 112, 566

Veins, jugular, Agonistic and antagonistic properties of the bradykinin B₂ receptor antagonist, Hoe 140, in isolated blood vessels from different species, 112, 683

Veins, mesenteric, Role of R-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202

Veins, portal, Action of niflumic acid on evoked and spontaneous calcium-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 112, 977

- Effects of Cl channel blockers on Ca-activated chloride and potassium currents in smooth muscle cells from rabbit portal vein, 111, 1333
- Release of Ca²⁺ from intracellular store in smooth muscle cells of rat portal vein by ATP-induced Ca²⁺ entry, 113, 457
- Study on the vascular reactivity and α_1 -adrenoceptors of portal hypertensive rats, 111, 439
- Sustained prejunctional facilitation of noradrenergic neurotransmission by adrenaline as a co-transmitter in the portal vein of freely moving rats, 113, 342

Veins, renal, human, Vasoconstrictor endothelin receptors characterized in human renal artery and vein *in vitro*, 113, 49

Veins, saphenous, Interactions of chloroethylclonidine with rauwolscine- and prazosin-sensitive adrenoceptors in dog saphenous vein, 113, 1263

Veins, saphenous, human, Modulation of noradrenaline release from the sympathetic nerves of the human saphenous vein and pulmonary artery by presynaptic EP₃- and DP-receptors, 111, 733

- A study of ATP as a sympathetic cotransmitter in human saphenous vein, 111, 65

Ventral tegmental area, Comparison between the pharmacology of dopamine receptors mediating the inhibition of cell firing in rat brain slices through the substantia nigra pars compacta and ventral tegmental area, 112, 873

- Effects of dihydropyridine calcium antagonists on rat midbrain dopaminergic neurones, 113, 831

Ventricle, rat, Functional studies on α_1 -adrenoceptor subtypes mediating inotropic effects in rat right ventricle, 111, 539

Ventricle, right, Evidence for a cooperation between adenosine A₂ receptors and β_1 -adrenoceptors on cardiac automaticity in the isolated right ventricle of the rat, 111, 1316

Ventricular cells, Modulation of cardiac L- Ca²⁺ channels by GTP γ S in response to isoprenaline, forskolin and photoreleased nucleotides, 111, 250

Ventricular fibrillation, Antiarrhythmic effects of BN-063, a newly synthesized adenosine A₁ agonist, on myocardial ischaemia in rats, 112, 1031

- Antifibrillatory effects of clofilium in the rabbit isolated heart, 113, 209
- Ventricular function**, Effects of the calcium channel blockers, diltiazem and Ro 40-5967, on systemic haemodynamics and plasma noradrenaline levels in conscious dogs with pacing-induced heart failure, 113, 395
- Ventricular muscles**, Class I and III antiarrhythmic actions of prazosin in guinea-pig papillary muscles, 111, 717
- Verapamil**, Differential inhibition of neuronal calcium entry and [³H]-D-aspartate release by the quaternary derivatives of verapamil and emopamil, 113, 379
- Differential potentiation by depolarization of the effects of calcium antagonists on contraction and Ca²⁺ current in guinea-pig heart, 113, 451
- The effects of phorbol 12,13-diacetate on responses of guinea-pig isolated trachea to methylxanthines, isoprenaline and ryanodine, 111, 769
- Veratridine**, Antagonism by riluzole of entry of calcium evoked by NMDA and veratridine in rat cultured granule cells: evidence for a dual mechanism of action, 113, 261
- Veratrine**, Prevention by specific chemical classes of α_1 -adrenoceptor antagonists of veratrine-contractures in rat left atria independently of α_1 -adrenoceptor blockade, 112, 195
- Vestibular compensation**, A dose-response analysis of the beneficial effects of the ACTH-(4-9) analogue, Org 2766, on behavioural recovery following unilateral labyrinthectomy in guinea-pig, 111, 358
- Virus**, The effect of leukocyte depletion on pulmonary M₂ muscarinic receptor function in parainfluenza virus-infected guinea-pigs, 112, 588
- Vitamin D-analogues**, Inhibition of glucocorticoid-induced epidermal and dermal atrophy with KH 1060 – a potent 20-epi analogue of 1,25-dihydroxyvitamin D₃, 113, 439
- Vitamin E**, Recovery after dietary vitamin E supplementation of impaired endothelial function in vitamin E-deficient rats, 112, 515
- Voltage clamp**, Competitive and non-competitive effects of 5-hydroxy-indole on 5-HT₂ receptors in N1E-115 neuroblastoma cells, 112, 541
- Inhibitory effects of capsaicin on acetylcholine-evoked responses in rat phaeochromocytoma cells, 113, 296
- Voltammetry**, Differential action of (+)-amphetamine on electrically evoked dopamine overflow in rat brain slices containing corpus striatum and nucleus accumbens, 111, 829

W

- W-7**, The effect of caffeine on prostaglandin output from the guinea-pig uterus, 113, 103
- Inhibition of olfactory cyclic nucleotide-activated current by calmodulin antagonists, 111, 469
- Water absorption**, Differential effects of luminal L-arginine and N^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Water fluxes**, Differential effects of luminal L-arginine and N^G-nitro-L-arginine on blood flow and water fluxes in rat ileum, 112, 304
- Water intake**, Mediation by nitric oxide formation in the preoptic area of endotoxin and tumour necrosis factor-induced inhibition of water intake in the rat, 111, 1328
- Water-lick test**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
- Water reabsorption**, α_{1b} -Adrenoceptors mediate renal tubular sodium and water reabsorption in the rat, 111, 819
- WB4101**, Pharmacological evidence of distinct α_1 -adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery, 113, 723
- WEB-2170**, Role of R-calcium channels in the response of the perfused arterial and venous mesenteric vasculature of the rat to platelet-activating factor, 112, 1202
- Whole-cell patch clamp**, Inhibitory effects of caffeine on secretagogue-induced catecholamine secretion from adrenal chromaffin cells of the guinea-pig, 111, 935
- Ion channel modulation by NS 1619, the putative BK_{Ca} channel opener, in vascular smooth muscle, 113, 1538
- Muscarinic acetylcholine response in pyramidal neurones of rat cerebral cortex, 112, 1160
- Whole-cell voltage clamp**, All-or-nothing responses to carbachol in single intestinal smooth muscle cells of rat, 112, 972

X

- Blockade of the inward rectifier potassium current by the Ca²⁺-ATPase inhibitor 2',5'-di(tert-butyl)-1,4-benzohydroquinone (BHQ), 112, 1118
- WIN 64338**, Effects of a novel nonpeptide bradykinin B₂ receptor antagonist on intestinal and airway smooth muscle: further evidence for the tracheal B₃ receptor, 112, 461

Y

- Y-23684**, The pharmacological properties of Y-23684, a benzodiazepine receptor partial agonist, 111, 1170
- Y-24180**, Pharmacological analysis of neutrophil chemotactic factor production by leucocytes and roles of PAF in allergic inflammation in rats, 111, 123
- Y-26763**, Effects of a newly synthesized K⁺ channel opener, Y-26763, on noradrenaline-induced Ca²⁺ mobilization in smooth muscle of the rabbit mesenteric artery, 111, 165
- Yellow star thistle**, Toxic inhibition of smooth muscle contractility by plant-derived sesquiterpenes caused by their chemically reactive α -methylenebutyrolactone functions, 112, 9
- Yohimbine**, Blockade of noradrenaline-induced constrictions by yohimbine and prazosin differs between consecutive segments of cutaneous arteries in guinea-pig ears, 113, 1105

Z

- Zaprinast**, Comparison of the ability of nicardipine, theophylline and zaprinast to restore cardiovascular haemodynamics following inhibition of nitric oxide synthesis, 112, 423
- The identification of apparently novel cyclic AMP and cyclic GMP phosphodiesterase activities in guinea-pig tracheal smooth muscle, 113, 3
- Prevention by phosphodiesterase inhibitors of antigen-induced contraction of guinea-pig colonic smooth muscle, 111, 259
- Zatebradine**, Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
- ZD 1542**, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272
- ZD 7288**, Haemodynamic actions of a novel sino-atrial node function modulator, ZENECA ZD7288, in the anaesthetized dog: a comparison with zatebradine, atenolol and nitrendipine, 113, 1064
- ZM 170809**, Comparative effects of anti-platelet agents as adjuncts to tissue plasminogen activator in a dog model of occlusive coronary thrombosis, 112, 272
- ZM 230487**, Effect of a 5-lipoxygenase inhibitor, ZM 230487, on cutaneous allergic inflammation in the guinea-pig, 111, 1205
- Zucker rat**, Potentiation of the anti-obesity effect of the selective β_3 -adrenoceptor agonist BRL 35135 in obese Zucker rats by exercise, 113, 1231
- Zymosan**, Differential effects of the PAF receptor antagonist UK-74,505 on neutrophil and eosinophil accumulation in guinea-pig skin, 113, 513

British Journal of Pharmacology

Instructions to Authors	3
Nomenclature Guidelines for Authors	11
Index to Volumes 111, 112 and 113	
Author Index	15
Subject Index	43

1994