

## **Pollution Studies in the Río Santiago Basin, Tributary of the Río de la Plata Estuary: Preliminary Risk Assessment Evaluation**

Alicia E. Ronco, M. Cecilia Sobrero, and Gustavo Bulus Rossini

Centro de Investigaciones del Medio Ambiente, Facultad de Ciencias Exactas,  
Universidad Nacional de La Plata 47 & 115, 1900-La Plata, Argentina

Recent developments in the field of ecotoxicological effects assessment of pollutants on aquatic systems make use of methods founded on statistical principles and requirements (Kooijman 1987; Van Straalen & Denneman 1989). Van Straalen and Kooijman methods estimate a hazardous concentration for  $p$  percent of species in a community (HC<sub>p</sub>) at a chosen protection level. Evaluations on the application of the methodology to different toxic compounds and test species using both chronic and acute toxicity data have been undertaken (Van Leeuwen 1990).

The application of the Van Straalen method to estimate the percentage of species protected at the concentration levels of several contaminants present in the Río Santiago was performed as an initial stage in the hazard assessment of the region. The area is characterized by contamination from a large petrochemical and oil refinery complex, a steel rolling mill and shipyards (Colombo et al. 1990; Catoggio 1991; Ronco et al. 1992; Porta & Ronco, 1992). The calculation of the effect of four polycyclic aromatic hydrocarbons (PAH) (Pyrene, Fluoranthene, Anthracene, Naphthalene) and phenol found in the water body was done mainly using acute toxicity data available in the literature. The assessment was performed at the condition of maximum stress by using the highest concentration level of each contaminant (Catoggio 1991; Ronco et al. 1992). Standardized acute toxicity tests with two small invertebrates and a higher plant seed were performed in a water sample. This sample was also chemically characterized. Results from the calculated assessment based on existent toxicity data for pure compounds are being compared to those from laboratory bioassays on an environmental sample.

---

Send reprint request to AE Ronco at the above address.




Figure 1. Study area location.

## MATERIALS AND METHODS

Acute toxicity data (effective dose concentrations for a 50% reduction of a given end point) for different species were chosen from available reported data (Table 1). Species were selected taking at least 3 different organization levels of aquatic organisms.

Acute toxicity tests were carried with *Daphnia obtusa* Kurz, 1874, emend Scourfield, 1942; *Artemia salina* (Artoxkit M) and *Lactuca sativa* L. A 24h EC50 (inhibition of the mobility) with *D. obtusa* was performed according to the ISO (1989) protocol. The standardized and inter calibrated test with *A. salina*, Artoxkit M, was performed according to Vanhaecke & Persoone (1984) measuring immobility as endpoint after 48 hours of exposure (LC50). A modified lettuce seed germination test (EPA 1989) was used measuring radicle protrusion. Effect on germination was expressed as 120 h EC50. All the tests were performed with three replications and six sample dilutions. LC50 calculations were done by a log-probit curve regression and correlation computer program.

The determination of the PAH content prior solvent extraction followed by a clean up procedure in a Florisil column (Lee et al. 1982) was done by HPLC using a Beckman-Altex, model 100A, with a 4.6x250 mm

Table 1. Acute toxicity data from literature. (A) Algae, (Am) Amphibian, (An) Annelid, (B) Bacterium, (C) Crustacean, (F) Fish, (I) Insect, (P) Protozoan.

| COMPOUND     | TEST SPECIES                          | EC50<br>mg/l |
|--------------|---------------------------------------|--------------|
| Anthracene   | <i>Paramecium caudatum</i> (P)        | 0.0001 (1)   |
|              | <i>Rana pipiens</i> (Am)              | 0.025 (1)    |
|              | <i>Musca sp.</i> (I)                  | 0.027 (1)    |
|              | <i>Lepomis macrochirus</i> (F)        | 0.012 (1)    |
| Fluoranthene | <i>Skeletonema costatum</i> (A)       | 45 (1)       |
|              | <i>Selenastrum capricornutum</i> (A)  | 54 (1)       |
|              | <i>Mysidopsis bahia</i> (C)           | 0.4 (2)      |
|              | <i>Neanthes arenaceodentata</i> (An)  | 0.5 (1)      |
|              | <i>Lepomis macrochirus</i> (F)        | 0.4 (1)      |
| Naphthalene  | <i>Callinectes sapidus</i> (C)        | 2.3 (3)      |
|              | <i>Panaeus aztecus</i> (C)            | 2.5 (3)      |
|              | <i>Cyprinodon variegatus</i> (F)      | 2.4 (3)      |
|              | <i>Pimephales promelas</i> (F)        | 7.8 (4)      |
|              | <i>Neantes arenaceodentata</i> (An)   | 3.8 (3)      |
| Pyrene       | <i>Daphnia magna</i> (C)              | 0.0057 (1)   |
|              | <i>Gambusia affinis</i> (C)           | 0.0026 (1)   |
|              | <i>Aedes aegypti</i> (I)              | 0.035 (1)    |
|              | <i>Pimephales promelas</i> (F)        | 0.026 (1)    |
| Phenol       | <i>Photobacterium phosphoreum</i> (B) | 23.5 (5)     |
|              | <i>Spirillum volutans</i> (B)         | 300 (6)      |
|              | <i>Daphnia magna</i> (C)              | 5.5 (7)      |
|              | <i>Jordanella floridae</i> (F)        | 5.1 (8)      |
|              | <i>Lepomis macrochirus</i> (F)        | 24.0 (4)     |

(1) Costamagna et al. 1990; (2) Suter & Rosen 1988; (3) IRPTC 1987; (4) Vitozzi & De Angelis 1991; (5) Indorato et al. 1984; (6) Dutka & Kwan 1984; (7) Obtained by the authors; (8) Holdway et al. 1991.

Supelco column containing Ultrasphere (5 $\mu$ m), with a Hitachi Vis-UV spectrophotometer as detector and a Hewlett Packard HP 3396A combined computing integrator and plotter.

A direct photometric method (APHA 1985) was used for determination of steam-distillable phenolic compounds, measuring absorbance at 500 nm in the presence of 4-aminoantipyrine and potassium ferricyanide.

The Van Straalen method (Van Straalen & Denneman 1989) was used to estimate the risk associated with the concentration level of each contaminant in the

Table 2. Percentage of species protected as calculated with the Van Straalen and Denneman extrapolation procedure at maximum concentration of toxicants in (I) EFFLUENT DUMPING SITE in CANAL OESTE (sampling campaign 1987-89); (II)down stream in the CANAL OESTE (sampling campaign 1987-89) and, in the RIO SANTIAGO (sampling campaign 1985-86) using acute toxicity from literature.

| COMPOUND     | CONCENTRATION<br>mg/l |       |       | EC50<br>mg/l | T    | HC5<br>mg/l          | % SPECIES<br>PROTECTED |      |      |
|--------------|-----------------------|-------|-------|--------------|------|----------------------|------------------------|------|------|
|              | I                     | II    | III   |              |      |                      | I                      | II   | III  |
| Anthracene   | 0.08                  | 0.018 | 0.007 | 0.0053       | 2240 | $2.4 \times 10^{-6}$ | 26.2                   | 47.4 | 47.4 |
| Fluoranthene | 1.1                   | 0.024 | 0.13  | 2.9          | 1230 | 0.0023               | 59.8                   | 87.9 | 78.2 |
| Naphtalene   | 0.07                  | 0.22  | 0.01  | 3.3          | 4    | 0.81                 | 100                    | 100  | 100  |
| Pyrene       | 0.3                   | 0.6   | 0.02  | 0.011        | 35   | $3.0 \times 10^{-4}$ | 6.0                    | 32.6 | 37.5 |
| Phenol       | 53                    | 15    | 40    | 21.8         | 91   | 0.24                 | 35.8                   | 55.9 | 40.1 |

Table 3. Percentage of species protected as calculated with the Van Straalen and Denneman extrapolation procedure at the concentration levels of the toxicants in Canal Oeste (II sampling point, April 1992).

| COMPOUND     | CONCENTRATION<br>mg/l | % SPECIES<br>PROTECTED |
|--------------|-----------------------|------------------------|
| Anthracene   | 0.018                 | 38.6                   |
| Fluoranthene | 0.01                  | 91.2                   |
| Naphthalene  | 0.2                   | 99.7                   |
| Pyrene       | 0.66                  | 3.2                    |
| Phenol       | 15                    | 55.9                   |

locations indicated in Figure 1. Calculations included the percentage of species protected at the different contaminant concentrations in the environment. The fraction of water life not protected by the HCp was arbitrarily chosen at 0.05 (HC5) as recommended by Van Leeuwen (1990). Four or five test species per toxicant were used in this calculations. Similar calculations were done using experimental results from acute toxicity determinations to a whole water sample.

## RESULTS AND DISCUSSION.

The estimation of the risk associated with the concentration levels of anthracene, fluoranthene, naphthalene, pyrene and phenol in the industries dumping sites and down stream the Canal Oeste and the run off to the Río Santiago basin shows a high impact on the aquatic life of this water bodies. Although chronic test data is recommended for the Van Straalen and Denneman extrapolation procedure, the scope of this work only limits to the initial stage in the regional hazard assessment. Risks related to industrial spills in the area evaluated with acute toxicity data can be a helpful tool complementary with the water quality parameters characterization.

Table 2 shows the calculated percentage of species protected for the maximum concentration levels of the contaminants in the three different locations seen in figure 1. T is an application factor which is already included in the calculation of the HC5.

Even though the very low solubility of PAHs, their

content in the water compartment in the study area associated with the suspended matter, added to the soluble fraction, can reach levels over this limit. Percentages of species protected could be as low as 6% in the industrial dumping sites, partly recovering down stream to 32% in the Canal and 37 % in Río Santiago. This is only considering the effects of single contaminants.

The results of the acute toxicity to three species in laboratory assays to whole water from the II sampling point show LC/EC50 values (expressed as % v/v) of 12.7 to *D.obtusa*, 40.0 to *A.salina* and 56,0 to *L.sativa*. Using these experimental LC/EC50 values in the Van Straalen and Denneman extrapolation procedure to calculate the hazard concentration, it is found that only dilutions of this water of 2.7% or larger would protect 95% of the species (HC5=2.7%; T=11.3). Making the inverse calculation for zero dilution (100% sample water) 19.1 % of the species will be protected.

The estimated percentages of species protected at the concentration levels of anthracene, fluoranthene, naphthalene, pyrene and phenol in the same water sample (Table 3) reach a minimum of 3.2% for a level of pyrene of 0.66 ppm.

Although the estimated percentage of species protected with experimental results is larger than the minimum value obtained from Table 2, this could be due to interactions between the toxicants and other components present in the water as for example dissolved and particulate matter.

These results support field observations in the area. Biota are almost absent within the course of the Canal Oeste and in the sector of Río Santiago where this canal flows into.

Acknowledgments. A.E.Ronco is a member of the Research Career of the National Research Council (CONICET). M.C.Sobrero and G.Bulus Rossini are fellows of CONICET and of the Province Research Council (CICPBA). The project has been supported by a CONICET grant.

## REFERENCES

- APHA (1985) Standard methods for the examination of water and wastewater, 16th edition, Am Publ Hlth Assoc, Washington DC  
Catoggio JA (1991) Contaminación del agua. Causas de

- la contaminación de aguas superficiales y subterráneas, Fundación JE Roulet - Fundación Nauman, Buenos Aires, Argentina, pp 137-155
- Colombo JC, Khalil MF, Arnac M, Horth AC, Catoggio JA (1990) Distribution of chlorinated pesticides and individual polychlorinated biphenyls in biotic and abiotic compartments of the Río de la Plata, Argentina. *Environ Sci Technol* 24:498-505
- Costamagna FM, Iannarilli R, Marcello I, Scaffa L (1990) Rapporti Istisan. Composti aromatici polinucleari. Proprietà chimico-fisiche, tossicologia, ecotossicologia, epidemiologia, valutazioni. Istituto Superiore di Sanità, Roma, pp 862, report 90/4
- Dutka BJ, Kwan KK (1984) Studies on a synthetic activated sludge toxicity screening procedure with comparison to three microbial toxicity tests. In Liu D, Dutka BJ (ed) Toxicity screening procedures using bacterial systems. Marcel Dekker Inc, New York and Basel, pp 125-138
- EPA/600/3-88/029 (1989) Protocols for short term toxicity screening of hazardous waste sites. US Environmental Protection Agency, Corvallis, pp 84-93
- Holdway DA, Dixon DG, Kaiser KLE (1991) The acute toxicity of pulse-dosed, para-substituted phenols to larval American flagfish (*Jordanella floridae*): a comparison with toxicity to photoluminescent bacteria and predicted toxicity using log  $K_{ow}$ . *Science Tot Environ* 104:229-237
- Indorato AM, Snyder KB, Usinowicz PJ (1984) Toxicity screening using Microtox analyzer. In Liu D, Dutka BJ (ed) Toxicity screening procedures using bacterial systems. Marcel Dekker Inc, New York and Basel, pp 37-53
- IRPTC (1987) International register of potentially toxic chemicals. WHO, Geneva
- ISO International standard (1989) Water quality - determination of the mobility of *Daphnia magna* Straus (Cladocera, Crustacea). Reference number ISO 6341:1989 (E)
- Kooijman SALM (1987) A safety factor for LC<sub>50</sub> values allowing for differences in sensitivity among species. *Water Res* 21:269-276
- Lee H-B, Chau ASY, Kawahara F (1982) Organochlorine pesticides. In Chau ASY, Afghan BK (ed). Analysis of pesticides in water, Vol 2. CRC Press Inc, Boca Raton, Florida, pp 21-46
- Porta AA, Ronco AE (1992) Cu(II) acute toxicity to the rotifer *Brachionus calyciflorus*, as affected by fulvic acids from fresh water origin. *Environ Pollut*, in press
- Ronco A, Porta A, Roca A (1992) Las aguas del Río Santiago. Otro caso de contaminación urbano-

- industrial. *Ciencia Hoy* 4:34-38
- Suter GW, Rosen AE (1988) Comparative toxicology for risk assessment of marine fishes and crustaceans. *Environm Sci Technol* 22:548-556
- Vanhaecke P, Persoone G (1984) The ARC-TEST: A standardized short-term routine toxicity test with *Artemia* nauplii. Methodology and evaluation. In: *Ecotoxicological testing for the marine environment*, eds. Persoone G, Jaspers E, Claus C. State Univ. Ghent and Inst. Mar. Scient. Res., Belgium, Vol 2, pp 143-157
- Van Leeuwen K (1990) Ecotoxicological effects assessment in the Netherlands: Recent developments. *Environ Management* 14:779-792
- Van Straalen NM, Denneman CAJ (1989) Ecotoxicological evaluation of soil quality criteria. *Ecotoxicol Environ Safety* 18:241-251
- Vitozzi L, De Angelis G (1991) A critical review of comparative acute toxicity data on fresh fish. *Aquatic Toxicol* 19:167-204

Received December 7, 1992; accepted April 20, 1993.