

Chemical Composition of Major Ions in Rainwater

P. R. Salve · A. Maurya · S. R. Wate ·
Sukumar Devotta

Received: 5 June 2007 / Accepted: 26 December 2007 / Published online: 15 January 2008
© Springer Science+Business Media, LLC 2008

Abstract The present study investigated the chemical composition of rainwater at Kabir nagar, Nari, Nagpur, Maharashtra, India. The rainwater samples were collected on event basis during June–July–August-2006 and were analyzed for pH, major anions Cl^- , NO_3^- , SO_4^{2-} and cations (Ca^{2+} , Mg^{2+} , Na^+ , K^+ , NH_4^+). The pH value varied from 6.0 to 7.3 (avg. 6.3 ± 0.3) indicating alkaline nature of rainwater. The pH of the rainwater was found well above the reference pH (5.6), showing alkalinity during the monsoon season. The average and standard deviation of ionic composition was found to be $98.1 \pm 10.6 \mu\text{eq l}^{-1}$. The total anions contribute 45.1% and cations 54.9%, respectively to rainwater. Neutralization factors (NF) followed a sequence of $\text{NF}_{\text{Ca}} > \text{NF}_{\text{Mg}} > \text{NF}_{\text{NH}_4}$ with factors of 1.1, 0.38 and 0.15 indicating the crustal components are responsible for neutralization of anions. The average ratio of $(\text{NO}_3^- + \text{Cl}^-)/\text{SO}_4^{2-}$ observed as 1.1 indicates that nitric and hydrochloric acid influences the acidity of rainwater. The ratio of $\text{NH}_4^+/\text{NO}_3^-$ and $\text{NH}_4^+/\text{SO}_4^{2-}$ was observed as 0.68 and 0.34 indicate that the possible compounds which may predominate in the atmosphere are NH_4NO_3 and $(\text{NH}_4)_2\text{SO}_4$. Ionic correlation was established to identify sources of origin. A good correlation was seen between Ca and Mg ($r = 0.95$); suggesting the common occurrence of these ions from crustal origin. Similarly, the acidic ions SO_4^{2-} and NO_3^- correlated well ($r = 0.60$) indicating their origin from similar sources. Other relatively significant correlations were observed between Ca and SO_4^{2-} ($r = 0.92$), Mg and SO_4^{2-} ($r = 0.83$), Ca and NO_3^- ($r = 0.09$), Ca and Cl ($r = 0.34$) and Mg and Cl ($r = 0.31$), and Mg and NO_3^-

($r = 0.71$). The observed rainwater ratio of Cl/Na (1.1) is closer to that of seawater ratio (1.16) indicates fractionation of sea-salt and modifications by non-marine constituents as the site is 834 km away from the sea coast. The nss-Ca contribution was observed as 95.7% suggesting their crustal origin whereas nss-Mg and nss-K shows their contribution as 87.9% and 83.2% indicating influence of soil sources. The nss- SO_4^{2-} contributed as 87.4% shows anthropogenic origin.

Keywords Chemical composition · Neutralization · Nonsea-salt · pH · Rainwater · Sources

Rainwater composition plays an important role in scavenging soluble components from the atmosphere and helps us to understand the relative contribution of different sources of atmospheric pollutants (Malik et al. 2007; Tham and Sakugawa 2007). The chemical composition of rainwater varies from site to site and region to region due to influence of local sources. Rainwater is an efficient pathway for removing the gases and particles from the atmosphere. Therefore, rainwater chemistry measurements play a critical role in defining both the level of acid deposition, the state of important biogeochemical cycles of the earth atmosphere system and to understand the relative importance of the different sources (Moody et al. 1991; Galloway et al. 1984; Salve et al. 2006). In India, the reported pH values of rainwater varies from 6 to 7.5, which is higher as compared to reference pH of rainwater (5.6) (Safai et al. 2004; Ali et al. 2004; Mouli et al. 2005; Kulshrestha et al. 2005). The acid rain events are also reported in Delhi (Balachandran and Khilare 2001), Dhanbad (Singh et al. 2007). Literature suggests that if the source is

P. R. Salve (✉) · A. Maurya · S. R. Wate · S. Devotta
National Environmental Engineering Research Institute,
Nehru Marg, Nagpur 440020, India
e-mail: salvepr1@rediffmail.com

influenced by anthropogenic activities, the nature of rainwater becomes acidic because anthropogenic activities contribute acidic gases like SO_2 and NO_x and a basic gas like NH_3 . When these acidic gases are dissolved in clouds, sulphuric and nitric acids are formed, respectively.

The study area fall under residential area and has narrow roads on which vehicular traffic activity takes place. A national highway lies about 0.7 km from the sampling site. There is thermal power plant and small scales industries which is 10 km away from it. The air pollutants encountered in the region are of both particulate and gaseous in nature, which is coming out as vehicular emission, thermal power plant and small scale industries. The thermal power plant emit particulate matters and various gaseous pollutants in to the atmosphere; causing the deterioration of air quality. An attempt has been made to investigate the chemical composition of major ions, understand rainwater chemistry, and generate data on rainwater composition at an urban environment.

Materials and Methods

The rainwater samples were collected on event basis during monsoon season (June–July–August, 2006), at Kabir nagar, near Nari bearing $21^\circ 11' 32''$ N and $79^\circ 06' 13''$ E fixed by Global Positioning System (GPS) located about 8 km from zero miles of Nagpur district, Maharashtra, India.

Samples were collected on the roof of the building about 5 m from ground level. The sampler was placed 1 m from the floor of the roof. Sampling was done manually on an event basis, using only wet collectors, which were fitted with a 2 l polypropylene bottle and a polypropylene funnel 22 cm in diameter. Sample collection equipment used on an event basis were washed with 10% HCl and then rinsed with double distilled water (DDW). The collectors were deployed as soon as the rain began and were withdrawn immediately after they were filled up or when the rain ceased. A total of 21 samples had sufficient volume for chemical analysis, which represented all of the rain events during the study period. Upon arrival at the laboratory, the pH was measured in an aliquot of the sample. The pH meter was calibrated before and after each measurement. After pH measurements, the remaining sample was filtered through a prewashed Whatman filter No. 41 (Balachandran and Khillare 2001; Singh et al. 2001) into precleaned polypropylene bottles. All the filtered samples were preserved at 4°C in a refrigerator until subjected to major inorganic ion analysis by ion chromatograph.

The major inorganic anions and cations were quantitatively determined by Dionex ion chromatograph, a Dionex IonPac AS11 for anions and CS12A for cations, a self-regenerating suppressed conductivity detector and gradient

pump. The gradient weak base eluent (1.7 mM sodium carbonate or 1.8 mM sodium bicarbonate) was used for anion detection, while weak acid eluent (2 mM MSA) for cation detection at a flow of 1 mL/min and air pressure 10 kg/cc. The cation and anion standards were procured from Dionex for calibration purpose. This is highly sensitive method for determination of micro level ions present in environmental samples. The limits of detection for cations and anions were 1 ppb in Dionex ion chromatograph. The analytical precision was maintained through running of known standards. The analytical precision was between 5% and 10% for different anions and cations measurements. The statistical analysis, Oneway ANOVA was applied to estimate the measurement uncertainty in analysis. The analytical variance for the data should comprise not more than, say, 20% of the total variance. The statistics were performed within and between batches using SPSS software and F test results are presented in Table 1. The F test statistic was applied to test the null hypothesis and is true for cations and anions. The calculated F_{value} were observed in the range of 0.014–0.520 which is less than F_{critical} of 2.0 for a stated level of confidence (typically 95%) means that

Table 1 Oneway ANOVA test for cations and anions

Ions	Variation	Sum of squares	df	Mean square	F
Cl	Between groups	0.617	1	0.69	0.090
	Within groups	273.9	40	6.84	
	Total	274.5	41		
SO_4	Between groups	0.829	1	0.829	0.031
	Within groups	1,086.02	40	27.15	
	Total	1,086.8	41		
NO_3	Between groups	0.200	1	0.200	0.075
	Within groups	106.51	40	2.663	
	Total	106.71	41		
Na	Between groups	0.006	1	0.006	0.014
	Within groups	193.7	40	4.88	
	Total	193.7	41		
K	Between groups	0.229	1	0.229	0.214
	Within groups	42.67	40	1.067	
	Total	42.901	41		
Ca	Between groups	17.1	1	17.1	0.520
	Within groups	1,314.52	40	32.86	
	Total	1,331.6	41		
Mg	Between groups	0.007	1	0.007	0.066
	Within groups	43.7	40	1.169	
	Total	46.82	41		
NH_4	Between groups	0.945	1	0.945	0.209
	Within groups	180.9	40	4.523	
	Total	181.88	41		

Table 2 Major ions composition of rainwater ($\mu\text{eq l}^{-1}$)

Component	Min	Max	Average	SD	%
n = 21 no. of samples					
pH	6.0	7.3	6.3	0.3	–
Cl	8.9	21.2	13.3	2.6	13.6
NO ₃	8.1	14.8	10.3	1.7	10.5
SO ₄	9.8	32.8	20.7	5.4	21.1
Na	8.2	15.6	12.1	2.2	12.4
Ca	22.5	42.5	29.1	5.9	29.7
K	1.1	4.9	2.3	1.0	2.3
Mg	1.8	7.1	3.3	1.2	3.3
NH ₄	3.2	12.4	7.0	2.2	7.2

the difference being tested is statistically significant at 95% confidence level (Ramsey et al. 1992).

Results and Discussion

The major ions in rainwater are presented in Table 2. The pH of rainwater is generally affected by acidic and alkaline ions in the rainwater. The average pH of rainwater is found to be alkaline value 6.3 ± 0.3 . The pH of individual rainwater events ranged from 6.0 to 7.3 the showing higher alkalinity which may be due to suspension of particles in the atmosphere which is rich in carbonates and bicarbonates of calcium buffers the acidity generated by mineral acid. The pH was alkaline in all the samples as compared to the reference level, 5.6 (Charlson and Rodhe 1982). Acidic pH reveals the presence of strong acids in rainwater while neutral or alkaline pH indicates neutralization of acids by wind blown soil dust and ammonia.

Rainwater quality is a function of its contents of both acid and alkaline. The average and standard deviation of ionic composition was found to be $98.1 \pm 10.6 \mu\text{eq l}^{-1}$. The ionic balance of rainwater samples showed a trend as $\text{SO}_4 > \text{Cl} > \text{NO}_3$ for anions and $\text{Ca} > \text{Na} > \text{NH}_4 > \text{Mg} > \text{K}$ for cations. Among the ionic constituents calcium contributed maximum (29.7%) followed by sulfate (21.1%), chloride (13.6%), sodium (12.4%), nitrate (10.5%), ammonium (7.2%), magnesium (3.3%) and potassium (2.3%). The total anions contribute 45.1% and cations 54.9%, respectively to rainwater. The slight cationic excess is probably due to some excluded anions. According to literature, acidity originates primarily from sulfuric acid and nitric acid and neutralization by Ca and Mg, so the ratios of $\text{SO}_4 + \text{NO}_3/\text{Ca} + \text{Mg}$ (anions/cations) can be considered as indicator for acidity, if anions/cation is less than one it indicates alkaline nature of rainwater and the ratios greater than unity indicates the presence of free anions which is responsible for precipitation acidity

Table 3 Ratio of different ionic species

Ratio	Values
NO ₃ /SO ₄	0.56
NH ₄ /NO ₃	0.68
NH ₄ /SO ₄	0.34
(Ca + NH ₄)/(NO ₃ + SO ₄)	1.2
(NO ₃ + Cl)/SO ₄	1.1
(SO ₄ + NO ₃)/(Ca + Mg)	0.95

(Kumar et al. 2002). The ratio of sum of SO₄ and NO₃ and sum of Ca and Mg was observed as 0.95 which lower than one indicating alkaline nature of rainwater.

Neutralization factors (NF) were also used to evaluate the neutralization of ions by crustal components and ammonia. The NF_{NH_4} was less than both NF_{Ca} and NF_{Mg} , the factors being 0.15 for NH₄, 0.38 for Mg and 1.1 for Ca, revealing that the crustal components are responsible for neutralization of anions. Studies reported on rainwater at Ahmedabad, and Pune reported NF of 0.73, 0.66 0.45 for NH₄, 1.77, 1.64, 1.25 for Ca and 0.11, 0.43 and 0.23 for Mg, respectively suggesting main neutralization was occurred due to Ca followed by Mg and then ammonium (Rastogi and Sarin 2005; Kulshrestha et al. 2003, Mouli et al. 2005). The ratio of different species is presented in Table 3.

The average ratio of $(\text{NO}_3 + \text{Cl})/\text{SO}_4$ observed as 1.1 indicates that nitric and hydrochloric acid influences the acidity of rainwater whereas ratio below one indicates the influence of sulphuric acid (Khemani et al. 1994). The ratio of $(\text{Ca} + \text{NH}_4)/(\text{NO}_3 + \text{SO}_4)$ is 1.2 indicates that Ca and NH₄ ions plays an important role neutralization of acidic ions in rainwater. The NO₃/SO₄ was observed as 0.56 indicate that about 46% of sulphuric acid and 54% of nitric acid contribution to the acidity in rainwater. The ratio of $(\text{SO}_4 + \text{NO}_3)/(\text{Ca} + \text{Mg})$ was observed as 0.95 which is below one indicating the alkaline nature of rainwater. The ratio of NH₄/NO₃ and NH₄/SO₄ was observed as 0.68 and 0.34 indicate that the possible compounds which may predominate in the atmosphere are NH₄NO₃ and (NH₄)₂SO₄ (Seinfeld 1986). Moreover ammonium nitrate

Table 4 Ionic correlation of rainwater

	Cl	NO ₃	SO ₄	Na	Ca	K	Mg	NH ₄
Cl	1.00	0.82	0.22	0.00	0.34	0.50	0.31	0.11
NO ₃		1.00	0.60	0.88	0.09	0.78	0.71	0.73
SO ₄			1.00	0.54	0.92	0.67	0.83	0.67
Na				1.00	0.45	0.90	0.13	0.33
Ca					1.00	0.21	0.95	0.56
K						1.00	0.71	0.06
Mg							1.00	0.05
NH ₄								1.00

dominates over ammonium sulphates compound (Parmar et al. 2000) whereas an ammonium sulphate dominates over ammonium nitrate (Khare et al. 2004). A correlation between ions in rainwater is presented in Table 4.

A significant correlation was seen between Ca and Mg ($r = 0.95$); suggesting the occurrence from crustal origin. Similarly, the acidic ions SO_4 and NO_3 correlated well ($r = 0.60$) indicating their origin from similar sources, because of the similarity in their behavior in precipitation and the co-emissions of their precursors SO_2 and NO_x . Other relatively significant correlations were observed between Ca and SO_4 ($r = 0.92$), Mg and SO_4 ($r = 0.83$), Ca and NO_3 ($r = 0.09$), Ca and Cl ($r = 0.34$) and Mg and Cl ($r = 0.31$), and Mg and NO_3 ($r = 0.71$). Most of these well-correlated pairs have common sources or occur in rainwater as a result of atmospheric chemical reactions probably from the reaction of acids H_2SO_4 , HNO_3 and HCl with alkaline compounds rich in Ca and Mg carried into the atmosphere by wind blown dust. This suggests that wind carried dust and soil play an important role in rainwater chemistry (Mouli et al. 2005). Ammonium compounds generally occurs in the atmosphere as $(\text{NH}_4)_2\text{SO}_4$ and NH_4NO_3 (Seinfeld 1986), which is supported by the significant correlation between NH_4 and NO_3 ($r = 0.73$) followed by NH_4 and SO_4 ($r = 0.67$).

An attempt has been made to distribute the ratio values with respect to sea salt are presented in Table 5. The enrichment factor of Mg, Ca, Mg and SO_4 was higher than one whereas Cl has value almost equivalent to one indicating significance influence of local sources. The observed rainwater ratio of Cl/Na (1.1) is closer to that of seawater ratio (1.16) indicates fractionation of sea-salt and modifications by non-marine constituents. The elevated K/Na, Mg/Na, Ca/Na SO_4 /Na and NO_3 /Na ratios indicate the possible contribution of other components, probably the soil. The observation of relative ratios among the ionic species suggests that the possible combinations are NaCl, CaSO_4 , MgSO_4 , MgCl_2 , HNO_3 , NH_4NO_3 , NH_4SO_4 and $(\text{NH}_4)_2\text{SO}_4$. The nonsea-salt (nss) contributions of ionic constituent, nss-Ca contribution was observed as 95.7% suggesting their crustal origin whereas nss-Mg and nss-K

shows their contribution as 87.9% and 83.2% indicating influence of soil sources. The nss- SO_4 contributed as 87.4% shows crustal and anthropogenic origin. Studies carried out by other researchers in India shows similar pattern (Rastogi and Sarin 2005; Das et al. 2005). The nss- NO_3 shows 99.998% indicating influence of non-marine sources in the region. This was supported by other study conducted at northern region of India (Hameed et al. 2006).

Acknowledgments The research study was carried out to assess the chemistry of rainwater during monsoon season. The work was supported by Project No. C-7-1425. The authors thank to D.R. Satapathy for reviewing the manuscript.

References

- Ali K, Momin GA, Tiwari S, Safai PD, Chate DM, Rao PSP (2004) Fog and precipitation chemistry at Delhi, North India. *Atmos Environ* 38:4215–4222
- Balachandran S, Khillare PS (2001) Occurance of acid rain over Delhi. *Environ Monit Assess* 71:165–176
- Charlson RJ, Rodhe H (1982) Factors controlling the acidity of natural rainwater. *Nature* 295:683–685
- Das R, Das SN, Misra VN (2005) Chemical composition of rainwater and dustfall at Bhubaneswar in the east coast of India. *Atmos Environ* 39:5908–5916
- Galloway JN, Likens GE, Hawley ME (1984) Acid deposition: natural versus anthropogenic components. *Science* 226:829–831
- Hameed AM, Obaidy JA, Joshi H (2006) Chemical composition of rainwater in a tropical urban area of northern India. *Atmos Environ* 40:6886–6891
- Khare P, Goel A, Patel D, Behari J (2004) Chemical characterization of rainwater at a developing urban habitat of Northern India. *Atmos Res* 69:135–145
- Khemani LT, Momin GA, Rao PSP, Pillai AG, Safai PD, Mohan K, Rao MG (1994) Atmospheric pollutants and their influence on acidification of rain water at an industrial location on the west coast of India. *Atmos Environ* 28:3145–3154
- Kulshrestha UC, Kulshrestha MJ, Sekhar R, Sastry GSR, Vairamani M (2003) Chemical characteristics of rainwater at an urban site of south-central India. *Atmos Environ* 37:3019–3026
- Kulshrestha UC, Granat L, Engardt M, Rodhe H (2005) Review of precipitation monitoring studies in India-a search for regional patterns. *Atmos Environ* 39:7403–7419
- Kumar R, Rani A, Singh SP, Kumari MK, Srivastava SS (2002) A long term study on chemical composition of rainwater at Dayalbagh, a Suburban site of semiarid region. *J Atmos Chem* 41:265–279
- Malik A, Singh VK, Singh KP (2007) Occurrence and distribution of persistent trace organics in rainwater in an urban region (India). *Bull Environ Contam Toxicol*. doi:10.1007/s00128-007-9290-8
- Moody JL, Pszeny AAP, Gaudry A, Keene WC, Galloway JN, Polian G (1991) Precipitation chemistry and its variability in the southern Indian Ocean Amsterdam Island 1980–1987. *J Geophys Res* 96(DII-20):769–786
- Mouli CP, Venkatamohan S, Reddy JS (2005) Rainwater chemistry at a regional representative urban site: influence of terrestrial sources on ionic composition. *Atmos Environ* 39:999–1008
- Parmar RS, Satsangi GS, Lakhani AL, Srivastava SS, Prakash S (2000) Simultaneous measurements of ammonia and nitric acid in ambient air at Agra (27 10N and 28 05E). *Atmos Environ* 35:5979–5988

Table 5 Ratio values of ions, EF, SS and NSS in rainwater

	Cl	K	Mg	Ca	SO_4	NO_3
Sea water	1.16	0.037	0.038	0.12	0.25	0.00002
Rainwater	1.1	0.2	0.3	2.4	1.7	0.00021
EF	0.95	5.09	7.08	20.0	6.2	10.50
SS (%)	48.6	16.8	12.1	4.3	12.6	0.002
NSS (%)	51.4	83.2	87.9	95.7	87.4	99.998

EF = Enrichment factor; SS = Sea-salt fraction; NSS = Nonsea-salt fraction

- Ramsey MH, Thompson M, Hale M (1992) Objective evaluation of precision requirements for geochemical analysis using robust analysis of variance. *J Geochem Expl* 44:23–36
- Rastogi N, Sarin MM (2005) Chemical characteristics of individual rain events from a semi-arid region in India: three-year study. *Atmos Environ* 39:3313–3323
- Safai PD, Rao PSP, Momin GA, Ali K, Chate DM, Praveen PS (2004) Chemical composition of precipitation during 1984–2002 at Pune, India. *Atmos Environ* 38:1705–1714
- Salve PR, Maurya A, Sinha R, Gawane AG, Wate SR (2006) Characterization and source identification of major inorganic ions in precipitation of Nagpur, India. *Bull Environ Contam Toxicol* 77:305–311
- Seinfeld JH (1986) *Atmospheric chemistry and physics of air pollution*, vol 219. John Wiley and Sons, New York
- Singh AK, Mondal GC, Suresh K, Singh KK, Kamal KP, Sinha A (2007) Precipitation chemistry and occurrence of acid rain over Dhanbad, coal city of India. *Environ Monit Assess* 125:99–110 doi: [10.1007/s10661-006-9243-4](https://doi.org/10.1007/s10661-006-9243-4)
- Tham YWF, Sakugawa H (2007) Preliminary study of the photolysis of fluorene in rainwater. *Bull Environ Contam Toxicol*. doi: [10.1007/s00128-007-9309-1](https://doi.org/10.1007/s00128-007-9309-1)