

the morphogenetic potential of embryonic tissues and their interaction during embryogenesis under normal conditions and in the presence of certain pathological processes, such as transplacental carcinogenesis.

LITERATURE CITED

1. T. S. Kolesnichenko, Vopr. Onkol., No. 12, 39 (1966).
2. T. S. Kolesnichenko and T. V. Nikonova, Byull. Eksp. Biol. Med., No. 8, 99 (1975).
3. A. L. Medvinskii and T. S. Kolesnichenko, Byull. Eksp. Biol. Med., No. 3, 379 (1981).
4. L. M. Shabad, T. S. Kolesnichenko, and Yu. D. Sorokina, Transplacental Carcinogenesis and Organ Cultures [in Russian], Moscow (1975).
5. G. M. Hodges, in: Organ Cultures in Biomedical Research (M. Balls and M. Monnickendam, eds.), Cambridge (1976), pp. 15-61.
6. M. S. Steinberg, Proc. Natl. Acad. Sci. USA, 48, 1769 (1962).

ERRATA

BBM-10, October, 1981, V. V. Zakusov et al. p. 1369. The first sentence of the section Experimental Results should read "EA in the course of 20-40 min caused an increase in LP to the noxious thermal stimulus on average by $99 \pm 4\%$ (from 13.6 ± 0.6 to 27.0 ± 1.1 sec) and inhibition of the second positive component of EP on average by $43 \pm 3\%$ from the initial."