

## Constitution of Mucilage from *Ulva Conglobata*. I. Separation of Rhamnobiase Derivative by Partial Methanolysis

Kiyo MITA

Home Economics, Hiroshima Women's University, Hiroshima

(Received February 19, 1969)

Partial methanolysis of the mucilage from *Ulva conglobata* gave methyl  $\alpha$ -L-rhamnoside (29.07%), methyl  $\alpha$ -D-xyloside (7.04%) and methyl  $\alpha$ -D-glucoside (5.13%). Separation of a new disaccharide (methyl  $\alpha$ -L-rhamnobiase, 3.4%) from methanolysate was achieved chromatographically. The biose was attacked by maltase but not by emulsin. The results and infrared spectrum show that the rhamnobiase derivative consists of rhamnose units linked by  $\alpha$ -link.

It was shown<sup>1)</sup> that an acid hydrolysis of the mucilage from *Ulva conglobata* collected in Saga prefecture gave L-rhamnose, D-xylose and D-glucose as neutral constituents and D-glucuronic acid and sulfuric acid as acidic constituents.

The present investigation was to determine the manner of the linkages of these constituents in the polysaccharide molecule.

Percival *et al.*<sup>2-7)</sup> have studied water-soluble sulfated polysaccharides which contain glucose, galactose, mannose, arabinose, xylose, rhamnose and uronic acid residues, from *Caulerpa filiformis*, *Acrosiphonia centralis*, *Cladophora rupestris*, *Ulva lactuca*, *Enteromorpha compressa*, *Codium fragile*, *Chaetomorpha capillaris*. Furthermore, partial hydrolysis of the above mucilages led to the isolation of 3-O- $\beta$ -D-xylopyranosyl-D-xylose, 4-O- $\alpha$ -D-glucopyranosyl-D-glucose, 4-O-D-xylopyranosyl-L-rhamnose, 4-O-D-glucopyranosyl-L-rhamnose and 4-O-glucuronosyl-L-rhamnose-2-sulfate.

Partial methanolysis was conducted by refluxing the suspension of the mucilage in a methanolic hydrogen chloride (1%) for 2 hr. The methanolysate was treated with a barium hydroxide solution (0.3 N) to effect saponification. The neutral methanolysate passed through a cation- and anion exchange resins was subjected to charcoal chromatography,

which led to the isolation of methyl  $\alpha$ -L-rhamnoside (29.07%), methyl  $\alpha$ -D-xyloside (7.04%), methyl  $\alpha$ -D-glucoside (5.13%), a mixture of methyl glycosides of disaccharides (5.5%) and methyl glycosides of oligosaccharides (32.5%).

The mixture of methyl glycosides of disaccharides was further resolved into its components by chromatography on a powdered filter paper column (Table 2). Fraction III-a in Table 2 gave only L-rhamnose by acidic hydrolysis. The fraction was not attacked at all by emulsin but by maltase. The results are very interesting because methyl  $\alpha$ -L-rhamnobiase has not yet been isolated from natural substance and its synthesis has not been reported.<sup>8)</sup>

Barker *et al.*<sup>9,10)</sup> have studied structures of carbohydrates by means of IR spectra and reported that  $\alpha$ -anomers give absorption at  $844 \pm 8$  cm<sup>-1</sup> and  $\beta$ -anomers at  $891 \pm 7$  cm<sup>-1</sup>. This seems to be one of the C-H deformation modes. The absorption bands are observed in all saccharides. Judging from the absorption band at 850 cm<sup>-1</sup>, the isolated rhamnobiase appears to be  $\alpha$ -anomer.

The IR spectrum of xyloside fraction indicated the presence of hydroxy groups from the absorption bands at 3400 and 1100–1000 cm<sup>-1</sup> and  $\alpha$ -anomer from the absorption band at 740 cm<sup>-1</sup> (type 3 of pyranose ring). The IR spectrum of glucoside fraction also identified as  $\alpha$ -anomer from the absorption bands: 3400, 1100–1000 (OH), 840 (type 2a of pyranose ring) and 745 cm<sup>-1</sup> (type 3 of pyranose ring). Furthermore, these  $\alpha$ -anomers were identified by direct comparison of the melting

1) K. Mita, *Nippon Kagaku Zasshi (J. Chem. Soc. Japan, Pure Chem. Sect.)*, **82**, 1664 (1961).

2) I. M. Mackie and E. Percival, *J. Chem. Soc.*, **1959**, 1151; **1960**, 2381; **1961**, 3010.

3) J. J. O'Donnell and E. Percival, *ibid.*, **1959**, 1739, 2168.

4) J. P. McKinnell and E. Percival, *ibid.*, **1962**, 2082, 3141.

5) J. Love, W. Mackie, J. W. McKinnell and E. Percival, *ibid.*, **1963**, 4177.

6) E. L. Hirst, W. Mackie and E. Percival, *ibid.*, **1965**, 2958.

7) Q. N. Haq and E. Percival, *Some Contemp. Stud. Mar. Sci.*, **1966**, 355.

8) S. Kamiya, S. Esaki and M. Hama, *Agr. Biol. Chem.*, **31**, 261, 397 (1967).

9) S. A. Barker, E. J. Bourne, M. Stacey and H. Whiffen, *J. Chem. Soc.*, **1954**, 171; *Chem. & Ind.*, **1953**, 196.

10) S. A. Barker, E. J. Bourne, R. Stephens and D. H. Whiffen, *J. Chem. Soc.*, **1954**, 3468, 4211.

point, the specific rotation and the IR spectrum with authentic samples.

### Experimental

**Materials.** The mucilage was obtained from *Ulva conglobata* (collected in Saga prefecture) as described in the previous paper.<sup>11</sup> All reagents were of analytical or practical grade. Reagents of practical grade were purified by recrystallization and distillation. All melting points are uncorrected.

**Apparatus.** The IR spectra were obtained with KBr disk on a Hitachi EPI-S2 infrared spectrophotometer.

**Partial Methanolysis of the Mucilage.** Powdered mucilage (20 g, moisture 12.2 %, ash 17.6 %) was swelled in a 1 % methanolic hydrogen chloride solution (300 ml) at room temperature for 12 hr and then refluxed for 2 hr. After filtration, the insoluble substance (7.0 g) was dried *in vacuo* at 40°C and refluxed again with a 1 % methanolic hydrogen chloride solution (200 ml) for 2 hr and then filtered. Finally, the insoluble substance (1.85 g) was washed with methanol and dried *in vacuo*.

All filtrates and washings were combined and neutralized with silver carbonate. After filtration, the filtrate was evaporated to a syrup (22.3 g) *in vacuo* at 40°C. This was saponified at 60°C for 2 hr with a 0.3 N barium hydroxide solution (350 ml), and excess barium hydroxide was neutralized with carbon dioxide. After filtration, the filtrate was evaporated to a syrup (18.05 g).

ration, the filtrate was evaporated to a syrup (18.05 g).

**Separation of the Neutral Methanolysate.** The saponified syrup was dissolved in water (10 %) and the aqueous solution was allowed to pass through columns (3×40 cm) of Amberlite IR 120 (H form; 210 ml) and then Amberlite IR 45 (OH form; 210 ml). Both columns were washed with water (5 l) at a flow rate of 200 ml/hr. The combined eluates and washings were evaporated in order to obtain the neutral syrup (8.45 g);  $[\alpha]_D^{20} -21.4^\circ$  (*c* 1.5, water),  $\text{OCH}_3$  15.31 %. The neutral fraction from the methanolysates gave four distinct colored spots with  $R_f$  values 0.10 (D: orange brown), 0.43 (C: orange brown), 0.55 (B: violet blue) and 0.64 (A: orange, main) on a paper chromatogram with *o*-aminophenol reagent,<sup>12</sup> as shown in Fig. 1.

**Adsorption Chromatography for the Neutral Methanolysate.** The neutral methanolysate (8.0 g) was dissolved in water (72 ml), placed on a charcoal-Celite 545 (1:1 w/w) column (5×34 cm) and eluted successively with water and then aqueous ethanol (2, 5, 7.5, 15 and 30 %) at room temperature according to the method of Whistler.<sup>14</sup>

The eluates were measured for specific rotation and chromatographed on papers. The chromatograms were sprayed with *o*-aminophenol<sup>12</sup> and potassium periodate-potassium permanganate reagents.<sup>15</sup> The results are summarized in Table 1.

**Separation and Identification of Methyl  $\alpha$ -D-Xyloside, Methyl  $\alpha$ -L-Rhamnoside and Methyl  $\alpha$ -D-Glucoside.** Fraction I in Table 1 gave three distinct spots on a paper chromatogram:  $R_f$  0.64 (orange; corresponding to methyl  $\alpha$ -L-rhamnoside), 0.55 (violet-blue; corresponding to methyl  $\alpha$ -D-xyloside) and 0.43 (brown-orange; corresponding to methyl  $\alpha$ -D-glucoside).

Fraction I (2.7 g) was further separated through a charcoal-Celite 545 column (2×35 cm), water being used as a mobile phase. Each eluate was examined by specific rotation and then chromatographed on paper. The eluates gave the same specific rotation and  $R_f$  values were combined and then evaporated to dryness. They were identified as follows.

**Methyl  $\alpha$ -D-Xyloside:** The xyloside fraction (white solid, 1.05 g) was dissolved in methanol-acetone (1:3




Fig. 1. Paper chromatogram of the neutral fraction of methanolysates.

Solvent system is *n*-butanol: ethanol: water: ammonia = 40:10:49:1 (v/v).<sup>13</sup>

- A: methyl  $\alpha$ -L-rhamnoside  
B: methyl  $\alpha$ -D-xyloside  
C: methyl  $\alpha$ -D-glucoside  
D: unknown

TABLE 1. CHROMATOGRAPHIC SEPARATION OF NEUTRAL METHANOLYSATE

Fraction	I	II	III	IV	V	VI
Solvent	W* <sup>1</sup>	2% E* <sup>1</sup>	5% E* <sup>1</sup>	7.5% E* <sup>1</sup>	15% E* <sup>1</sup>	30% E* <sup>1</sup>
Vol. (l)	20	12	20	21	11	17
Yield (g)	2.7	0.55	0.5	0.5	1.1	1.0
$[\alpha]_D^{20}(\text{H}_2\text{O})$	—	-87.31°	-89.77°	-84.17°	-57.25°	-53.17°
$\text{OCH}_3$ (%)	—	17.5	6.43	3.58	3.70	4.22
$R_f$ value	{ 0.64 0.55 0.43	0.64	{ 0.30 0.24	{ 0.14* <sup>2</sup> 0.10	0.10	0.10

\*<sup>1</sup> W: water, E: ethanol, \*<sup>2</sup> trace

- 11) K. Mita, *Japanese Soc. Sci. Fish.*, **27**, 239 (1961).  
12) H. Hirase, C. Araki and S. Nakanishi, *This Bulletin*, **26**, 183 (1953).  
13) K. Arai, *Nippon Kagaku Zasshi (J. Chem. Soc. Japan, Pure Chem. Sect.)*, **82**, 1416 (1961).

- 14) R. L. Whistler and D. F. Durso, *J. Am. Chem. Soc.*, **72**, 677 (1950).  
15) M. L. Wolfrom and J. S. Miller, *Anal. Chem.*, **28**, 1037 (1956).

v/v) on heating and then filtered. The filtrate gave white crystals (1.0 g),  $[\alpha]_D^{25} +143.96^\circ$  (*c* 2.7, water), mp  $103.5^\circ\text{C}$  and identified as methyl  $\alpha$ -D-xyloside by means of mixed melting point and comparison of the IR spectrum (KBr): 3400, 1100–1000 (OH),  $740\text{ cm}^{-1}$  (type 3 of pyranose ring) with that of the authentic sample.

Found:  $\text{OCH}_3$ , 18.55%. Calcd for  $\text{C}_5\text{H}_9\text{O}_4(\text{OCH}_3)$ :  $\text{OCH}_3$ , 18.9%.

**Methyl 2,3,4-Tri-O-acetyl- $\alpha$ -L-rhamnopyranoside:** After dissolving the methyl  $\alpha$ -L-rhamnoside fraction (1.1 g) in pyridine (10 ml), acetic anhydride (20 ml) was added. After 24 hr the solution was dropped into ice water and the white solid (0.88 g) separated was filtered. Recrystallization from aqueous-methanol (1 : 2 v/v) gave white crystals,  $[\alpha]_D^{14} -8.10^\circ$  (*c* 0.29, methanol), mp  $84-85^\circ\text{C}$  (lit  $86-87^\circ\text{C}$ ).<sup>16</sup> identified as methyl 2,3,4-tri-O-acetyl- $\alpha$ -L-rhamnopyranoside by means of mixed melting point and comparison of the IR spectrum (KBr): 1745, 1375 and  $1225\text{ cm}^{-1}$  (OAc),  $850\text{ cm}^{-1}$  (type 2a of pyranose ring) with the authentic sample.

Found:  $\text{OCH}_3$ , 18.00%. Calcd for  $\text{C}_6\text{H}_{11}\text{O}_5(\text{OCH}_3)$ :  $\text{OCH}_3$ , 17.4%.

**Methyl  $\alpha$ -D-Glucoside:** The glucoside fraction (white solid: 0.58 g) was dissolved in methanol-acetone (1 : 2 v/v) on heating and then filtered. The filtrate gave white crystals (0.55 g) mp  $163-164^\circ\text{C}$ ,  $[\alpha]_D^{25} +150.30^\circ$  (*c* 0.24, water) which recrystallized from the same mixed solvent. The crystals were identified as methyl  $\alpha$ -D-glucoside by means of mixed melting point and comparison of the IR spectrum (KBr):  $840\text{ cm}^{-1}$  (type 2a of pyranose ring),  $745\text{ cm}^{-1}$  (type 3 of pyranose ring) with that of the authentic sample.

Found:  $\text{OCH}_3$ , 15.40%. Calcd for  $\text{C}_6\text{H}_{11}\text{O}_5(\text{OCH}_3)$ :  $\text{OCH}_3$ , 15.97%.

**Separation and Identification of Methyl  $\alpha$ -L-Rhamnobioid.** A solution (3 ml) of *n*-butanol-water (6 : 1 v/v) containing fraction III (0.5 g) was chromatographed on a column ( $2 \times 50\text{ cm}$ ) of cellulose powder with *n*-butanol saturated with water according to the method of Hough<sup>17</sup> and the eluates (5 ml each) were collected automatically and analyzed by means of specific rotation and paper chromatography. The

eluates containing the same components were combined, and then *n*-butanol was distilled off *in vacuo*. The residues dried on potassium hydroxide in a dessiccator gave white crystals.

TABLE 2. CHROMATOGRAPHIC SEPARATION OF FRACTION III ON CELLULOSE POWDER

Fraction	Eluate No.	Yield (g)	$R_f$ value
III-a	51—58	0.11	0.31
III-b	59—100	0.33	0.31, 0.24
III-c	101—110	0.04	0.24

Fraction III-a in Table 2 recrystallized from methanol-water (4 : 1 v/v) gave needles, mp  $123-125^\circ\text{C}$ ,  $[\alpha]_D^{25} -16.0^\circ$  (*c* 2.0, water),  $R_f$  0.31 (orange brown by *o*-aminophenol reagent).

Found: C, 48.11; H, 7.39%. Calcd for  $\text{C}_{13}\text{H}_{24}\text{O}_9$ : C, 48.14; H, 7.40%.

Found:  $\text{OCH}_3$ , 8.89%. Calcd for  $\text{C}_{12}\text{H}_{21}\text{O}_8(\text{OCH}_3)$ :  $\text{OCH}_3$ , 9.56%.

Fraction III-a was hydrolyzed with 0.1 N sulfuric acid solution (5 ml) in a boiling water bath for 1 hr, neutralized with barium carbonate and then filtered. The filtrate was vacuum evaporated and chromatographed. It gave only rhamnose spot ( $R_f$  0.37).

**Enzymic Hydrolysis of Fraction III-a.** Fraction III-a (20 mg) was hydrolyzed with  $\beta$ -glucosidase (emulsin, 0.1 g) in an acetic acid-sodium acetate buffer solution (pH 4.1) at  $30^\circ\text{C}$  in a water bath for 1 hr. After filtration, the filtrate was chromatographed on paper. The  $R_f$  value did not change at all before treatment with  $\beta$ -glucosidase. However, Fraction III-a gave rhamnose by hydrolysis with maltase in a phosphoric buffer solution (pH 6.4) at  $37^\circ\text{C}$  for 1.5 hr.

Fractions III-b and III-c in Table 2 were hydrolyzed with 0.1 N sulfuric acid solution as Fraction III-a and then chromatographed. The hydrolysates from both fractions gave two spots corresponding to xylose and rhamnose by paper chromatography.

The author is indebted to Professor T. Matsuura of Hiroshima University for his continued advice and encouragement, to Professor C. Araki and Dr. K. Arai of Kyoto Tech. University for their helpful discussion and generous gift of the authentic samples.

16) E. Fischer, M. Bergmann and A. Rabe, *Ber.*, **53**, 2362 (1920).

17) L. Hough, J. K. N. Jones and W. H. Wadman, *J. Chem. Soc.*, **1949**, 2511.