

Mechanism of Decomposition of Carboxylic Sulfurous Anhydrides^{*1}

Michio KOBAYASHI, Hiroshi MINATO and Yoshikuni OGI

Department of Chemistry, Tokyo Metropolitan University, Setagaya, Tokyo




(Received September 11, 1969)

Ethyl benzoyloxysulfinate, prepared from ¹⁸O-labeled benzoic acid and ethyl chlorosulfite, was decomposed and the products were analyzed. The ¹⁸O analyses showed that; 1) prior to decomposition the ¹⁸O of the mixed anhydride was scrambled due to the formation and the internal return of an ion-pair, $\text{C}_6\text{H}_5\text{C}^+-\text{OSOEt}$; 2) ethyl benzoate, benzoic anhydride and diethyl

sulfite were produced *via* an ionic chain mechanism involving successive attack of ethoxide anion and benzoate anion on the carbonyl carbon and sulfinyl sulfur of the mixed anhydride.

In the thermal decomposition of mixed anhydride (I), formed from a carboxylic acid and an alkyl hydrogen sulfite, products are sulfur dioxide, a carboxylic ester, a carboxylic anhydride, and dialkyl sulfite, as shown in Eqs. (1) and (2).¹⁾ In a previous paper²⁾ we described the isolation and the infrared spectra of I with various R and R', and discussed the mechanism of the formation of a carboxylic ester by use of ¹⁸O-labeled methyl and phenyl benzoyloxysulfonates.

In order to clarify the mechanism of the formation of all the decomposition products, the ¹⁸O contents of all the products from the decomposition




of ¹⁸O-labeled ethyl benzoyloxysulfinate (Ia, R = -Ph, R' = -C₂H₅) have been analyzed in detail, and the results are presented in this paper.


Results and Discussion

When ¹⁸O-labeled sodium benzoate and ethyl chlorosulfite were mixed and the sirupy mixture formed was heated at 110–120°C, it decomposed

with vigorous evolution of sulfur dioxide. The sulfur dioxide was collected in a dry ice-acetone trap, and was converted to benzyl phenyl sulfone for the ¹⁸O analysis according to the equation.



After the decomposition was complete, the products were separated by repeated fractional distillation. In order to distinguish the carbonyl oxygen and the ether oxygen in the benzoic anhydride, the anhydride was converted to benzoic acid and benzanilide by reaction with aniline in dichloromethane as shown in the equation.⁵⁾


TABLE 1. SPECIFICITY OF ¹⁸O-LABELING

Sample	% ¹⁸ O-excess	% ¹⁸ O-calcd
Starting PhCO ₂ Na	1.30	(1.30) 1
PhCOOC ₂ H ₅ *	0.67	0.65 1/2
(EtO) ₂ SO*	0.31	0.22 1/6
PhSO ₂ CH ₂ Ph (←SO ₂ *)	0.60	0.65 1/2
PhCOCPh*	1.00	1.08 5/6
PhCO ₂ H**	1.00	1.07 13/16
PhCONHPh**	1.00	1.14 7/8




* Direct reaction products.

** Derived from benzoic anhydride.

^{*1} Organic Sulfur Compound. XX.1) M. Kobayashi and A. Yamamoto, This Bulletin, **39**, 961 (1966).2) M. Kobayashi and R. Kiritani, *ibid.*, **39**, 1782 (1966).3) D. B. Denney and M. A. Greenbaum, *J. Amer. Chem. Soc.*, **79**, 3701 (1957).



The attack of the ethoxide anion on the sulfinyl sulfur atom of Ia will yield ethyl sulfite and benzoate anion, and the attack of the benzoate anion thus formed on the carbonyl carbon of Ia will yield benzoic anhydride, according to Eqs. (11) and (12). The analyses of the ^{18}O content of the products should determine whether or not Eq. (11) or (12) represent the reaction taking place in this medium. The benzoic anhydride was converted to benzanilide and benzoic acid according to Eq. (13). The result of the ^{18}O analyses of benzoic anhydride, benzoic acid, ethyl sulfite and sulfur dioxide (see Table 1) shows that Eqs. (11) and (12) actually represent the reaction taking place in this medium.



Experimental

^{18}O -Labeled Benzoic Acid. According to a previous method²⁾ a mixture of 25 g of benzotrichloride and of H_2^{18}O (1.30 excess- ^{18}O atm %) was refluxed and shaken for 20 hr in a bath at 110°C . After the hydrogen shaken for 20 hr in a bath at 110°C . After the hydrogen chloride formed and excess water were removed under reduced pressure, and the crystals of the benzoic acid remaining were converted to sodium benzoate by addition of the equimolar amount of a sodium hydroxide solution. The sodium salt was recrystallized from distilled water, ground to fine powder, and then dried in a vacuum at 100°C for 2 hr. Yield 14.2 g.

Ethyl Chlorosulfite. To 100 g of thionyl chloride 35.8 g of methanol was slowly added while the temperature was kept low with ice-cooling. The mixture was allowed to stand at room temperature for three days. Distillation under reduced pressure gave 61.5 g of ethyl chlorosulfite (bp $34.5^\circ\text{C}/23$ mmHg). Yield 61.5%.

Thermal Decomposition of Ethyl Benzoyloxy-sulfinate (Ia). When 17.85 g of ethyl chlorosulfite

was added to 10 g of well-dried sodium benzoate- ^{18}O , a slightly exothermic reaction occurred. The milky reaction mixture was then heated at 110 – 130°C for 2 hr. The sulfur dioxide which evolved vigorously during this period was dried with calcium chloride and introduced into an ethereal solution of phenylmagnesium bromide.⁵⁾ The yield of sodium benzenesulfinate was 6.63 g. After the salt was recrystallized from absolute methanol and dried, it was converted to phenyl benzyl sulfone by heating it with benzyl bromide in a sealed tube at 110°C for 5 hr.

The residual reaction mixture was refluxed at 190°C for 2 hr to complete the decomposition. The cooled reaction mixture was treated with 40 ml of methylene chloride, and the sodium chloride precipitated was filtered off. After the solvent was removed in a vacuum, repeated fractional distillation of the residue under reduced pressure yielded 2.1 g of diethyl sulfite (bp $68^\circ\text{C}/30$ mmHg), 0.7 g of ethyl benzoate (bp 105 – $108^\circ\text{C}/30$ mmHg), and 2.1 g of benzoic anhydride (bp 190 – $200^\circ\text{C}/30$ mmHg).

The benzoic anhydride obtained above was converted to benzanilide and benzoic acid by allowing a mixture of 1.0 g of benzoic anhydride and 0.5 g of aniline to stand at room temperature for a few minutes. White crystals of benzanilide which precipitated were filtered off and recrystallized from ethanol; yield, 500 mg. The filtrate was shaken well with a sodium carbonate solution, and the aqueous layer yielded, upon acidification with a diluted hydrochloric acid, crystals of benzoic acid, which were recrystallized from *n*-hexane; yield, 300 mg.

Reaction between Ia and Piperidine. To 500 ml of a carbon tetrachloride solution containing 2 g of benzoic acid and 2.1 g of ethyl chlorosulfite was added 2 g of triethylamine drop by drop with ice-cooling. When 1.7 g of piperidine was added to this mixture, the solution turned pale yellow. After it was allowed to stand overnight at room temperature, 50 ml of water was added to remove triethylamine hydrochloride. The organic layer separated was dried over magnesium sulfate. After the solution was removed in a vacuum, the residual yellow-brown liquid was distilled under reduced pressure and 0.5 g of an oil boiling at $90^\circ\text{C}/5$ mmHg was obtained, which was found to be II by comparison of its infrared spectrum with that of the authentic sample prepared from ethyl chlorosulfate and piperidine.

Reaction between Ia and *p*-Chloroaniline. To 30 ml of benzene solution containing 2 g of benzoic acid and 2.1 g of ethyl chlorosulfite was added 2 g of triethylamine with ice-cooling. The triethylamine hydrochloride precipitated was filtered off, and 30 ml of a carbon tetrachloride solution containing 2 g of *p*-chloroaniline was added to the filtrate. After the mixture was allowed to stand at room temperature for one hour, white crystals of *p*-chlorobenzanilide precipitated; yield, 130 mg.

Isolation of Ia. When 5 g of well-dried sodium benzoate and 10 g of ethyl chlorosulfite were mixed, an exothermic reaction takes place and the mixture became sirupy. Distillation in a high vacuum yielded a crystalline distillate at $94^\circ\text{C}/4 \times 10^{-3}$ mmHg, which was found to be Ia on the basis of its infrared spectrum and the finding that upon addition of an ethereal solution of

5) C. F. H. Allen and S. Converse, "Organic Syntheses," Coll. Vol. I, 226 (1956).

p-chloroaniline it yielded crystalline *p*-chlorobenzanilide.

¹⁸O-Analyses. The ¹⁸O analyses were carried out according to the method of Rittenberg and Ponticorvo.⁶⁾

6) D. Rittenberg and L. Ponticorvo, *Int. J. Appl. Rad. Isotopes*, **1**, 208 (1956).

Each sample was pyrolyzed in the presence of mercuric chloride and mercuric cyanide at 400°C for 4 hr, and the ¹⁸O atom % of the carbon dioxide produced was determined by use of a Hitachi RMU-6D type mass spectrometer.