



4) There has been recently reported by Dornow that α -anilino- α -hydroxyiminoacetophenone was obtained in good yield by the reaction of α -bromo- α -hydroxyiminoacetophenone with aniline. A. Dornow and W. Sassenberg, *Ann. Chem.*, **594**, 185 (1955).

phenone when it is treated with base. Similarly, by the reaction of α -bromo- α -hydroxyiminoacetophenone (IVa) with sodium salt of dibenzoylmethane, dibenzoylfuroxane (VIII) and dibenzoylmethane (IX) were obtained in 62% and 90% yields. In this reaction, sodium salt of dibenzoylmethane behaved as a base to accept hydrogen bromide from α -bromo- α -hydroxyiminoacetophenone instead of a nucleophilic reagent.




Next, the reactions of pyridinium bromides with isopropyl nitrite were tried with the expectation that the nucleophilic displacement by bromide ion at α -carbon of pyridinium salt would not occur because of the lower activity of α -carbon of the pyridinium salts than that of the sulfonium salts described before. When a mixture of phenacylpyridinium bromide (Xa) and isopropyl nitrite (II) in ethanol was allowed to stand at room temperature, the crystalline, (α -hydroxyimino-phenacyl)pyridinium bromide (XIa) was obtained in a quantitative yield. Similarly (α -hydroxyiminoacetyl)pyridinium bromide (XIb) was quantitatively obtained by the reaction of acetylpyridinium bromide (Xb) with isopropyl nitrite (II). They were identified by the IR spectra and the elemental analyses. Whereas, ethoxycarbonylhydroxyiminomethylpyridinium bromide could not be isolated by the reaction of ethoxycarbonylmethylpyridinium bromide with isopropyl nitrite and the starting materials were recovered quantitatively.



Then, the reactions of (α -hydroxyiminophenacyl)-pyridinium bromide, produced by the above mentioned reaction, with nucleophilic reagents were investigated. When a mixture of (α -hydroxyiminophenacyl)pyridinium bromide (XIa) and sodium thiophenolate in ethanol was allowed to stand at room temperature, α -hydroxyimino- α -phenylthioacetophenone (VI) was obtained in 71% yield. This was identical with that prepared by the reaction of α -bromo- α -hydroxyiminoacetophenone and sodium thiophenolate. Similarly, α -anilino- α -hydroxyiminoacetophenone (XII) was produced in 65% yield by the reaction of (α -hydroxyiminophenacyl)-pyridinium bromide (XIa) with two equimolar amounts of aniline in ethanol at room tem-

perature. These reactions indicate that pyridine can be easily eliminated from the pyridinium salt by the nucleophilic displacement of the nucleophiles, such as aniline and sodium thiophenolate.



Experimental

Reaction of Dimethylphenacetyl-sulfonium Bromide with Isopropyl Nitrite.

A mixture of dimethylphenacetyl-sulfonium bromide (1.31 g, 0.005 mol) and excess isopropyl nitrite (0.89 g, 0.01 mol) was stirred in dichloromethane (100 ml) for 12 hr at room temperature. The evolution of dimethyl sulfide was observed by its characteristic odor accompanied with disappearance of the crystals of the sulfonium salt. Evaporation of the solvent under reduced pressure gave white crystals, α -bromo- α -hydroxyiminoacetophenone (mp 138–139°C), 1.07 g (93%), recrystallized from cyclohexane. Similarly, various α -bromo- α -hydroxyimino compounds were obtained by the reactions of sulfonium bromides with isopropyl nitrite. The results are listed in Table 1. The α -bromo- α -hydroxyimino compounds were identified by their IR spectra and elemental analyses.

TABLE 1. THE REACTION OF SULFONIUM BROMIDES WITH ISOPROPYL NITRITE

Sulfonium bromide	Reaction time	Solvent	Yield	Mp, °C
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{S}^+-\text{CH}_2-\text{C}-\text{C}_6\text{H}_5 \\ \\ \text{Br}^- \\ \text{O} \end{array} $	12 hr	CH ₂ Cl ₂	93%	138–139
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{S}^+-\text{CH}_2-\text{C}-\text{CH}_3 \\ \\ \text{Br}^- \\ \text{O} \end{array} $	1 day	CH ₂ Cl ₂	quant.	119–120
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{S}^+-\text{CH}_2-\text{C}-\text{COC}_2\text{H}_5 \\ \\ \text{Br}^- \\ \text{O} \end{array} $	1 day	CH ₂ Cl ₂	quant.	92–93

Reaction of α -Bromo- α -hydroxyiminoacetophenone with Pyridine.

Into a solution of α -bromo- α -hydroxyiminoacetophenone (0.43 g, 0.0019 mol) in ether (50 ml), a solution of pyridine (0.15 g, 0.0019 mol) in ether (10 ml) was added drop by drop with stirring. After stirring for 30 min at room temperature, white precipitate deposited. The precipitate was collected by filtration and recrystallized from ethanol to give (α -hydroxyiminophenacyl)pyridinium bromide (mp 143.5°C (dec.)) in a quantitative yield.

Found: C, 51.07; H, 3.36; N, 9.31%. Calcd for C₁₃H₁₁BrN₂O₂: C, 50.83; H, 3.61; N, 9.12%.

Reaction of α -Bromo- α -hydroxyiminoacetophenone with Sodium Thiophenolate.

A solution of sodium thiophenolate (0.66 g, 0.005 mol) in ethanol (50 ml) was added dropwise to a solution of α -bromo- α -hydroxyiminoacetophenone (1.14 g, 0.005 mol) in ethanol (50 ml) with stirring. After stirring for 12 hr at room temperature, solvent was evaporated. The residue was washed with water (100 ml) and extracted with ether (100 ml). Then, the solvent was evaporated under reduced pressure to give yellow crystals. Recrystallization from cyclohexane gave α -hydroxyimino- α -phenylthioacetophenone (mp 110–111°C) in an almost quantitative yield.

Found: C, 65.49; H, 4.56; N, 5.35; S, 12.20%. Calcd for $C_{14}H_{11}NO_2S$: C, 65.36; H, 4.31; N, 5.45; S, 12.44%.

Reaction of α -Bromo- α -hydroxyiminoacetophenone with Triethylamine.

Into a solution of α -bromo- α -hydroxyiminoacetophenone (1.12 g, 0.005 mol) in ether (50 ml), an equimolar amount of triethylamine was added with stirring. The reaction mixture was stirred for 1 day at room temperature. After removal of the solvent, the residue was chromatographed on silica gel and elution with benzene gave dibenzoylfuroxane (mp 86–87°C, 48%).

Found: C, 65.51; H, 3.25; N, 9.27%. Calcd for $C_{16}H_{10}N_2O_4$: C, 65.30; H, 3.43; N, 9.52%.

Reaction of Phenacylpyridinium Bromide with Isopropyl Nitrite. A mixture of phenacylpyridinium bromide (2.78 g, 0.01 mol) and excess amount of isopropyl nitrite (1.78 g, 0.02 mol) in ethanol (50 ml) was stirred for 1 day at room temperature. After removal of the solvent, white crystals were obtained.

Recrystallization from ethanol gave (α -hydroxyiminophenacyl)pyridinium bromide (mp 143.5°C (dec.)) in a quantitative yield. By a similar procedure, (α -hydroxyiminoacetyl)pyridinium bromide (mp 73.5–74.0°C) was obtained quantitatively.

Found: C, 39.48; H, 3.59; N, 11.36%. Calcd for $C_8H_9BrN_2O_2$: C, 39.20; H, 3.70; N, 11.43%.

Reaction of (α -Hydroxyiminophenacyl)pyridinium Bromide with Sodium Thiophenolate.

To a suspension of (α -hydroxyiminophenacyl)pyridinium bromide (3.07 g, 0.01 mol) in ethanol (50 ml), a solution of sodium thiophenolate (1.32 g, 0.01 mol) in ethanol (20 ml) was added with stirring. After stirring for 12 hr, crystals of (α -hydroxyiminophenacyl)pyridinium bromide disappeared. The residue was chromatographed on silica gel after removal of the solvent under reduced pressure, and elution with dichloromethane gave yellow crystals, α -hydroxyimino- α -phenylthioacetophenone (mp 110–111°C), 1.82 g (71%).

Reaction of (α -Hydroxyiminophenacyl)pyridinium Bromide with Aniline.

A mixture of (α -hydroxyiminophenacyl)pyridinium bromide (3.07 g, 0.01 mol) and two equimolar amounts of aniline (1.86 g, 0.02 mol) in ethanol (100 ml) was stirred for 5 days at room temperature. After removal of the solvent, the residue was washed with water (50 ml) and extracted with ether (50 ml). Then, the solvent was evaporated and crystalline solid of α -anilino- α -hydroxyiminoacetophenone (mp 145.0–146.5°C) was obtained in 65% yield after removal of the solvent under reduced pressure.