

BULLETIN OF THE CHEMICAL SOCIETY OF JAPAN, VOL. 44, 858—859 (1971)

Syntheses of Fused Heterocycles *via* Cycloaddition of Hetaryne. Studies on Heteroaromaticity. Part XLVII¹⁾

Tadashi SASAKI, Ken KANEMATSU, and Masayuki UCHIDE

Institute of Applied Organic Chemistry, Faculty of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya

(Received October 14, 1970)

In comparison with abundant examples for the cycloaddition reactions of dehydrobenzene (benzyne), only occasional reports have been made concerning the same reactions of dehydroaromatic heterocycles (hetarynes).²⁾ Recently, the electronic structures of hetarynes have been discussed using the extended Hückel theory,³⁾ by which the larger relative stability is predicted for 3-pyridyne compared with other pyridyne isomers.

These facts prompted us to report one-step syntheses of the fused bicyclic heteroaromatic ring system by the cycloaddition reactions of this hetaryne.

Expecting that the intermediate 3-pyridyne (II) might be trapped with dipolarophiles or dienophiles, the lead tetraacetate oxidation of 1-aminotriazolo-[4,5-*c*]pyridine (I)⁴⁾ was carried out in the presence of dienophilic five-membered heterocycles or some dipolarophiles.

These results are described in this note.

Results and Discussion

Only one example of 1,2-cycloaddition of dehydroheterocycle is recorded;⁵⁾ cyclopentadiene adds to 3-pyridyne in a 1,2-fashion rather than in a 1,4-fashion of Diels-Alder reaction. On the other hand, attempts to add 3-pyridyne to anthracene remain unsuccessful.⁶⁾

In contrast, lead tetraacetate oxidation of compound I was carried out in the presence of dimethylfulvene, norbornadiene, and furan at room temperature for 1 hr. Only the *endo*-oxide III (picrate, mp 178°C) was obtained from furan in a 74% yield, which was identified by the IR and NMR spectral comparison with those of an authentic sample prepared by heating 3-pyridinediazonium-4-carboxylate in furan.⁷⁾ A recent report described the lead tetraacetate oxidation of I in the presence of tetracyclone which affords 5,6,7,8-tetraphenylisoquinoline.⁴⁾ Neither 1,2-cycloadducts nor 1,4-Diels-Alder adducts could be isolated from the reaction of 3-pyridyne with dimethylfulvene or norbornadiene.

Similarly, 1,3-dipolar cycloaddition reactions of 3-pyridyne with dipolarophiles were also attempted.




Chart 1.

There is only one example²⁾ known for a 1,3-addition of azide to a dehydroheterocyclic intermediate as shown by the following scheme.




Chart 2.

Treatment of a mixture of I and *N*-phenylsydnone in 1 : 1.3 mol ratio with lead tetraacetate in tetrahydrofuran at room temperature led to the formation of 1,3-dipolar cycloadduct IV, mp 132–133°C, in a 30% yield. The NMR spectrum of IV shows signals of the ring protons at τ 0.60 (1H, s), 1.52 (1H, s), 1.77 (1H, d, $J=7.0$ Hz), and overlapping signals at τ 1.96–2.51 (1H-ring proton and 5H-phenyl protons, m). Similar oxidation of a mixture of compound I and benzaldehyde phenylhydrazone^{8,9)} in 1 : 1.1 mol ratio with an excess of lead tetraacetate afforded the expected cycloadduct V as pale yellow oil (picrate, mp 234–237°C) in a 15% yield. However, attempted cycloaddition of diphenyldiazomethane or benzonitrile oxide with 3-pyridyne was unsuccessful.

The orientation in the addition reaction of 1,3-dipolar *N*-phenylsydnone or diphenylnitrilimine with 3-pyridyne is not clear at present. However, the nucleophilic attack of the dipolar on 3-pyridyne might occur preferentially at the C-4 position as a consequence of its lower total electron density (3.97 against 4.35 electrons)³⁾ as shown by the following chart.

The structures of the 1,3-dipolar cycloadducts IV and V were tentatively assigned as 2-phenyl-2*H*-pyrazolo[4,3-*c*]pyridine and 1,3-diphenyl-1*H*-pyrazolo-

1) Part of series, XLVI., T. Sasaki and T. Yoshioka, *This Bulletin*, **43**, 2989 (1970).

2) R. W. Hoffmann, "Dehydrobenzene and Cycloalkynes," Academic Press, New York, N. Y. (1967), p. 199, 303, 306.

3) W. Adams, A. Grimison, and R. Hoffmann, *J. Amer. Chem. Soc.*, **91**, 2570 (1969).

4) G. W. J. Fleet and I. Fleming, *J. Chem. Soc., C*, **1969**, 1758.

5) T. Kauffmann, J. Hansen, K. Udeloft, and R. Winthwein, *Angew. Chem.*, **76**, 590 (1964).

6) T. Kauffmann, *ibid.*, **77**, 557 (1965).

7) T. Kauffmann and F. P. Boettcher, *Chem. Ber.*, **95**, 949 (1962).

8) The nitrilimines have been trapped during the oxidation of aldehyde hydrazones; W. A. F. Gladstone, J. B. Aylward, and R. O. C. Norman, *J. Chem. Soc., C*, **1969**, 2587.

9) T. Sasaki and T. Yoshioka, *This Bulletin*, **43**, 1254 (1970).

[4,3-*c*]pyridine, respectively. Such a fused ring system as pyrazolo[4,3-*c*]pyridine is the first in the fused heterocyclic series.

Attempted 1,3-dipolar cycloaddition reactions of 3-pyridinediazonium-4-carboxylate with dipolarophilic nitrilimines were unsuccessful.




Chart 3.

Thus, it can be concluded that the cycloaddition reactivity of 3-pyridyne is generally much lower than that of benzyne, since the stereospecific 2+4 cycloaddition, the 2+2 nonconcerted cycloaddition, and

the 1,3-cycloaddition for benzyne have been reported to give the corresponding cycloadducts in excellent yields.^{2,10)}

Experimental¹¹⁾

Lead Tetraacetate Oxidation of 1-Aminotriazolo[4,5-*c*]pyridine in the Presence of Furan. Compound I (390 mg) was dissolved in furan (5.1 g) and this solution was dropwise added to a stirred suspension of lead tetraacetate (1.4 g) in dichloromethane (20 ml) at 20°C. The reaction was always instantaneous and exothermic with evolution of nitrogen. Work-up involved filtration, evaporation of the solution, and chromatography (silicic acid, 100 mesh, Mallinckrodt) of the residue using benzene as an eluent.

Colorless oil of an *endo*-oxide III (317 mg, 75%) was obtained from the eluent; picrate, mp 178°C (lit.⁷⁾ mp 178°C).

Found: C, 48.28; H, 2.69; N, 14.97%. Calcd for C₁₅H₁₀O₈N₄ (picrate): C, 48.13; H, 2.69; N, 14.97%.

Reaction of 3-Pyridyne with *N*-Phenylsydnone. A solution of compound I (400 mg) and *N*-phenylsydnone (670 mg) in tetrahydrofuran (40 ml) was added dropwise to a stirred suspension of lead tetraacetate (1.2 g) in tetrahydrofuran (20 ml) at 20–25°C. After 1 hr, work-up as described above afforded an adduct IV (168 mg, 30%) as colorless prisms, mp 132–133°C.

Found: C, 73.56; H, 4.71; N, 21.74%. Calcd for C₁₂H₈N₃: C, 73.83; H, 4.65; N, 21.53%.

Reaction of 3-Pyridyne with 1,3-Diphenylnitrilimine. A solution of benzaldehyde phenylhydrazone (226 mg) and I (135 mg) in dichloromethane (20 ml) was added during 50 min to a stirred suspension of lead tetraacetate (1.5 g) in dichloromethane (50 ml) at below 10°C. The resulting thick suspension was further stirred at room temperature for 1 hr. Work-up as above afforded an adduct V (80 mg) as pale yellow oil; picrate, mp 234–237°C.

Found: C, 57.86; H, 3.29; N, 16.51%. Calcd for C₂₄H₁₆N₆O₇ (picrate): C, 57.60; H, 3.22; N, 16.80%.

10) For leading reference, see M. Jones, Jr., and R. H. Levin, *J. Amer. Chem. Soc.*, **91**, 6411 (1969).

11) The melting points were measured with a Yanagimoto micromelting point apparatus and are uncorrected. Microanalyses were performed on a Yanagimoto C.H.N.-Corder, Model MT-1. The NMR spectra were taken with a JEOLCO., Model JNM-MH-60 NMR spectrometer and with a Varian A-60 spectrometer with tetramethylsilane as an internal standard. The chemical shifts are expressed in τ values. The IR spectra were taken with a JASCO Model IR-S spectrometer.