

BULLETIN OF THE CHEMICAL SOCIETY OF JAPAN, VOL. 44, 1677—1682 (1971)

**Substituted Benzopyranopyridine and Pyrimidine Ring Syntheses by the Ternary Condensation of Ethyl Cyanoacetate, Salicylaldehyde, and Certain Aldehydes in the Presence of Ammonium Acetate**

Akio SAKURAI, Hiroshi MIDORIKAWA, and Yasuo HASHIMOTO\*

*The Institute of Physical and Chemical Research, Wako-shi, Saitama*

*\*Tokyo Electrical Engineering College, Nishikicho, Chiyoda-ku, Tokyo*

(Received December 5, 1970)

Various 1-substituted benzopyranopyridines were readily prepared by the condensation of ethyl cyanoacetate and salicylaldehyde (or 3-methoxysalicylaldehyde) with aliphatic aldehydes (propion, *n*-butyl, *n*- and isovaleraldehyde) in the presence of ammonium acetate. On the other hand, condensation with aromatic aldehydes such as benzaldehyde and *o*-, *m*-, or *p*-substituted benzaldehydes gave 2-aryl-benzopyranopyrimidines.


In a previous paper,<sup>1)</sup> the synthesis of 2- or 1,2-

1) A. Sakurai, H. Midorikawa, and Y. Hashimoto, This Bulletin, **43**, 2925 (1970).

disubstituted benzopyranopyridines by the condensation of ethyl cyanoacetate, salicylaldehyde, and ketones in the presence of ammonium acetate has been reported.

The present paper will deal with the syntheses of 1-alkylbenzopyranopyridines (from aliphatic aldehyde) and 2-arylbenzopyranopyrimidines (from aromatic aldehyde) by the reaction of ethyl cyanoacetate and salicylaldehyde or 3-methoxysalicylaldehyde with various aldehydes in the presence of ammonium acetate.


The reaction of ethyl cyanoacetate and salicylaldehyde or 3-methoxysalicylaldehyde with aliphatic aldehydes was carried out by refluxing a mixture of the ester,



Scheme 1

salicylaldehyde, aliphatic aldehyde, and ammonium acetate (molar ratio 1:1:1-1.5) in ethanol. In this case, the condensation of the amino group of 3-amidinocoumarin<sup>1)</sup> (I) with the carbonyl group of the aldehyde, cyclization between the methylene group adjacent to the aldehyde carbonyl and the 4-position of the coumarin ring, and subsequent dehydrogenation took place to form 1-alkyl-4-amino-5-oxo-[1]-benzopyrano[3,4-c]pyridines (II).

However, when isobutyraldehyde was employed as the aldehyde reactant, the corresponding product of the type II was not obtained. The NMR spectrum (in  $\text{CF}_3\text{CO}_2\text{H}$ ) of IIa (from propionaldehyde) showed only a methyl singlet at 2.9 ppm, while IIb (from *n*-butyraldehyde) showed a methyl triplet at 1.55 ppm and a methylene quartet at 3.35 ppm in a higher field than the aromatic-ring region. These facts indicated that the methylene group adjacent to the aldehyde carbonyl was involved in this cyclization. On the other hand, the condensation with aromatic aldehydes (benzaldehyde or its derivatives) gave 2-aryl-4-amino-5-oxo-[1]benzopyrano[4,3-*d*]pyrimidines (IV). In this case, 1 mol each of ethyl cyanoacetate, salicylaldehyde, and benzaldehyde (or its derivatives) reacted with 2 mol of ammonia to give IV, with the elimination of 1 mol of ethanol and 2 mol each of water and hydrogen. The compound IVa was also obtained by the condensation of 3-amidinocoumarin or 3-cyanocoumarin with benzaldehyde in the presence of ammonium acetate



Scheme 2




Table 1  
1-Alkyl-4-amino-5-oxo-[1]benzopyrano [3,4-*C*] pyridine (II)

|     | R                | R'                                      | MP, °C  | Yield, % | Formula                                                       | Calcd, % |      |       | Found, % |      |       |
|-----|------------------|-----------------------------------------|---------|----------|---------------------------------------------------------------|----------|------|-------|----------|------|-------|
|     |                  |                                         |         |          |                                                               | C        | H    | N     | C        | H    | N     |
| IIa | H                | CH <sub>3</sub>                         | 221-222 | 34       | C <sub>13</sub> H <sub>10</sub> O <sub>2</sub> N <sub>2</sub> | 69.01    | 4.46 | 12.38 | 68.93    | 4.66 | 12.36 |
| b   | H                | C <sub>2</sub> H <sub>5</sub>           | 207-209 | 19       | C <sub>14</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub> | 69.99    | 5.03 | 11.66 | 69.92    | 5.07 | 11.66 |
| c   | H                | <i>n</i> -C <sub>3</sub> H <sub>7</sub> | 186-187 | 20       | C <sub>15</sub> H <sub>14</sub> O <sub>2</sub> N <sub>2</sub> | 70.85    | 5.55 | 11.02 | 70.32    | 5.57 | 10.80 |
| d   | H                | <i>i</i> -C <sub>3</sub> H <sub>7</sub> | 176-177 | 5        | C <sub>15</sub> H <sub>14</sub> O <sub>2</sub> N <sub>2</sub> | 70.85    | 5.55 | 11.02 | 70.87    | 5.41 | 11.16 |
| e   | H                | <i>n</i> -C <sub>4</sub> H <sub>9</sub> | 173-174 | 20       | C <sub>16</sub> H <sub>16</sub> O <sub>2</sub> N <sub>2</sub> | 71.62    | 6.01 | 10.44 | 71.74    | 5.93 | 10.42 |
| f   | OCH <sub>3</sub> | CH <sub>3</sub>                         | 212-215 | 35       | C <sub>14</sub> H <sub>12</sub> O <sub>3</sub> N <sub>2</sub> | 65.62    | 4.72 | 10.93 | 65.37    | 4.50 | 10.96 |
| g   | OCH <sub>3</sub> | C <sub>2</sub> H <sub>5</sub>           | 214-215 | 19       | C <sub>15</sub> H <sub>14</sub> O <sub>3</sub> N <sub>2</sub> | 66.65    | 5.22 | 10.37 | 66.40    | 5.24 | 10.30 |
| h   | OCH <sub>3</sub> | <i>n</i> -C <sub>3</sub> H <sub>7</sub> | 200-202 | 20       | C <sub>16</sub> H <sub>16</sub> O <sub>3</sub> N <sub>2</sub> | 67.59    | 5.67 | 9.85  | 67.57    | 5.69 | 9.71  |
| i   | OCH <sub>3</sub> | <i>i</i> -C <sub>3</sub> H <sub>7</sub> | 211-212 | 12       | C <sub>16</sub> H <sub>16</sub> O <sub>3</sub> N <sub>2</sub> | 67.59    | 5.67 | 9.85  | 67.10    | 5.64 | 9.78  |
| j   | OCH <sub>3</sub> | <i>n</i> -C <sub>4</sub> H <sub>9</sub> | 181-182 | 19       | C <sub>17</sub> H <sub>18</sub> O <sub>3</sub> N <sub>2</sub> | 68.44    | 6.08 | 9.39  | 68.29    | 6.03 | 9.29  |




Table 2  
1-Alkyl-4, 4-diacetamino-5-oxo-[1] benzopyrano [3,4-C] pyridine (III)

|      | R                | R'                              | MP, °C  | Yield, % | Formula                                                       | C     | Calcd, %<br>H | N    | C     | Found, %<br>H | N    |
|------|------------------|---------------------------------|---------|----------|---------------------------------------------------------------|-------|---------------|------|-------|---------------|------|
| IIIa | H                | CH <sub>3</sub>                 | 176-179 | 55       | C <sub>17</sub> H <sub>14</sub> O <sub>4</sub> N <sub>2</sub> | 65.80 | 4.55          | 9.03 | 65.50 | 4.64          | 9.21 |
| b    | H                | C <sub>2</sub> H <sub>5</sub>   | 132-135 | 86       | C <sub>18</sub> H <sub>16</sub> O <sub>4</sub> N <sub>2</sub> | 66.66 | 4.97          | 8.64 | 66.96 | 5.02          | 8.86 |
| c    | H                | n-C <sub>3</sub> H <sub>7</sub> | 104-108 | 88       | C <sub>19</sub> H <sub>18</sub> O <sub>4</sub> N <sub>2</sub> | 67.44 | 5.36          | 8.28 | 67.66 | 5.28          | 8.42 |
| d    | OCH <sub>3</sub> | CH <sub>3</sub>                 | 228-230 | 45       | C <sub>18</sub> H <sub>16</sub> O <sub>5</sub> N <sub>2</sub> | 63.52 | 4.74          | 8.23 | 63.40 | 4.69          | 8.37 |
| e    | OCH <sub>3</sub> | C <sub>2</sub> H <sub>5</sub>   | 177-180 | 89       | C <sub>19</sub> H <sub>18</sub> O <sub>5</sub> N <sub>2</sub> | 64.40 | 5.12          | 7.91 | 64.48 | 5.33          | 8.05 |
| f    | OCH <sub>3</sub> | n-C <sub>3</sub> H <sub>7</sub> | 160-163 | 90       | C <sub>20</sub> H <sub>20</sub> O <sub>5</sub> N <sub>2</sub> | 65.21 | 5.47          | 7.61 | 65.54 | 5.52          | 7.92 |

TABLE 3. INFRARED SPECTRAL DATA FOR THE COMPOUNDS II

|     | $\nu$ NH <sub>2</sub> | $\nu$ C=O | $\delta$ NH <sub>2</sub> | Aromatic ring          | $\delta$ CH |
|-----|-----------------------|-----------|--------------------------|------------------------|-------------|
| IIa | 3430, 3270, 3150      | 1700      | 1625                     | 1610, 1590, 1570, 1545 | 760         |
| b   | 3420, 3260, 3130      | 1690      | 1620                     | 1610, 1590, 1565, 1540 | 755, 745    |
| c   | 3410, 3270, 3130      | 1700      | 1625                     | 1610, 1590, 1570, 1545 | 765, 750    |
| d   | 3420, 3260, 3120      | 1720      | 1630                     | 1610, 1590, 1565, 1545 | 765, 750    |
| e   | 3410, 3260, 3120      | 1700      | 1630                     | 1610, 1570, 1550       | 770         |
| f   | 3420, 3270, 3140      | 1695      | 1620                     | 1610, 1590, 1570, 1550 | 780         |
| g   | 3410, 3260, 3130      | 1700      | 1625                     | 1610, 1590, 1570, 1550 | 785         |
| h   | 3410, 3260, 3140      | 1695      | 1620                     | 1610, 1570, 1545       | 785         |
| i   | 3400, 3260, 3150      | 1695      | 1620                     | 1610, 1590, 1565, 1540 | 780         |
| j   | 3410, 3260, 3140      | 1700      | 1625                     | 1610, 1590, 1570, 1550 | 785         |

TABLE 4. INFRARED SPECTRAL DATA<sup>a)</sup> FOR THE COMPOUNDS III

|      | $\nu$ O-C=O | $\nu$ N-C=O | Aromatic ring          | $\delta$ CH |
|------|-------------|-------------|------------------------|-------------|
| IIIa | 1725        | 1700,       | 1610, 1590, 1575, 1540 | 770         |
| b    | 1735        | 1715, 1700  | 1600, 1590, 1570, 1540 | 770         |
| c    | 1735        | 1720, 1695  | 1610, 1590, 1570, 1540 | 770         |
| d    | 1725        | 1705        | 1610, 1590, 1575, 1545 | 785         |
| e    | 1730        | 1685        | 1610, 1590, 1570, 1540 | 790, 785    |
| f    | 1725        | 1695, 1680  | 1610, 1580, 1570, 1540 | 785         |

a) All spectra were taken in potassium bromide disks; cm<sup>-1</sup>

(Scheme 2).

The infrared spectra of IV revealed slightly lower shifted absorption bands for a primary amino group at about 3400 and 3300 cm<sup>-1</sup> and at about 1700 cm<sup>-1</sup> for a coumarin-type carbonyl group. This shows the presence of intramolecular C=O...H<sub>2</sub>N bonding, as is shown in the compounds obtained by condensation with ketones.<sup>1)</sup> When heated with acetic anhydride in pyridine, type IV compounds were converted to their corresponding monoacetylated derivatives (V). On

TABLE 5. NMR SPECTRAL DATA<sup>a)</sup> FOR THE COMPOUNDS II AND III

|      | CH <sub>3</sub>       | OCH <sub>3</sub> | CH <sub>2</sub>             | -CH-           | Ring -CH=                       |
|------|-----------------------|------------------|-----------------------------|----------------|---------------------------------|
| IIa  | 2.9 (s, 3H)           |                  |                             |                | 7.5-8.2 (m, 4H), 8.6-8.8(m, 5H) |
| b    | 1.55(t, 3H)           |                  | 3.35(q, 2H)                 |                | 7.5-8.8 (m, 5H)                 |
| c    | 1.2 (t, 3H)           |                  | 1.9 (sx, 2H), 3.25 (t, 2H), |                | 7.5-8.8 (m, 5H)                 |
| g    | 1.5 (t, 3H)           | 4.1 (s, 3H)      | 3.3 (q, 2H)                 |                | 7.5-7.7, 8.0-8.3 (m, 2H each)   |
| h    | 1.2 (t, 3H)           | 4.12(s, 3H)      | 1.95(sx, 2H), 3.25(t, 2H)   |                | 7.5-7.7, 8.0-8.3 (m, 2H each)   |
| i    | 1.5 (d, 6H)           | 4.1 (s, 3H)      |                             | 3.7-4.3(m, 1H) | 7.4-7.8, 7.9-8.4 (m, 2H each)   |
| IIIb | 1.6 (t, 3H)           | 2.7 (s, 3H each) | 3.5 (q, 2H)                 |                | 7.5-8.1, 8.4-8.8 (m, 5H)        |
|      | 2.4, 2.7 (s, 3H each) |                  |                             |                |                                 |

a) Parts per million downfield from tetramethylsilane in CF<sub>3</sub>CO<sub>2</sub>H; s=singlet, d=doublet, t=triplet, q=quartet, sx=sextet, m=multiplet

the other hand, type II compounds gave diacetylated derivatives (III) under the same reaction conditions. In the infrared spectra of III, carbonyl stretching bands shifted to frequencies higher by from 20 to 30  $\text{cm}^{-1}$  than those of the compounds II. However, the carbonyl bands of V appeared in the region nearly the same

as or slightly lower than those of the compounds IV, and the imino stretching band always appeared at 3250  $\text{cm}^{-1}$ . These data, therefore, can reasonably be explained by the presence of intramolecular hydrogen bonding in the compounds II and IV.




Table 6  
2-Aryl-4-amino-5-oxo-1H-benzopyrano[4,3-d]pyrimidine (IV)

|     | R              | Ar | MP, °C  | Yield, % | Formula                                                                             | C     | Calcd, % |       |  | C     | Found, % |       |  |
|-----|----------------|----|---------|----------|-------------------------------------------------------------------------------------|-------|----------|-------|--|-------|----------|-------|--|
|     |                |    |         |          |                                                                                     |       | H        | N     |  |       | H        | N     |  |
| IVa | H              | Ph | 266-268 | 20       | $\text{C}_{17}\text{H}_{11}\text{O}_2\text{N}_3$                                    | 70.58 | 3.83     | 14.53 |  | 70.33 | 4.01     | 14.47 |  |
| b   | H              |    | 277-280 | 13       | $\text{C}_{18}\text{H}_{13}\text{O}_2\text{N}_3$                                    | 71.27 | 4.32     | 13.86 |  | 71.06 | 4.10     | 13.68 |  |
| c   | H              |    | 211-213 | 8        | $\text{C}_{18}\text{H}_{13}\text{O}_3\text{N}_3$                                    | 67.70 | 4.11     | 13.16 |  | 67.81 | 4.04     | 13.34 |  |
| d   | H              |    | 262-264 | 11       | $\text{C}_{18}\text{H}_{13}\text{O}_3\text{N}_3$                                    | 67.70 | 4.11     | 13.16 |  | 67.77 | 4.12     | 13.11 |  |
| e   | H              |    | 229-231 | 17       | $\text{C}_{20}\text{H}_{17}\text{O}_2\text{N}_3$                                    | 72.49 | 5.17     | 12.68 |  | 72.36 | 4.98     | 12.56 |  |
| f   | H              |    | 318-321 | 17       | $\text{C}_{19}\text{H}_{14}\text{O}_3\text{N}_4$<br>$\frac{1}{2}\text{H}_2\text{O}$ | 61.12 | 4.55     | 15.01 |  | 61.34 | 4.40     | 15.20 |  |
| g   | H              |    | 254-256 | 6        | $\text{C}_{19}\text{H}_{16}\text{O}_2\text{N}_4$                                    | 68.66 | 4.85     | 16.86 |  | 68.42 | 4.51     | 16.89 |  |
| h   | H              |    | 254-257 | 25       | $\text{C}_{17}\text{H}_{10}\text{O}_4\text{N}_4$                                    | 61.08 | 3.02     | 16.76 |  | 61.82 | 3.27     | 16.62 |  |
| i   | H              |    | 225-228 | 10       | $\text{C}_{19}\text{H}_{15}\text{O}_4\text{N}_3$                                    | 65.32 | 4.33     | 12.03 |  | 65.14 | 4.27     | 11.98 |  |
| j   | H              |    | 247-248 | 12       | $\text{C}_{19}\text{H}_{15}\text{O}_4\text{N}_3$                                    | 65.32 | 4.33     | 12.03 |  | 65.47 | 4.23     | 12.29 |  |
| k   | H              |    | 185-186 | 13       | $\text{C}_{21}\text{H}_{19}\text{O}_4\text{N}_3$                                    | 66.83 | 5.07     | 11.14 |  | 66.80 | 4.55     | 11.18 |  |
| l   | H              |    | 322-324 | 13       | $\text{C}_{17}\text{H}_9\text{O}_3\text{N}_3\text{Br}_2$                            | 44.09 | 1.96     | 9.07  |  | 44.18 | 2.19     | 8.96  |  |
| m   | $\text{OCH}_3$ | Ph | 283-284 | 20       | $\text{C}_{18}\text{H}_{13}\text{O}_3\text{N}_3$                                    | 67.70 | 4.11     | 13.16 |  | 67.43 | 3.97     | 13.06 |  |
| n   | $\text{OCH}_3$ |    | 263-265 | 15       | $\text{C}_{19}\text{H}_{15}\text{O}_3\text{N}_3$                                    | 68.46 | 4.54     | 12.61 |  | 68.07 | 4.55     | 12.35 |  |
| o   | $\text{OCH}_3$ |    | 234-236 | 15       | $\text{C}_{21}\text{H}_{19}\text{O}_3\text{N}_3$                                    | 69.79 | 5.30     | 11.63 |  | 69.50 | 5.11     | 11.48 |  |
| p   | $\text{OCH}_3$ |    | 300-303 | 40       | $\text{C}_{18}\text{H}_{12}\text{O}_5\text{N}_4$                                    | 59.34 | 3.32     | 15.38 |  | 59.30 | 3.24     | 15.83 |  |
| q   | $\text{OCH}_3$ |    | 327-330 | 12       | $\text{C}_{20}\text{H}_{17}\text{O}_5\text{N}_3$                                    | 63.32 | 4.52     | 11.08 |  | 62.52 | 4.24     | 11.59 |  |
| r   | $\text{OCH}_3$ |    | 251-253 | 28       | $\text{C}_{20}\text{H}_{17}\text{O}_5\text{N}_3$                                    | 63.32 | 4.52     | 11.08 |  | 63.06 | 4.38     | 11.30 |  |
| s   | $\text{OCH}_3$ |    | 222-225 | 26       | $\text{C}_{22}\text{H}_{21}\text{O}_5\text{N}_3$                                    | 64.85 | 5.20     | 10.31 |  | 64.91 | 5.17     | 10.46 |  |




Table 7  
2-Aryl-4-acetamino-5-oxo-[1]benzopyrano [4,3-d] pyrimidine (V)

|    | R                | Ar | MP, °C  | Yield, % | Formula                                                       | C     | Calcd, % |       |  | Found, % |      |       |
|----|------------------|----|---------|----------|---------------------------------------------------------------|-------|----------|-------|--|----------|------|-------|
|    |                  |    |         |          |                                                               |       | H        | N     |  | H        | N    |       |
| Va | H                | Ph | 252-253 | 87       | C <sub>19</sub> H <sub>13</sub> O <sub>3</sub> N <sub>3</sub> | 68.87 | 3.96     | 12.68 |  | 68.80    | 3.48 | 12.78 |
| b  | H                |    | 257-260 | 88       | C <sub>20</sub> H <sub>15</sub> O <sub>4</sub> N <sub>3</sub> | 66.47 | 4.18     | 11.63 |  | 66.44    | 4.23 | 11.53 |
| c  | H                |    | 330-332 | 77       | C <sub>21</sub> H <sub>16</sub> O <sub>4</sub> N <sub>4</sub> | 64.94 | 4.15     | 14.43 |  | 64.22    | 4.51 | 14.48 |
| d  | H                |    | 260-263 | 88       | C <sub>19</sub> H <sub>12</sub> O <sub>5</sub> N <sub>4</sub> | 60.64 | 3.21     | 14.89 |  | 59.95    | 3.45 | 15.05 |
| e  | H                |    | 235-237 | 89       | C <sub>22</sub> H <sub>19</sub> O <sub>3</sub> N <sub>3</sub> | 70.76 | 5.13     | 11.25 |  | 70.79    | 4.99 | 11.45 |
| f  | OCH <sub>3</sub> | Ph | 267-268 | 70       | C <sub>20</sub> H <sub>15</sub> O <sub>4</sub> N <sub>3</sub> | 66.47 | 4.18     | 11.63 |  | 66.19    | 4.45 | 11.75 |
| g  | OCH <sub>3</sub> |    | 303-305 | 89       | C <sub>20</sub> H <sub>14</sub> O <sub>6</sub> N <sub>4</sub> | 59.11 | 3.47     | 13.79 |  | 58.69    | 3.71 | 13.85 |
| h  | OCH <sub>3</sub> |    | 238-239 | 91       | C <sub>23</sub> H <sub>21</sub> O <sub>4</sub> N <sub>3</sub> | 68.47 | 5.25     | 10.42 |  | 68.94    | 5.21 | 10.60 |

TABLE 8. INFRARED SPECTRAL DATA<sup>a)</sup> FOR THE COMPOUNDS IV

|     | $\nu$ NH <sub>2</sub> | $\nu$ C=O  | $\delta$ NH <sub>2</sub> | Aromatic ring                | $\delta$ CH   |
|-----|-----------------------|------------|--------------------------|------------------------------|---------------|
| IVa | 3480, 3360            | 1705       | 1630                     | 1615, 1595, 1555, 1535       | 765, 700      |
| b   | 3430, 3330            | 1695       | 1630                     | 1615, 1605, 1580, 1550, 1540 | 825, 770      |
| c   | 3400, 3260            | 1710       |                          | 1615, 1600, 1555, 1540       | 770, 765      |
| d   | 3430, 3330            | 1695       | 1625                     | 1615, 1600, 1580, 1540       | 830, 775      |
| e   | 3430, 3330            | 1695       | 1625                     | 1610, 1575, 1550, 1535       | 825, 775, 770 |
| f   | 3420, 3290, 3200      | 1710, 1680 | 1640                     | 1615, 1600, 1550, 1530       | 825, 775      |
| g   | 3420, 3345            | 1700       | 1625                     | 1615, 1600, 1565, 1545       | 820, 775      |
| h   | 3450, 3340            | 1705       | 1625                     | 1610, 1600, 1555, 1545       | 770, 750, 710 |
| i   | 3470, 3410, 3290      | 1690       | 1625                     | 1615, 1605, 1555, 1540, 1520 | 825, 765      |
| j   | 3420, 3290            | 1705       | 1630                     | 1615, 1605, 1590, 1555, 1540 | 820, 770      |
| k   | 3420, 3300            | 1700       | 1630                     | 1615, 1605, 1585, 1555, 1540 | 815, 770      |
| l   | 3480, 3380            | 1705       | 1625                     | 1610, 1590, 1560, 1535,      | 770           |
| m   | 3510, 3370            | 1700       |                          | 1615, 1595, 1555, 1535       | 785, 740      |
| n   | 3450, 3340            | 1715       | 1630                     | 1615, 1565, 1540             | 825, 795      |
| o   | 3470, 3350            | 1715       | 1620                     | 1610, 1565, 1555, 1535       | 830, 800      |
| p   | 3430, 3350            | 1715       | 1620                     | 1610, 1565, 1555, 1535       | 790, 750      |
| q   | 3460, 3330            | 1695       | 1630                     | 1615, 1605, 1570, 1540       | 790, 735      |
| r   | 3420, 3290            | 1705       | 1635                     | 1600, 1570, 1540, 1525       | 795           |
| s   | 3430, 3310            | 1700       | 1625                     | 1615, 1600, 1585, 1565, 1535 | 790           |

a) All spectra were taken in potassium bromide disks; cm<sup>-1</sup>

TABLE 9. INFRARED SPECTRAL DATA FOR THE COMPOUNDS V

|    | $\nu$ NH   | $\nu$ O=C=O, NH-C=O | Aromatic ring                      | $\delta$ CH   |
|----|------------|---------------------|------------------------------------|---------------|
| Va | 3250       | 1700, 1685          | 1610, 1600, 1590, 1570, 1550       | 765, 700      |
| b  | 3250       | 1680                | 1610, 1600, 1590, 1570, 1545, 1520 | 825, 780      |
| c  | 3360, 3250 | 1700, 1680          | 1615, 1600, 1580, 1550, 1530       | 830, 775      |
| d  | 3250       | 1700, 1690          | 1610, 1590, 1575, 1550, 1535       | 830, 780, 770 |
| f  | 3250       | 1695, 1685          | 1615, 1605, 1590, 1580, 1555       | 795, 750      |
| g  | 3250       | 1690                | 1610, 1590, 1580, 1555, 1530       | 805, 745, 705 |

TABLE 10. NMR SPECTRAL DATA<sup>a)</sup> FOR THE COMPOUNDS IV AND V

|     | CH <sub>3</sub>                             | OCH <sub>3</sub> | CH <sub>2</sub> | -CH-            | Ring -CH=               | NH                     |
|-----|---------------------------------------------|------------------|-----------------|-----------------|-------------------------|------------------------|
| IVa |                                             |                  |                 |                 | 7.55-8.9(m, 10H)        | 9.75(br, 1H)           |
| c   |                                             | 4.4 (s, 3H)      |                 |                 | 7.2-8.3, 8.7-8.9(m, 9H) | 9.48(br, 1H)           |
| d   |                                             | 4.05(s, 3H)      |                 |                 | 7.15-8.7(m, 9H)         | 9.6 (br, 1H)           |
| e   | 1.35(d, 6H)                                 |                  |                 | 2.75-3.4(m, 1H) | 7.5-8.8(m, 8H)          | 8.5, 9.6(s, 1H each)   |
| f   | 2.52(s, 3H)                                 |                  |                 |                 | 7.6-8.8(m, 9H)          | 9.3, 9.7(s, 1H each)   |
| g   | 3.6 (s, 6H)                                 |                  |                 |                 | 7.6-8.1, 8.5-8.8(m, 9H) | 9.8 (br, 1H)           |
| j   |                                             | 4.12(d, 6H)      |                 |                 | 7.2-8.8(m, 8H)          | 9.65(br, 1H)           |
| k   | 1.55(t, 6H)                                 | 4.4 (q, 4H)      |                 |                 | 7.1-8.8(m, 8H)          | 9.6 (br, 1H)           |
| m   |                                             | 4.1 (s, 3H)      |                 |                 | 7.4-8.3(m, 8H)          | 8.45, 9.5(br, 1H each) |
| o   | 1.35(d, 6H)                                 | 4.1 (s, 3H)      |                 | 2.7-3.4(m, 1H)  | 7.4-7.8, 8-8.4(m, 7H)   | 8.45, 9.6(br, 1H each) |
| r   |                                             | 4.1 (s, 9H)      |                 |                 | 7.2-8.2(m, 6H)          | 8.6, 9.6(br, 1H each)  |
| s   | 1.6 (t, 6H)                                 | 4.15(s, 3H)      | 4.45(q, 4H)     |                 | 7.1-8.7(m, 7H)          | 9.65(br, 1H)           |
| Va  | 2.73(s, 3H, CO-CH <sub>3</sub> )            |                  |                 |                 | 7.5-8.95(m, 9H)         | 12.22(br, 1H)          |
| e   | 1.4(d, 6H) 2.7 (s, 3H, CO-CH <sub>3</sub> ) |                  |                 | 2.8-3.4(m, 1H)  | 7.5-9(m, 8H)            | 12.25(br, 1H)          |
| f   | 2.74(s, 3H, CO-CH <sub>3</sub> )            | 4.17(s, 3H)      |                 |                 | 7.6-8.6(m, 8H)          | 12.27(br, 1H)          |
| h   | 1.4(d, 6H) 2.7 (s, 3H, CO-CH <sub>3</sub> ) | 4.1 (s, 3H)      |                 | 2.8-3.4(m, 1H)  | 7.5-8.6(m, 7H)          | 12.2 (br, 1H)          |

a) Parts per million downfield from tetramethylsilane in CF<sub>3</sub>CO<sub>2</sub>H; s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, br=broad

### Experimental

All the melting points are uncorrected. The infrared spectra were determined by means of potassium bromide disks. The NMR spectra were determined in trifluoroacetic acid at 60 Mc, using tetramethylsilane as the internal standard. The chemical shifts are reported as parts per million downfield from TMS.

**Reaction of Ethyl Cyanoacetate and Salicylaldehyde or 3-Methoxysalicylaldehyde with Aldehydes.** A mixture of ethyl cyanoacetate (0.03 mol), salicylaldehyde or 3-methoxysalicylaldehyde (0.03 mol), aldehyde (0.03 mol), and ammonium acetate (0.04 mol) in ethanol (10 ml) was refluxed for 0.5-1.5 hr. A pale yellow crystalline matter precipitated out during the reaction. The experimental results and spectral data are summarized in Tables 1,3,5,6,8, and 10.

**Reaction of 3-Amidinocoumarin (1) and Benzaldehyde.** A mixture of 1 (2.82 g), benzaldehyde (1.59 g), and ammonium acetate (1.54 g) in ethanol (5 ml) was refluxed for 0.5 hr. A crystalline precipitate was formed during the reaction; this was collected, and washed with ethanol and then with water. Recrystallization from pyridine afforded 0.9 g of pale yellow crystals; mp 265-267°C. By their infrared

spectra and the results of elemental analyses, this compound was identified with IVa.

**Reaction of 3-Cyanocoumarin and Benzaldehyde.** A mixture of 3-cyanocoumarin (2.55 g), benzaldehyde (1.59 g), and ammonium acetate (1.54 g) in pyridine (5 ml) was heated for 1 hr. The pale yellow crystals which precipitated were then recrystallized from pyridine to give 0.8 g of almost white crystals; mp 262-264°C. This compound was proved to be identical with IVa by a study of their infrared spectra.

**Reaction of the II and IV Compounds and Acetic Anhydride.** To a solution of II or IV (0.001 mol) dissolved in pyridine (3-5 ml), acetic anhydride (5-8 ml) was added, the mixture was then refluxed for 2-4 hr. After standing overnight at room temperature, the crystalline precipitate thus formed was washed with dilute methanol. The experimental results and spectral data are summarized in Tables 2,4,7,9, and 10.

The authors wish to express their thanks to Dr. Taro Hayashi and Dr. Tatsuo Takeshima for their kind advice. Thanks are also due to Dr. Haruo Homma and his staff for their microanalyses, to Mr. Jun Uzawa for his measurements of the NMR spectra, and to Mr. Hironori Ogawa for his measurements of the IR spectra.