

Phosphorylation by Oxidation-Reduction Condensation.¹⁾ Preparation of Active Phosphorylating Reagents

Teruaki MUKAIYAMA and Mitsunori HASHIMOTO

Laboratory of Organic Chemistry, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo


(Received April 30, 1971)

Active phosphorylating reagents such as imidazolylphosphonate,^{2,3)} 2-pyridyl ester of phosphoric acid,^{4,5)} and phosphoramidate,⁶⁾ especially phosphoromorpholidate,⁷⁾ are considered to be important intermediates in nucleotide synthesis. In the present experiment, the preparation of these active phosphorylating reagents such as imidazolylphosphonate, 2-pyridyl ester of phosphoric acid and phosphoromorpholidate by the oxidation-reduction condensation reaction¹⁾ was studied.

First, the preparation of imidazolylphosphonate was attempted. In a typical reaction, triphenylphosphine (2 mmol) was added rapidly into a mixture of *p*-chlorophenyl dihydrogen phosphate (1 mmol), imidazole (5 mmol), triethylamine (1 mmol) and 2,2'-dipyridyl disulfide (2 mmol) in 5 ml of THF at room temperature. After stirring for 20 min, acetone solution of sodium iodide (1.5 mmol) was added to the reaction mixture and cooled down to 0°C. After being kept standing for 1 hr at 0°C, the precipitated white needle crystals were filtered off and washed with cold acetone. After removal of the solvent *in vacuo*, the sodium salt of *p*-chlorophenyl imidazolylphosphonate was obtained in a quantitative yield. UV: $\lambda_{\max}$ 270 m μ .

Found: C, 38.50; H, 2.51; N, 10.00%. Calcd for C₉H₇O₃N₂PClNa: C, 38.53; H, 2.52; N, 9.99%.

In the case of adenosine 5'-monophosphate, DMF was used in place of THF in the above experiment, and the corresponding adenosine 5'-imidazolylphosphonate was obtained in a quantitative yield without producing symmetrical pyrophosphate (AppA). *p*-Chlorophenyl-2-pyridyl phosphate was isolated as its cyclohexylammonium salt from *p*-chlorophenyl dihydrogen phosphate and 2-hydroxypyridine in high yield according to the following procedure. *p*-Chlorophenyl dihydrogen phosphate (1 mmol) was allowed to react with 2-hydroxypyridine (5 mmol), triphenylphosphine (2 mmol) and 2,2'-dipyridyl disulfide (2 mmol) in THF at room temperature for 3 hr. Cyclohexylamine (1.5 mmol) was then added to the reaction



mixture cooled in an ice-bath and kept standing for 2 hr. The precipitated crystals were filtered off, washed with cold THF and dried *in vacuo*. The corresponding cyclohexylammonium salt of *p*-chlorophenyl-2-pyridyl phosphate, mp 164°C, was obtained in 80% yield. UV: $\lambda_{\max}$ 262 m μ .⁸⁾ Found: C, 52.95; H, 5.85; N, 7.40%. Calcd for C₁₇H₂₂O₄N₂ClP: C, 53.10; H, 5.76; N, 7.29%.

Next, it was found that adenosine 5'-phosphoromorpholidate, an important intermediate in the synthesis of pyrophosphates such as FAD (flavin Adenine dinucleotide), was obtained in a quantitative yield by this method. Triphenylphosphine (2 mmol) was added into the suspended DMF solution of adenosine monophosphate (1 mmol), morpholine (5 mmol), triethylamine (1 mmol) and 2,2'-dipyridyl disulfide (2 mmol) at room temperature. After the mixture was stirred for 1 hr, the suspended solution turned clear. By this time, paper chromatography and paper electrophoresis showed the presence of only one spot corresponding to adenosine 5'-phosphoromorpholidate. Determination by UV absorption after separation with paper chromatography showed that the adenosine 5'-phosphoromorpholidate was obtained in a quantitative yield. $\lambda_{\max}^{\text{pH } 7}$ 260 m μ (ϵ 1.59 $\times$ 10⁴). R_f 0.41, isopropanol - concd. ammonium hydroxide - water (7 : 1 : 2).

In conclusion, it is noted that the present oxidation-reduction condensation produces active phosphorylating reagents such as imidazolylphosphonate, 2-pyridyl ester of phosphoric acid and phosphoromorpholidate in high yields in a short reaction time by simply mixing triphenylphosphine, 2,2'-dipyridyl disulfide and phosphate with imidazole, 2-hydroxypyridine or morpholine, respectively.

The authors wish to express their hearty thanks to Dr. Tsujiaki Hata for his advice.

1) T. Mukaiyama and M. Hashimoto, This Bulletin, **44**, 196 (1971).

2) R. Cramer, H. Schaller, and H. A. Staab, *Chem. Ber.*, **94**, 1612 (1961).

3) R. Cramer and H. Neunhoeffer, *ibid.*, **95**, 1664 (1962).

4) K. H. Scheit and W. Kampe, *ibid.*, **98**, 1045 (1965).

5) W. Kampe, *ibid.*, **99**, 593 (1966).

6) J. G. Moffatt and H. G. Khorana, *J. Amer. Chem. Soc.*, **80**, 3756 (1958).

7) J. G. Moffatt and H. G. Khorana, *ibid.*, **83**, 663 (1961).

8) W. Kampe, *Chem. Ber.*, **98**, 1031 (1965).