

a) Molar ratio. b) The structure could not be determined. c) Yield of IV.

Scheme 1. Conversion of TDZ to 1,3,5-triazine.

reduced by the effect of the protonated imino group, and so the attack of the amine may be easier (Fig. 1).

Experimental

2-Methoxycarbonimidoyl-3-imino-5-ethoxy- Δ^4 -1,2,4-thiadiazoline (I). To a stirred mixture of III⁴⁾ (6.27 g, 40 mmol) and acetonitrile (40 ml), we added a solution of *N*-chloro-*O*-methylisourea (4.34 g, 40 mmol) in acetonitrile (20 ml). The temperature was maintained below 5 °C during the reaction. After 1.75 h of continued stirring, the reaction mixture was evaporated to dryness under reduced pressure. The title compound was obtained by washing the residue with water 6.45 g; 80%; mp 138–143 °C. Recrystallization from ethanol gave a pure product; mp 143 °C. Picrate: mp 137 °C (from water). Found: C, 33.32; H, 2.97; N, 22.78%. Calcd for C₁₂H₁₃N₇SO₈: C, 33.42; H, 3.04; N, 22.73%.

2-Amidino-3-imino-5-ethoxy- Δ^4 -1,2,4-thiadiazoline (IIa). The title compound was prepared according to approximately the same procedure as that described previously.³⁾ Recrystallization from methanol–water gave pure IIa; mp 177–180 °C. Picrate: mp 288 °C (from water). Found: C, 31.77; H, 2.93; N, 26.80%. Calcd for C₁₁H₁₂N₈SO₈: C, 31.74; H, 2.91; N, 26.92%.

Amidination of I. **Amidination of I with Benzylamine:** A solution of I (2.02 g, 10 mmol), benzylamine (2.14 g, 20 mmol), and benzylamine hydrochloride (1.44 g, 10 mmol) in ethanol (25 ml) was refluxed for 3 h. After the reaction mixture had been evaporated to dryness under reduced pressure, the residue was washed with ether (10 ml). TDZ, IIc containing sulfur was obtained by washing the residue with water: 1.39 g; 50%; mp 137–140 °C (from MeOH). The addition of petroleum ether to the ethereal washings afforded 2-amino-4-benzylamino-6-ethoxy-1,3,5-triazine (IVc) (0.47 g; 19%; mp 181–183 °C (from DMF). Found: C, 58.61; H, 6.26; N, 28.45%. Calcd for C₁₂H₁₅N₅O: C, 58.76; H, 6.16; N, 28.55%. Picrate: mp 174–175 °C (from EtOH). Found: C, 45.45; H, 3.83; N, 23.51%. Calcd for C₁₈H₁₈N₈O₈: C, 45.57; H, 3.82; N, 23.62%.

Amidination of I with *n*-Propylamine: Using the same procedure as above (20 mmol scale), after the sulfur had been removed, the filtrate was concentrated. The residual oily material was crystallized from water (20 ml), subsequent recrystallization from ethanol (15 ml) afforded two kinds of products. IIb: 0.92 g; 20%; mp 162–163 °C (from EtOH). 2-Amino-4-ethoxy-6-*n*-propylamino-1,3,5-triazine (IVb): 1.53

g; 39%; mp 132–134 °C (from aq MeOH). Found: C, 48.59; H, 7.45; N, 35.31%. Calcd for C₈H₁₅N₅O: C, 48.72; H, 7.67; N, 35.51%.

2-Amino-4-ethoxy-6-methylamino-1,3,5-triazine (IVd) was obtained in a similar manner by the reaction of I with methylamine: 10%; mp 171–172 °C (170–171 °C)⁶⁾ (from water). Picrate: dp 202–203 °C (from water). Found: C, 36.26; H, 3.74; N, 27.70%. Calcd for C₁₂H₁₄N₈O₈: C, 36.19; H, 3.54; N, 28.13%.

2-Carbamoyl-3-imino-5-ethoxy- Δ^4 -1,2,4-thiadiazoline (V). A slow stream of dry hydrogen chloride was passed through a solution of I (2.02 g, 10 mmol) in methanol (30 ml) at 50 °C for 40 min. A small amount of a brown precipitate was removed by filtration, and the filtrate was evaporated to dryness under reduced pressure. The title compound was obtained by washing the residue with ether; 1.84 g, 98%; dp > 300 °C (from DMF–EtOH).

Reaction of TDZ with Free Amines. **Reaction of I with Aqueous Ammonia:** A solution of I (1.01 g, 5 mmol) and 5% ammonium hydroxide (8 ml, 20 mmol) in ethanol (15 ml) was refluxed for 3 h. After the precipitated sulfur had been filtered off, the filtrate was cooled and I was precipitated; 0.46 g, 46%. After the filtration of I, the concentration of the filtrate afforded 2-amino-4-ethoxy-6-methoxy-1,3,5-triazine (IVe): 0.46 g, 54%; mp 102–103 °C. Found: C, 41.92; H, 6.03; N, 32.62%. Calcd for C₆H₁₀N₄O₂: C, 42.35; H, 5.92; N, 32.92%.

Reaction of IIc with Benzylamine: A mixture of IIb (0.55 g, 2 mmol), benzylamine (0.72 g, 4 mmol), and benzylamine hydrochloride (0.29 g, 2 mmol) was refluxed in ethanol (6 ml) for 3 h. After the reaction mixture had been concentrated, the residue was washed with ether and water; 1,3,5-triazine (IVc) was thus obtained; 0.49 g; quant.

References

- 1) Part 104 of "Studies of Cyanamide Derivatives."
- 2) T. Fuchigami and K. Odo, *Chem. Lett.*, **1973**, 917.
- 3) T. Fuchigami and K. Odo, *Bull. Chem. Soc. Jpn.*, **48**, 310 (1975).
- 4) T. Fuchigami and K. Odo, *Bull. Chem. Soc. Jpn.*, **49**, 3165 (1976).
- 5) J. Controulis and C. Banks, *J. Am. Chem. Soc.*, **67**, 1946 (1945).
- 6) W. Pearlmann, J. Mitsulski, and C. Banks, *J. Am. Chem. Soc.*, **71**, 3248 (1949).